Analysis of coding and non-coding transcriptome of peripheral B cells reveals an altered interferon response factor (IRF)-1 pathway in multiple sclerosis patients

Viviana Annibali, Renato Umeton, Antonia Palermo, Martina Severa, Marilena Paola Etna, Simona Giglio, Silvia Romano, Michela Ferraldeschi, Maria Chiara Buscarinu, Andrea Vecchione, Anita Annese, Claudia Policano, Rosella Mechelli, Raffaella Pizzolato Umeton, Arianna Fornasiero, Daniela Francesca Angelini, Gisella Guerrera, Luca Battistini, Eliana Marina Coccia, Marco Salvetti, Giovanni Ristori

PII: S0165-5728(18)30216-9
DOI: doi:10.1016/j.jneuroim.2018.09.005
Reference: JNI 476843

To appear in: *Journal of Neuroimmunology*

Received date: 2 May 2018
Revised date: 13 August 2018
Accepted date: 10 September 2018

Please cite this article as: Viviana Annibali, Renato Umeton, Antonia Palermo, Martina Severa, Marilena Paola Etna, Simona Giglio, Silvia Romano, Michela Ferraldeschi, Maria Chiara Buscarinu, Andrea Vecchione, Anita Annese, Claudia Policano, Rosella Mechelli, Raffaella Pizzolato Umeton, Arianna Fornasiero, Daniela Francesca Angelini, Gisella Guerrera, Luca Battistini, Eliana Marina Coccia, Marco Salvetti, Giovanni Ristori , Analysis of coding and non-coding transcriptome of peripheral B cells reveals an altered interferon response factor (IRF)-1 pathway in multiple sclerosis patients. Jni (2018), doi:10.1016/j.jneuroim.2018.09.005

Analysis of coding and non-coding transcriptome of peripheral B cells reveals an altered interferon response factor (IRF)-1 pathway in Multiple Sclerosis patients.

Viviana Annibali[1+], Renato Umeton[2,9+], Antonia Palermo[3], Martina Severa[4], Marilena Paola Etna[4], Simona Giglio[5], Silvia Romano[1], Michela Ferraldeschi[1], Maria Chiara Buscarinu[1], Andrea Vecchione[5], Anita Annese[1], Claudia Policano[1], Rosella Mechelli[1], Raffaella Pizzolato Umeton[6], Arianna Fornasiero[1], Daniela Francesca Angelini[7],Gisella Guerrera[7], Luca Battistini[7], Eliana Marina Coccia[4], Marco Salvetti[1,8*] and Giovanni Ristori[1*]

[1]Center for Experimental Neurological Therapies, Sant'Andrea Hospital, Department of Neurosciences, Mental Health and Sensory Organs (NESMOS), Faculty of Medicine and Psychology, Sapienza University of Rome, Rome, Italy.

[2]Department of Informatics, Dana-Farber Cancer Institute, Boston, MA, United States.

[3]Department of Mathematics and Computer Science, University of Calabria.

[4]Department of Infectious, Parasitic and Immune-mediated Disease, Istituto Superiore di Sanità, Rome, Italy.

[5]Division of Pathology, Department of Clinical and Molecular Medicine, Sant'Andrea Hospital, Faculty of Medicine and Psychology, Sapienza University of Rome, Rome, Italy.

[6]Department of Neurology, University of Massachusetts Memorial Medical Center, Worcester, MA, USA.

[7]Neuroimmunology Unit, Fondazione Santa Lucia, Rome, Italy.

[8]IRCCS Istituto Neurologico Mediterraneo (INM) Neuromed (M.S.), Pozzilli, IS, Italy

[9]Massachusetts Institute of Technology, Cambridge, MA, United States.

[+]Contributed equally to this work: Viviana Annibali, Renato Umeton

[*]Corresponding authors: Giovanni Ristori: giovanni.ristori@uniroma1.it; Telephone: +39 06 33776044; Marco Salvetti: marco.salvetti@uniroma1.it; Telephone: +390633775994: Fax: +390633775900; Sant'Andrea Hospital, Via di Grottarossa 1035-1039, 00189 Rome, Italy.