



## Molecular confirmation of *Hymenolepis hibernia* in field mice (*Apodemus sylvaticus*) from St Kilda has potential to resolve a host-parasite relationship

Neil Sargison<sup>a,\*</sup>, Jeremy Herman<sup>b</sup>, Jill Pilkington<sup>c</sup>, Peter Buckland<sup>a</sup>, Kathryn Watt<sup>c</sup>, Alex Chambers<sup>a,c,1</sup>, Umer Chaudhry<sup>a</sup>

<sup>a</sup> University of Edinburgh, Royal (Dick) School of Veterinary Studies and Roslin Institute, Easter Bush Veterinary Centre, Roslin, Midlothian, EH25 9RG, UK

<sup>b</sup> National Museums of Scotland, Natural Sciences Department, Chambers Street, Edinburgh, EH1 1JF, UK

<sup>c</sup> University of Edinburgh, Institute of Evolutionary Biology, Ashworth Laboratories, Kings Buildings, West Mains Road, Edinburgh, EH9 3JT, UK

### ARTICLE INFO

#### Keywords:

*Hymenolepis hibernia* cyclophyllidean tapeworm

*Apodemus sylvaticus* field mouse

Host parasite relationship

Maximum likelihood tree

Phylogeographic variation

### ABSTRACT

Insular wildlife populations provide opportunities to examine biological questions in systems that are relatively closed and potentially tractable, striking examples being the long-term studies of ecology and evolution in the red deer and feral sheep populations on the Hebridean islands of Rum and St Kilda. In the case of parasitology, understanding of parasitic infections insular wildlife populations in conjunction with knowledge of their origins has the potential to add a fresh perspective to disease control in humans and domestic animals. In the case of parasitology, understanding infections of insular wildlife populations, in conjunction with knowledge of their origins, has the potential to add a fresh perspective to disease control in humans and domestic animals. With this in mind, gross and molecular examination for the presence of cyclophyllidean tapeworms was performed on the viscera and rectal contents of 17 preserved specimens of *Apodemus sylvaticus* field mice and on the naturally voided faeces of a further four mice on the remote archipelago of St Kilda. Molecular speciation of hexacanth embryos extracted from the faeces of two mice, using nucleotide sequence analysis of the ribosomal cytochrome c-oxidase subunit-1, confirmed infection with *Hymenolepis hibernia*. Phylogenetic analysis showed that these were genetically distinct from *Hymenolepis diminuta*, previously reported in the insular *A. sylvaticus* mice, and from other published *H. hibernia* haplotypes. There was insufficient hymenolepidid tapeworm phylogeographic variation to resolve the origins of the co-evolved St Kilda mice, primarily due to a lack of published *H. hibernia* Cox-1 sequence data across the parasite's geographical range. Nevertheless, the Maximum Likelihood haplotype tree shows the potential for molecular parasitology to resolve a host-parasite relationship once more data become available. Morphological diagnostic features of zoonotic *H. hibernia* eggs are also described.

### 1. Introduction

St Kilda is an uninhabited archipelago lying in the north Atlantic Ocean about 40 miles west of the Outer Hebrides, and comprising the islands of Soay, Hirta, Dun and Boreray. Hirta, the largest of these islands, with about 670 ha of exposed hill and moorland, was continuously inhabited for thousands of years. The islanders kept sheep and cattle to supplement their diet of seabirds and cereal crops and for wool to trade and pay the landlord, along with large numbers of dogs. In 1930 the human population of St Kilda was evacuated along with their animals. In 1932, 107 feral Soay sheep were relocated onto Hirta

from the smaller neighbouring island of Soay, to maintain the pasture. These sheep have been left unmanaged since then, and the only other terrestrial mammals present on St Kilda have been mice and visiting humans.

Two years after the 1930 evacuation of St Kilda, the resident population of house mice (*Mus musculus*) became extinct (Harrisson and Moy-Thomas, 1933). Without the human inhabitants, these commensal animals were unable to compete with the less strongly anthropophilous field mice (*Apodemus sylvaticus*), which have survived on the islands of Hirta and Dun (Morton-Boyd, 1956; Berry, 1969; Berry and Tricker, 1969). Neither of these species would have survived the Last Glacial

\* Corresponding author.

E-mail addresses: [neil.sargison@ed.ac.uk](mailto:neil.sargison@ed.ac.uk) (N. Sargison), [J.Herman@nms.ac.uk](mailto:J.Herman@nms.ac.uk) (J. Herman), [J.Pilkington@ed.ac.uk](mailto:J.Pilkington@ed.ac.uk) (J. Pilkington), [buckland05@hotmail.com](mailto:buckland05@hotmail.com) (P. Buckland), [Kathryn.Watt@ed.ac.uk](mailto:Kathryn.Watt@ed.ac.uk) (K. Watt), [Alex.Chambers@agresearch.co.nz](mailto:Alex.Chambers@agresearch.co.nz) (A. Chambers), [uchaudhr@exseed.ed.ac.uk](mailto:uchaudhr@exseed.ed.ac.uk) (U. Chaudhry).

<sup>1</sup> Current address. AgResearch, Tennent Drive, Palmerston North 4410, New Zealand.

<https://doi.org/10.1016/j.ijppaw.2018.09.007>

Received 18 July 2018; Received in revised form 22 September 2018; Accepted 28 September 2018

2213-2244/ © 2018 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Maximum about 24 thousand years ago (Johnsen et al., 1992) in this location, hence both would have been introduced more recently, and presumably inadvertently, by humans (Corbet, 1961). The source of these animals has been a matter of considerable interest to zoologists and conservationists, with some evidence for introductions from Scandinavia that have been attributed to Vikings. However there may have been earlier and later introductions from mainland Britain or elsewhere (Berry, 1969; Jones et al., 2010; Herman et al., 2017). At various times, both the St Kilda house mouse and the field mouse have been considered as distinct species or subspecies, due to their size and pelage colour, however genetic evidence does not support any substantial differentiation from populations elsewhere in the British Isles or northern and western Europe (Jones et al., 2010; Herman et al., 2017) and this distinction consequently seems unwarranted. The St Kilda field mice survive on seeds, invertebrates and scavenged carcasses of sheep and seabirds (Berry, 1969). Previous parasitological studies have found that they are infected with adult *Hymenolepis diminuta* (Morton-Boyd, 1959) and larval metacestodes of *Taenia taeniaeformis* (referred to as *Taenia crassicolis* or *cysticercus fasciolaris*) (Waterston, 1906; Morton-Boyd, 1959).

The present study was undertaken to investigate the possibility that the St Kilda field mouse might be a definitive host for *Taenia hydatigena*, accounting for the high prevalence of cysticerci tenuicollis previously identified in the St Kilda Soay sheep (Torgerson et al., 1995), but this highly speculative hypothesis was not supported. Nevertheless, the study provided an opportunity for the post-mortem examination of the intestinal tracts from a series of preserved specimens of St Kilda field mice, showing the presence of adult cyclophyllidean tapeworms. In addition, examination of mouse faeces revealed the presence of taeniid eggs and permitted molecular confirmation of their species identity. The serendipitous and novel identification of *Hymenolepis hibernia* provided an opportunity to investigate the phylogenetic origins of the parasite, with reference to those of its wildlife host and to consideration of its zoonotic potential (Nkouawa et al., 2016).

## 2. Materials and methods

### 2.1. Mouse samples used to identify adult tapeworm infections

Viscera from carcasses of 17 St Kilda field mice, preserved in the vertebrate collections of the National Museums of Scotland, were examined. These had been collected between June 2011 and June 2012 from locations throughout Hirta and kept frozen at  $-20^{\circ}\text{C}$ , albeit with occasional cycles of thawing and re-freezing. Any tapeworms or tapeworm segments that were found were recovered and transferred to 70% ethanol. The rectal contents were collected for coprological examination. Naturally-voided faecal samples were also obtained opportunistically from an additional four locations in the Village Bay area on the island of Hirta.

### 2.2. Parasitological examination of faeces

Faecal samples were suspended in water. Sedimented material was then processed through a series of sieves. Material deposited on 53  $\mu\text{m}$  and 30  $\mu\text{m}$  sieves was retained and examined microscopically for the presence of cyclophyllidean cestode eggs. Eggs were picked and transferred to 70% ethanol.

### 2.3. Genomic DNA isolation, PCR amplification and sequence analysis of mitochondrial cytochrome c oxidase subunit-1 (Cox-1)

Aliquots of about 100 cyclophyllidean cestode eggs with hexacanth onchospheres were lysed in single 0.2 ml tubes containing 50  $\mu\text{l}$  of proteinase K lysis buffer and stored at  $-80^{\circ}\text{C}$  as previously described (Redman et al., 2008). 1  $\mu\text{l}$  of a 1:5 dilution of neat lysate was used as PCR template and the same dilutions of lysate buffer made in parallel

were used as negative controls. A fragment of 396 bp of the mitochondrial cytochrome c oxidase subunit-1 (Cox-1), was amplified using forward (CeCox-For- TTTTGGGCATCCTGAGGTTTAT) and reverse primers (CeCox-Rev- TAAAGAAAGAACATAATGAAAATG) (Lavikainen et al., 2008). PCR reaction conditions consisted of 1x thermopol reaction buffer (NEB BioLab), 2 mM  $\text{MgSO}_4$ , 200  $\mu\text{M}$  dNTPs, 0.2  $\mu\text{M}$  forward and reverse primers and 1U of Phusion high fidelity DNA polymerase (Finenzyme). The thermo-cycling parameters consisted of an initial 98  $^{\circ}\text{C}$  for 30 s followed by 40 cycles of 98  $^{\circ}\text{C}$  for 10 s, 54  $^{\circ}\text{C}$  for 30 s and 72  $^{\circ}\text{C}$  for 2 min with a single final extension cycle of 72  $^{\circ}\text{C}$  for 7 min. DNA templates for direct sequencing of the Cox-1 region were cleaned using QIAquick PCR Purification Kit (Cat No./ID: 28104) following the manufacturers' protocols. Amplicons were sequenced from both ends using the same primers used for PCR amplification for each region with BigDye Terminator Cycle Sequencing (Applied Biosystems). Sequences of Cox-1 were edited to remove primers and poor quality sequence on both ends using Geneious Pro 5.4 software (Kearse et al., 2012). Sequences showing 100% base pair similarity were grouped into haplotypes using the CD-HIT Suite software (Huang et al., 2010). Haplotype and GenBank reference sequences for *Hymenolepis* spp. were imported into MEGA 6 (Tamura et al., 2013) and a Maximum Likelihood tree was inferred, using the Hasegawa-Kishino-Yano (1985) nucleotide substitution model with gamma distribution of rates across sites (HKY + G), as selected with the Bayesian Information Criterion. Model parameters were estimated from the data and branch support was estimated from 1000 bootstrap pseudoreplicates of the data. The Maximum Likelihood tree was rooted using a closely related outgroup sequence from *H. contortus* (GenBank accession no. AF070785).

## 3. Results

### 3.1. Gross parasitological tapeworm identification

Cyclophyllidean tapeworms or segments were identified in the small intestines of five of the 17 mice that were examined postmortem. These were in poor condition and mostly fragmented, but could be separated into two morphological groups based on their segment dimensions and estimated overall length of about 2 mm wide and 6 cm long, or about 4 mm wide and up to 31 cm long. One of the smaller tapeworms had an intact, unarmed scolex (Fig. 1A), but the scolices were not found for any others. The most intact larger tapeworm, lacking a scolex, is shown in Fig. 1F. Cyclophyllidean tapeworm eggs were only recovered on the 53  $\mu\text{m}$  sieve, from the faeces of each of the five mice in which adult tapeworms were found and from one of the four naturally-voided faecal samples, collected from a cleat (or cleit; an ancient small stone storage hut). Eggs in the faeces of three mice (numbered 5, 8 and 10) in which the smaller tapeworms were identified and in the faeces collected from a cleat (labelled X) were typically anoplocephalid, being approximately pyramidal-shaped, with a maximum width of 62–68  $\mu\text{m}$  (Fig. 1B and D) and containing a similarly shaped onchosphere, about 38  $\mu\text{m}$  across (Fig. 1E). Eggs in the faeces of the three mice (numbered 7, 8 and 9) and in the naturally-voided faeces (lettered X) were near spherical, mean 68  $\mu\text{m}$  in diameter, with a thick, dark outer shell, a granular and non-striated embryophore about 20  $\mu\text{m}$  deep, granular, and a thin onchospherical membrane (Fig. 1C, G, I and 1K) surrounding a 36–38  $\mu\text{m}$  spherical hexacanth onchosphere (Fig. 1H and J and 1L). The embryonic hooks were relatively long, and consistently arranged in offset, radiating pairs. Cyclophyllidean co-infections were, therefore, identified in mouse 8 and in the naturally-voided faeces from mouse X. However, the lack of non-overlapping and definitive morphological diagnostic characters for cyclophyllidean eggs and the poor condition of the adult tapeworms prevented identification to generic or specific level.

Download English Version:

<https://daneshyari.com/en/article/11007629>

Download Persian Version:

<https://daneshyari.com/article/11007629>

[Daneshyari.com](https://daneshyari.com)