

Lesser lower extremity mechanical loading associates with a greater increase in serum cartilage oligomeric matrix protein following walking in individuals with anterior cruciate ligament reconstruction

Brittney A. Luc-Harkey, Jason R. Franz, Anthony C. Hackney, J. Troy Blackburn, Darin A. Padua, Brian Pietrosimone

PII: S0268-0033(18)30100-1
DOI: doi:10.1016/j.clinbiomech.2018.09.024
Reference: JCLB 4613

To appear in: *Clinical Biomechanics*

Received date: 7 February 2018
Accepted date: 25 September 2018

Please cite this article as: Brittney A. Luc-Harkey, Jason R. Franz, Anthony C. Hackney, J. Troy Blackburn, Darin A. Padua, Brian Pietrosimone , Lesser lower extremity mechanical loading associates with a greater increase in serum cartilage oligomeric matrix protein following walking in individuals with anterior cruciate ligament reconstruction. Jclb (2018), doi:10.1016/j.clinbiomech.2018.09.024

Lesser Lower Extremity Mechanical Loading Associates with a Greater Increase in Serum Cartilage Oligomeric Matrix Protein following Walking in Individuals with Anterior Cruciate Ligament Reconstruction

Brittney A. Luc-Harkey[1]
Jason R. Franz[2]
Anthony C. Hackney[3]
J. Troy Blackburn[3]
Darin A. Padua[3]
Brian Pietrosimone[3]

[1]Neurological Clinical Research Institute, Department of Neurology, Massachusetts General Hospital, Boston, MA.[2]Joint Department of Biomedical Engineering, University of North Carolina at Chapel Hill and North Carolina State University, Chapel Hill, NC.
[3]Department of Exercise and Sports Science, University of North Carolina at Chapel Hill, Chapel Hill, NC.

Corresponding Author:
Brittney A. Luc-Harkey, PhD
Neurological Clinical Research Institute
Department of Neurology
Massachusetts General Hospital
165 Cambridge Street
Boston MA 02114
The United States of America
Email: bharkey@mgh.harvard.edu
Phone: 617.742.4703

Abstract Word Count: 250 words

Manuscript Word Count: 3,411 words