

## Accepted Manuscript

Title: Seroprevalence and risk factors of lumpy skin disease in Ethiopia

Authors: Wassie Molla, Klaas Frankena, Getachew Gari, Menbere Kidane, Dereje Shegu, Mart C.M. de Jong



PII: S0167-5877(18)30535-X  
DOI: <https://doi.org/10.1016/j.prevetmed.2018.09.029>  
Reference: PREVET 4555

To appear in: *PREVET*

Received date: 4-8-2018  
Revised date: 18-9-2018  
Accepted date: 29-9-2018

Please cite this article as: Molla W, Frankena K, Gari G, Kidane M, Shegu D, de Jong MCM, Seroprevalence and risk factors of lumpy skin disease in Ethiopia, *Preventive Veterinary Medicine* (2018), <https://doi.org/10.1016/j.prevetmed.2018.09.029>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Seroprevalence and risk factors of lumpy skin disease in Ethiopia

Wassie Molla<sup>a,\*</sup>, Klaas Frankena<sup>b</sup>, Getachew Gari<sup>c</sup>, Menbere Kidane<sup>c</sup>, Dereje Shegu<sup>c</sup>, Mart C.M. de Jong<sup>d</sup>

<sup>a</sup> College of Veterinary Medicine and Animal Sciences, University of Gondar, P.O. Box 196, Gondar, Ethiopia

<sup>b</sup> Adaptation Physiology, Wageningen University & Research, P. O. Box 338, 6700AH Wageningen, The Netherlands

<sup>c</sup> National Animal Health Diagnostic and Investigation Centre (NAHDIC), Sebeta, Ethiopia

<sup>d</sup> Quantitative Veterinary Epidemiology, Wageningen University & Research, P.O. Box 338, 6700 AH Wageningen, The Netherlands

\*Corresponding author. Tel.: +251 918030334

E-mail address: Mollawassie@yahoo.com

### Highlights

- LSD seroprevalence is high at animal level and very high at herd level.
- Cattle populations with many adult animals and that live in wet areas are at highest risk for LSD.
- Cattle in frequent contact with other animals and with other animal species have lower risk to be infected with LSDV.
- The estimated intracluster correlation coefficient for LSD equals 0.104.

Download English Version:

<https://daneshyari.com/en/article/11019409>

Download Persian Version:

<https://daneshyari.com/article/11019409>

[Daneshyari.com](https://daneshyari.com)