

Accepted Manuscript

A new method using 1,3,5-triazine as an umpolung hydrogen cyanide equivalent toward the syntheses of isoquinolinone and 2-pyridone derivatives

Joji Hayashida, Shinya Yoshida

PII: S0040-4039(18)31110-9
DOI: <https://doi.org/10.1016/j.tetlet.2018.09.029>
Reference: TETL 50268

To appear in: *Tetrahedron Letters*

Received Date: 1 August 2018
Revised Date: 6 September 2018
Accepted Date: 11 September 2018

Please cite this article as: Hayashida, J., Yoshida, S., A new method using 1,3,5-triazine as an umpolung hydrogen cyanide equivalent toward the syntheses of isoquinolinone and 2-pyridone derivatives, *Tetrahedron Letters* (2018), doi: <https://doi.org/10.1016/j.tetlet.2018.09.029>



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A new method using 1,3,5-triazine as an umpolung hydrogen cyanide equivalent toward the syntheses of isoquinolinone and 2-pyridone derivatives.

Joji Hayashida* and Shinya Yoshida

Process Chemistry Labs, Astellas Pharma Inc., 160-2 Akahama, Takahagi, Ibaraki 318-0001, Japan

*Corresponding authors:

Joji Hayashida: joji.hayashida@astellas.com

Download English Version:

<https://daneshyari.com/en/article/11020147>

Download Persian Version:

<https://daneshyari.com/article/11020147>

[Daneshyari.com](https://daneshyari.com)