

## Accepted Manuscript

Posttraumatic Stress Disorder Clusters and Suicidal Ideation

Lily A. Brown , Ateka Contractor , Kathy Benhamou

PII: S0165-1781(18)30859-X  
DOI: <https://doi.org/10.1016/j.psychres.2018.09.030>  
Reference: PSY 11744

To appear in: *Psychiatry Research*

Received date: 7 May 2018  
Revised date: 4 August 2018  
Accepted date: 14 September 2018

Please cite this article as: Lily A. Brown , Ateka Contractor , Kathy Benhamou , Posttraumatic Stress Disorder Clusters and Suicidal Ideation, *Psychiatry Research* (2018), doi: <https://doi.org/10.1016/j.psychres.2018.09.030>



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**HIGHLIGHTS**

- Suicidal ideation (SI) was positively correlated with all PTSD factors in both studies.
- Negative mood/cognitions and arousal symptom clusters were more strongly correlated with SI than other clusters in Study 1
- PTSD symptom clusters contribute to SI in community and treatment-seeking samples.
- Findings suggest a need to assess SI when PTSD cluster scores are elevated.
- Interventions may benefit from targeting NACM and AAR symptom severity.

Download English Version:

<https://daneshyari.com/en/article/11024082>

Download Persian Version:

<https://daneshyari.com/article/11024082>

[Daneshyari.com](https://daneshyari.com)