

## Accepted Manuscript

### Attention Deficit-Hyperactivity Disorder in Adult Bipolar Disorder Patients

Marco Pinna , Caterina Visioli , Carlo Mario Rago ,  
Mirko Manchia , Leonardo Tondo , Ross J. Baldessarini

PII: S0165-0327(18)31169-8  
DOI: <https://doi.org/10.1016/j.jad.2018.09.038>  
Reference: JAD 10120



To appear in: *Journal of Affective Disorders*

Received date: 28 May 2018  
Revised date: 17 August 2018  
Accepted date: 15 September 2018

Please cite this article as: Marco Pinna , Caterina Visioli , Carlo Mario Rago , Mirko Manchia , Leonardo Tondo , Ross J. Baldessarini , Attention Deficit-Hyperactivity Disorder in Adult Bipolar Disorder Patients, *Journal of Affective Disorders* (2018), doi: <https://doi.org/10.1016/j.jad.2018.09.038>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Highlights**

- We studied the rates of ADHD in 703 type-I and type-II BD patients
- The lifetime prevalence of ADHD in BD patients was of 24.6%
- BD patients with ADHD differ for many factors from those without
- BD patients with ADHD had several unfavorable clinical and functional outcomes
- Co-occurrence of bipolar disorder (BD) and attention deficit-hyperactivity disorder (ADHD) is not uncommon

Download English Version:

<https://daneshyari.com/en/article/11025327>

Download Persian Version:

<https://daneshyari.com/article/11025327>

[Daneshyari.com](https://daneshyari.com)