


Accepted Manuscript

Repeat-Dose Ketamine Augmentation for Treatment-Resistant Depression with Chronic Suicidal Ideation: A Randomized, Double Blind, Placebo Controlled Trial

Dawn F. Ionescu , Kate H. Bentley , Matthias Eikermann , Norman Taylor , Oluwaseun Akeju , Michaela B. Swee , Kara J. Pavone , Samuel R. Petrie , Christina Dording , David Mischoulon , Jonathan E. Alpert , Emery N. Brown , Lee Baer , Matthew K. Nock , Maurizio Fava , Cristina Cusin



PII: S0165-0327(18)30778-X
DOI: <https://doi.org/10.1016/j.jad.2018.09.037>
Reference: JAD 10119

To appear in: *Journal of Affective Disorders*

Received date: 18 April 2018
Revised date: 17 August 2018
Accepted date: 15 September 2018

Please cite this article as: Dawn F. Ionescu , Kate H. Bentley , Matthias Eikermann , Norman Taylor , Oluwaseun Akeju , Michaela B. Swee , Kara J. Pavone , Samuel R. Petrie , Christina Dording , David Mischoulon , Jonathan E. Alpert , Emery N. Brown , Lee Baer , Matthew K. Nock , Maurizio Fava , Cristina Cusin , Repeat-Dose Ketamine Augmentation for Treatment-Resistant Depression with Chronic Suicidal Ideation: A Randomized, Double Blind, Placebo Controlled Trial, *Journal of Affective Disorders* (2018), doi: <https://doi.org/10.1016/j.jad.2018.09.037>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Controlled studies of ketamine for treatment-resistant depression (TRD) are needed.
- We enrolled 26 outpatients with TRD and current, chronic suicidal ideation (SI).
- Ketamine did not produce significantly better effects on depression than placebo.
- Ketamine did not produce significantly better effects on SI than placebo.
- Severely ill, chronically depressed patients may need higher doses of ketamine.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/11025343>

Download Persian Version:

<https://daneshyari.com/article/11025343>

[Daneshyari.com](https://daneshyari.com)