

Accepted Manuscript

Employing exogenous melatonin applying confers chilling tolerance in tomato fruits by upregulating *ZAT2/6/12* giving rise to promoting endogenous polyamines, proline, and nitric oxide accumulation by triggering arginine pathway activity

Morteza Soleimani Aghdam, Zisheng Luo, Abbasali Jannatizadeh, Morteza Sheikh-Assadi, Yavar Sharafi, Boukaga Farmani, Javad Rezapour Fard, Farhang Razavi

PII: S0308-8146(18)31731-X
DOI: <https://doi.org/10.1016/j.foodchem.2018.09.157>
Reference: FOCH 23644

To appear in: *Food Chemistry*

Received Date: 9 July 2018
Revised Date: 23 September 2018
Accepted Date: 24 September 2018

Please cite this article as: Aghdam, M.S., Luo, Z., Jannatizadeh, A., Sheikh-Assadi, M., Sharafi, Y., Farmani, B., Fard, J.R., Razavi, F., Employing exogenous melatonin applying confers chilling tolerance in tomato fruits by upregulating *ZAT2/6/12* giving rise to promoting endogenous polyamines, proline, and nitric oxide accumulation by triggering arginine pathway activity, *Food Chemistry* (2018), doi: <https://doi.org/10.1016/j.foodchem.2018.09.157>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Employing exogenous melatonin applying confers chilling tolerance in tomato fruits by upregulating ZAT2/6/12 giving rise to promoting endogenous polyamines, proline, and nitric oxide accumulation by triggering arginine pathway activity

Morteza Soleimani Aghdam^{1*}, Zisheng Luo^{2}, Abbasali Jannatizadeh¹, Morteza Sheikh-Assadi³, Yavar Sharafi⁴, Boukaga Farmani⁵, Javad Rezapour Fard⁶, Farhang Razavi⁷**

¹*Department of Horticultural Science, Imam Khomeini International University, Qazvin, Iran, E-mail address: jannatizadeh@ut.ac.ir*

²*Zhejiang University, College of Biosystems Engineering and Food Science, Key Laboratory of Agro-Products Postharvest Handling Ministry of Agriculture, Zhejiang Key Laboratory for Agro-Food Processing, Hangzhou, 310058, People's Republic of China.*

³*Department of Horticultural Science, University College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran, E-mail address, m.asadi66@ut.ac.ir*

⁴*Department of Horticulture, Faculty of Agriculture, Shahed University, Tehran, Iran, E-mail address, y.sharafi@shahed.ac.ir*

⁵*Department of Food Science and Technology, Faculty of Ahar Agriculture, University of Tabriz, Tabriz, Iran, E-mail address, bfarmani@tabrizu.ac.ir*

⁶*Department of Horticultural Sciences, Faculty of Agricultural Sciences, Urmia University, Urmia, Iran, E-mail address, j.rezapourfard@urmia.ac.ir*

⁷*Department of Horticulture, Faculty of Agriculture, University of Zanjan, Zanjan, Iran, E-mail address, razavi.farhang@znu.ac.ir*

***Corresponding author:** E-mail address: aghdam@ut.ac.ir (Morteza Soleimani Aghdam).

**** Corresponding author.** E-mail address, luozisheng@zju.edu.cn (Zisheng Luo)

Download English Version:

<https://daneshyari.com/en/article/11027427>

Download Persian Version:

<https://daneshyari.com/article/11027427>

[Daneshyari.com](https://daneshyari.com)