Accepted Manuscript



Biofilm-derived spores of *Clostridioides (Clostridium) difficile* exhibit increased thermotolerance compared to planktonic spores

D.S. Pickering, M.H. Wilcox, C.H. Chilton

PII:	S1075-9964(18)30175-6
DOI:	10.1016/j.anaerobe.2018.10.003
Reference:	YANAE 1954
To appear in:	*Anaerobe*
Received Date:	30 August 2018
Accepted Date:	04 October 2018

Please cite this article as: D.S. Pickering, M.H. Wilcox, C.H. Chilton, Biofilm-derived spores of *Clostridioides (Clostridium) difficile* exhibit increased thermotolerance compared to planktonic spores, *Anaerobe* (2018), doi: 10.1016/j.anaerobe.2018.10.003

Biofilm-derived spores of *Clostridioides (Clostridium) difficile* exhibit increased thermotolerance compared to planktonic spores

Pickering DS[1*], Wilcox MH[1, 2], Chilton CH[1]

1. Healthcare Associated Infections Research Group, Leeds Institute for Biomedical and Clinical Sciences, University of Leeds, West Yorkshire, UK.
2. Microbiology, Leeds Teaching Hospitals Trust, Leeds, UK.

***corresponding author**

Email; umdsp@leeds.ac.uk