


## Accepted Manuscript

Seroprevalence and risk factors for *Neospora caninum* infection in cattle from the eastern Antioquia, Colombia

Horwald Alexander Bedoya Llano , Marcelo Sales Guimarães , Rodrigo Martins Soares , Gina Polo , Andréa Caetano da Silva

PII: S2451-943X(17)30148-5  
DOI: [10.1016/j.vas.2018.03.001](https://doi.org/10.1016/j.vas.2018.03.001)  
Reference: VAS 20



To appear in: *Veterinary and Animal Science*

Received date: 20 September 2017  
Revised date: 2 March 2018  
Accepted date: 20 March 2018

Please cite this article as: Horwald Alexander Bedoya Llano , Marcelo Sales Guimarães , Rodrigo Martins Soares , Gina Polo , Andréa Caetano da Silva , Seroprevalence and risk factors for *Neospora caninum* infection in cattle from the eastern Antioquia, Colombia, *Veterinary and Animal Science* (2018), doi: [10.1016/j.vas.2018.03.001](https://doi.org/10.1016/j.vas.2018.03.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- First risk factor-study on *N. caninum* infection in Antioquia department, Colombia.
- Horizontal transmission is an important mode of infection in the studied region.
- Abortion, age, origin and poor hygienic practices associated with manual milking are risk factors for *N. caninum* infection.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/11033987>

Download Persian Version:

<https://daneshyari.com/article/11033987>

[Daneshyari.com](https://daneshyari.com)