Double-stranded RNA injected into female Black Tiger shrimp (Penaeus monodon) prior to spawning does not transfer to progeny

M. Rao, J.A. Cowley, B.S. Murphy, C.N. Stratford, M.J. Sellars

PII: S0044-8486(18)31203-1
DOI: doi:10.1016/j.aquaculture.2018.09.059
Reference: AQUA 633583

To appear in: *aquaculture*

Received date: 5 June 2018
Revised date: 27 September 2018
Accepted date: 28 September 2018

Please cite this article as: M. Rao, J.A. Cowley, B.S. Murphy, C.N. Stratford, M.J. Sellars , Double-stranded RNA injected into female Black Tiger shrimp (Penaeus monodon) prior to spawning does not transfer to progeny. Aqua (2018), doi:10.1016/j.aquaculture.2018.09.059

Double-stranded RNA injected into female Black Tiger shrimp (*Penaeus monodon*) prior to spawning does not transfer to progeny

Rao, M.[1], Cowley, J.A.[1], Murphy, B.S.[2], Stratford, C.N.[2], Sellars, M.J.[1,*]

CSIRO Agriculture and Food, Aquaculture Program

[1] Queensland Bioscience Precinct, 306 Carmody Road, St Lucia, QLD 4067, Australia

[2] Bribie Island Research Centre, 144 North Street, Woorim, QLD 4507, Australia

[*]Corresponding author:

Melony J Sellars

Phone +61 437 025 821

Email Melony.Sellars@csiro.au

Keywords: dsRNA, RNAi, *Penaeus monodon,* shrimp, virus, Black Tiger prawn, Giant Tiger prawn, Leader prawn