



Theoretical study of the stability of small triply charged carbon clusters C_n^{3+} ($n = 3-12$)

G. Sánchez-Sanz^a, S. Díaz-Tendero^{a,*}, F. Martín^{a,b}, M. Alcamí^a

^a Departamento de Química, Módulo 13, Facultad de Ciencias, Universidad Autónoma de Madrid, 28049 Madrid, Spain

^b Instituto Madrileño de Estudios Avanzados en Nanociencias (IMDEA-Nanociencia), Cantoblanco, 28049 Madrid, Spain

ARTICLE INFO

Article history:

Received 10 June 2010

Received in revised form 3 September 2010

Accepted 3 September 2010

Available online 16 September 2010

Keywords:

Carbon cluster

Triply charged

Theoretical study

Dissociation energy

Ionization potential

ABSTRACT

Using density functional theory, coupled cluster and multireference methods, dissociation energies and 3rd ionization potentials for, respectively, triply charged and neutral carbon clusters have been evaluated. The results show that the smaller C_n^{3+} clusters are metastable, i.e., they present a fragmentation channel with negative dissociation energy. The lowest dissociation channel always corresponds to evaporation of a singly charged carbon atom. Good agreement with available experimental data is found for most two-fragment channels. The third ionization potential of the corresponding neutral species decreases with cluster size.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Small carbon clusters have been widely studied over the last two decades both theoretically and experimentally (see, e.g., the reviews [1–4]) due to their interest in astrophysical problems (see, e.g., [5–11]). Indeed, molecules exclusively formed by carbon atoms have been detected in the interstellar space and they have importance in the chemistry of carbon stars [12,13], comets [14], and interstellar molecular clouds [5,6,15–18]. Knowing the geometric, energetic and spectroscopic properties of these species is thus of great interest in order to interpret and analyze astrophysical data. In achieving this goal, theory has always played a fundamental role. Among the large number of theoretical studies devoted to neutral species, one must mention the work of Martin and coworkers [19–26]. Using high-level *ab initio* quantum chemistry methods, these authors have predicted structural, rotational, vibrational and electronic properties of neutral C_n clusters up to $n = 18$. Structural and spectroscopic properties of C_4 , C_5 and C_6 clusters have also been theoretically studied by Masso et al. [27–29]. Positively charged carbon clusters have been much less studied theoretically. Giufreda et al. [30] have used density functional theory (DFT) and coupled cluster (CC) methods to evaluate structural, rotational, vibrational, and electronic properties of linear and cyclic singly charged C_n^+ clusters with $n = 4-19$. For small doubly charged carbon

clusters, multireference calculations were carried out by Hogreve, namely for C_2^{2+} [31], C_3^{2+} [32], C_4^{2+} [33] and C_5^{2+} [34]. In Ref. [35], this work was extended to larger C_n^{2+} clusters ($n = 3-9$) by using DFT and CC methods. A semiempirical tight-binding model has also been employed for the study of neutral and (multi) charged clusters [36].

Fragmentation is the main deexcitation channel of highly excited carbon clusters [3,37,38]. Thus, to obtain information on the stability of these clusters, experimentalists have focused on the detection of the different fragments arising from neutral [39–41], singly negatively charged [42,43], singly positively charged [44–45], and multiply charged carbon clusters [41]. In the latter work, branching ratios for all the fragmentation channels resulting from the collision of swift C_n^+ projectiles with noble gases were determined. As shown in Ref. [56], comparison of the measured fragmentation branching ratios with theoretical calculations allows one, in some cases, to extract the energy deposited in the collision. Very recently, the cluster internal energy resulting from the collision has been evaluated for carbon clusters C_n^{q+} with $n = 5-10$ and $q = 2-4$ [57].

Despite the experimental efforts to understand the stability of small highly charged carbon clusters, there is a lack of theoretical information for charges larger than +2. The aim of this paper is to provide energy data for small triply charged carbon clusters, such as dissociation energies, and the sequence of ionization potentials that lead to the triply charged species. The present data are thus of great importance to help in the interpretation of recent [58] and current experimental work on the fragmentation of C_n^{3+} clusters.

* Corresponding author.

E-mail address: sergio.diaztenero@uam.es (S. Díaz-Tendero).

These clusters are formed in collisions of singly charged C_n^+ projectiles with noble gases, in which the former are doubly ionized and excited.

The paper is organized as follows. The theoretical methods used in our calculations are described in Section 2. Results and discussion are presented in Section 3. Comparison between our results and recent experimental data is also included. We end with some conclusions in Section 4.

2. Computational details

In the present work we have used different methodologies for the description of triply charged small carbon clusters: DFT, CC method, Complete Active Space Self Consistent Field (CASSCF) and Complete Active Space Self Consistent Field including second-order perturbation theory (CASPT2).

For the DFT calculations, we have chosen the B3LYP functional which combines the Becke's three parameter nonlocal hybrid exchange potential [59] with the nonlocal correlation functional of Lee et al. [60]. C_n^{3+} clusters are open-shell systems, i.e., they have unpaired electrons and the unrestricted density functional theory (UDFT) has been used. In a previous work [35], we selected this functional for the description of doubly charged small carbon clusters showing that the results obtained are in good agreement with CC results. In addition, this functional has been previously used with success in the description of dissociation energies and fission barriers of (highly) charged fullerenes [61–65]. We thus consider that B3LYP is a reasonable choice; we have used it in combination with the 6-311+G(3df) basis set, which is a triple split valence basis, supplemented with a diffuse function (+), and three d-type and one f-type polarization functions. B3LYP/6-311+G(3df) has been used in the geometry optimization of all the species considered in this work.

Coupled cluster calculations, including single and double excitations, and triple excitations in a perturbative way, CCSD(T), have been carried out over the DFT optimized geometries (obtained at the B3LYP/6-311+G(3df) level) to obtain more accurate values of the energy.

DFT and CC methods are based on a single-reference wave function. To complement the theoretical description and to better understand the results obtained, we have also performed calculations for the smaller clusters using a multireference based approach. In particular, we have carried out calculations using the CASSCF method, which includes non dynamical correlation. In a second step, we have included the electronic dynamic correlation in the CASSCF wave function within the second-order perturbation theory (CASPT2). CASSCF calculations in combination with the same 6-311+G(3df) basis set have also been used to optimize the geometry of the smaller clusters. Single point CASPT2/6-311+G(3df) calculations were carried out to obtain a more accurate value of the energies. In these multireference methodologies, the choice of the active space is crucial. We have used the following selected active spaces (number of electrons, number of orbitals): (5,6) and (7,9) for C_3^{3+} and C_4^{3+} , respectively. Thus in CASSCF and CASPT2 calculations all valence electrons were correlated.

DFT and CCSD(T) calculations were performed with the Gaussian03 package [66] and the CASSCF and CASPT2 ones with the MOLPRO program [67].

3. Results and discussion

Theoretical results for carbon clusters may present serious deficiencies when single-reference methods are used. It has been shown that static correlation may be important in these systems and DFT methods often yield to instabilities in the wavefunction

(symmetry breaking, internal instabilities or RHF \rightarrow UHF instabilities) or very large spin contaminations [35]. CCSD(T) also presents serious deficiencies in some particular cases, reflected in very large values of the T1 diagnostic [35]. The multireference calculations have allowed us to: (i) analyze the instabilities and spin contamination problems that may appear in DFT calculations, (ii) obtain a deeper understanding on the wave function properties, and (iii) investigate the multiconfigurational nature of C_n^{3+} clusters. In addition, a systematic study of the stability of different isomers, their geometry, and vibrational frequencies was carried out. A detailed analysis of these technical aspects are beyond the scope of the present work and will be published elsewhere [68]. In this article, we will focus on the results of experimental interest and, from the comparison between the different theoretical methods, we will be able to show the accuracy that can be expected from our calculated data.

3.1. Ionization potentials

Third ionization potentials have been evaluated at the B3LYP/6-311+G(3df) and CCSD(T)/6-311+G(3df)//B3LYP/6-311+G(3df) levels for cluster size up to $n=12$. The results are presented in Fig. 1 together with the first and second ionization potentials taken from previous work [35]. Since, in the latter work, results were only reported for C_n clusters with sizes up to $n=9$, for completeness, in the present work, we have extended the calculations of first and second ionization potentials up to $n=12$ (see Fig. 1). CASSCF and CASPT2 calculations have been restricted to clusters with four or less atoms. It can be seen that DFT leads to first and second IPs that are ~ 0.4 and ~ 0.8 eV, respectively, above the CCSD(T) results. In contrast, it leads to third IPs ~ 0.7 eV below the CCSD(T) results. The ionization potentials resulting from the CASSCF and CASPT2 calculations differ by less than 1 eV in most cases.

As already discussed in [35], the first IP decreases very slowly with cluster size. In contrast, the second ionization potential decreases rapidly with the number of atoms. Our calculations show that this is also the case for the third IP, since the larger the cluster the easier the charge is accommodated.

3.2. Dissociation energies

In this case CASSCF and CASPT2 calculations have been restricted to clusters with four or less atoms and to fragmentation channels corresponding to evaporation of a neutral or a charged monomer. Fig. 1 shows the corresponding dissociation energies obtained with the different methods.¹ In general, a reasonable agreement between multireference methods and DFT is obtained. As a general rule, multireference methods lead to slightly lower dissociation energies than single-reference methods (DFT and CCSD(T)). However, the observed general trends are basically the same for all dissociation channels investigated here. The differences between DFT and CCSD(T) results are of the order of 1 eV and, in most cases, the former are slightly above the latter.

Being confident that the DFT-B3LYP approach leads to reasonable results, we have extended the above calculations to all the possible two-fragment dissociation channels: $C_n^{3+} \rightarrow C_{n-x}^{(3-q)+} + C_x^{q+}$. The results are shown in Fig. 2 as function of the size of the original cluster, n . As a general trend, the dissociation energy for the ejection of a neutral fragment [$C_n^{3+} \rightarrow C_{n-x}^{3+} + C_x$] decreases with cluster size, while for the other channels [$C_n^{3+} \rightarrow C_{n-x}^{(3-q)+} + C_x^{q+}$, $q > 0$] it increases rather monotonically. This can be explained in terms of

¹ C_2^{3+} is unstable in its ground state: it dissociates without barrier in $C_2^+ + C^+$ due to Coulomb explosion and, therefore, it cannot be detected experimentally. Thus, the dissociation energy of the channels involving C_2^{3+} is not given.

Download English Version:

<https://daneshyari.com/en/article/1193411>

Download Persian Version:

<https://daneshyari.com/article/1193411>

[Daneshyari.com](https://daneshyari.com)