




Spectral and thermal characterization of 3-acetyl-5-azophenyl-4-hydroxy-6-methyl-pyran-2-one and its metal complexes

Susannah Seth^{a,*}, K.K. Aravindakshan^b^a P.G. and Research Department of Chemistry, Malabar Christian College, Kozhikode 673 001, Kerala, India^b Department of Chemistry, University of Calicut, Kozhikode 673 635, Kerala, India

HIGHLIGHTS

- A novel diazo-coupled compound with phenylazo group at a rather unusual position.
- The synthesized compound was found to be acting as a potential ligand.
- Five chelates of phenylazo dehydroacetic acid were synthesized and characterized.
- Analytical, spectral and thermal data confirm the structure of the compounds.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 30 September 2012

Received in revised form 5 April 2013

Accepted 10 April 2013

Available online 23 April 2013

Keywords:

Phenylazo dehydroacetic acid

Complexes

Mass spectrum

Thermogravimetry

ABSTRACT

Five chelates of 3-acetyl-5-azophenyl-4-hydroxy-6-methyl-pyran-2-one (phenylazo dehydroacetic acid) with Cr(III), Fe(III), Ni(II), Cu(II) and Zn(II) have been synthesized and characterized by elemental analysis, magnetic susceptibility measurements, electronic, ¹H NMR, FAB mass, IR-spectral and thermal (TG/DTG) analytical techniques. In the present work it has been found that oxygen of the deprotonated —OH group and one of the azo-nitrogens of the ligand take part in coordination. The Cr(III), Fe(III) and Ni(II) complexes were found to be having octahedral geometry and the Cu(II) and Zn(II) tetrahedral.

© 2013 Elsevier B.V. All rights reserved.

Introduction

Dehydroacetic acid, 3-acetyl-4-hydroxy-6-methyl-2Hpyran-2-one (DAA) is found to be occurring in anthers of *Solanandra nitida* (solanaceae). It possesses interesting biological properties such as fungicidal and antibacterial activity and is used as a cosmetic preservative too. It can also act as a complexing ligand. Furthermore,

this compound has several reactive functional groups and can serve as a reagent in organic synthesis. The synthesis and characterization of two macrocyclic ligands, by the condensation of dehydroacetic acid with 1,2-phenylenediamine and 1,3-propylenediamine [1] have also been reported. DAA in solution exists as a completely enolised species [2]. Like phenols, it can couple with diazonium salts at C-5 and some reports [3,4] are available on the condensation of DAA with diazotised derivatives of aniline, substituted anilines, benzidine, etc. Even though some studies are available on metal chelate of DAA [5], no work has so far been

* Corresponding author. Tel.: +91 8086733577; fax: +91 4952768219.

E-mail address: susannahseth@yahoo.co.in (S. Seth).

reported on chelates of azo dehydroacetic acid (ADAA). ADAA formed stable complexes with Cr(III), Fe(III), Ni(II), Cu(II) and Zn(II). In all the complexes one of the azo nitrogens took part in coordination together with oxygen of the enolic OH. The Zn(II) complex was diamagnetic and all others showed expected magnetic moment values. All complexes were deep red/brown in color.

Experimental

Materials and methods

All the chemicals used for synthetic purposes were of Analar grade. Solvents like methanol, acetic acid, acetone, dichloromethane and chloroform were purchased from E Merck and used as received. Commercial ethanol was purified by distillation and used. The metal salts used for the synthesis of complexes were acetate of Ni(II) and Cu(II) and chlorides of Cr(III), Fe(III) and Zn(II).

Synthesis of phenylazo dehydroacetic acid (ADAA)

Aniline (0.01 mol) was diazotised and this solution of diazonium salt was added slowly drop wise to a well cooled alkaline solution of DAA (0.02 M). The mixture was allowed to stand for 0.5 h. The product, phenylazo dehydroacetic acid, (ADAA) was filtered, washed successively with very dilute HCl and water, recrystallized from methanol and stored over anhydrous MgSO₄ in a desiccator.

Synthesis of metal complexes

To a methanolic solution of ADAA, the metal salt solution, also in methanol was added drop wise with stirring. The mixture was refluxed on a steam bath for about 1 h. The metal salt to ligand ratio was kept at 1:2/1:3 for bivalent/trivalent metals, respectively. The precipitated complexes were filtered, washed with water, methanol and dried. These were recrystallized from CH₂Cl₂–CH₃OH mixture and stored over anhydrous MgSO₄ in a desiccator.

Physical measurements

The elemental percentage analyses of C, H and N were carried out using Vario EL III CHN analyzer. Some of the metal percentage analyses were conducted using ICP-AES spectrometer using argon as the conductor gas for plasma state at a core temperature of 9000 K. The other metal percentages were determined by adopting standard [6] procedure. UV–visible spectra were recorded using 10^{−3} M solutions of ligands and complexes in methanol. IR spectra

of the compounds were recorded by their thin KBr discs. The FAB mass spectra were obtained using argon as the FAB gas, accelerating voltage, 10 kV and *m*-nitrobenzyl alcohol (NBA) as matrix. The metal percentage analyses in some cases were conducted using ICP-AES spectrometer using argon as the conductor gas for plasma state at a core temperature of 9000 K. The other metal percentages were determined by adopting standard procedure. The NMR spectra were obtained using CDCl₃ as solvent and TMS as internal standard. Magnetic susceptibility was determined at room temperature on Sherwood Magway susceptibility instrument which was calibrated using Hg[Co(NCS)₄]. Thermogravimetric studies, TG/DTG were carried out in nitrogen atmosphere with a heating rate of 10 °C/min.

Results and discussion

The elemental percentages of the ligand, ADAA were found to be in fine agreement with molecular formula C₁₄N₂O₃H₁₂. ADAA formed stable complexes with Cr(III), Fe(III), Ni(II), Cu(II) and Zn(II). In all the complexes one of the azo nitrogens took part in coordination together with oxygen of the enolic OH. The Zn(II) complex was diamagnetic and all others showed expected magnetic moment values. All complexes were deep red/brown in color. The molar conductivity of the complexes in DMF solutions (10^{−3} M) at room temperature indicated that these are non-electrolytes. The analytical data are entered in Table 1.

Spectral studies

Electronic spectra and magnetic moment interpretation

The electronic spectrum of ADAA was conspicuous with the band at 278 nm which is characteristic of an azo form. The other bands of importance were at 260 nm (carbonyl) and 370 nm (N=N–). The magnetic moment value of 3.76 B.M. for the Chromium(III) complex suggested [7] an octahedral geometry. The prominent electronic spectral bands include 380 nm (⁴T_{1g}(F) → A_{2g}(F)) and 570 nm (⁴A_{2g}(F) → ⁴T_{2g}(F)) which also pointed to an octahedral geometry Fe(III) complex spectrum was difficult to interpret due to CT bands. The magnetic moment value of 5.85 B.M. indicated an octahedral geometry. The spectrum of Ni(II) complex had almost all of the peaks expected for an octahedral geometry. The bands at 376 nm (³A_{2g} → ³T_{1g}(P)) and 680 nm (³A_{2g} → ³T_{1g}(F)) were very conspicuous. The magnetic moment value of 2.92 B.M. also confirmed the octahedral geometry. The copper(II) complex exhibited bands at 316 nm and 682 nm which were characteristic overlapped bands of copper complexes. The band at 420 nm was typical [8] of a tetrahedral geometry. The

Table 1
Analytical and physical data of the ligand (LH) and its metal complexes.

Sl. No.	Compound	Color	Yield (%)	MP (°C)	Elemental analysis found (calc.%)					Magnetic moment μ B.M.	Electronic spectral data λ _{max} (nm)
					C	H	N	M	Cl		
1	C ₁₄ N ₂ O ₄ H ₁₂ (LH)	Brick red	60	50	61.70 (61.77)	4.26 (4.41)	9.95 (10.29)				260, 278, 370
2	[Cr(L) ₂ Cl(H ₂ O)]	Dark brown	65	80	51.53 (51.89)	4.10 (4.32)	8.98 (8.65)	8.65 (8.03)	5.69 (5.48)	3.76	276, 314, 380, 570
3	[Fe(L) ₂ Cl(H ₂ O)]	Dark brown	65	140	51.15 (51.58)	4.26 (4.29)	8.14 (8.59)	8.13 (8.58)	5.71 (5.45)	5.85	278, 310, 370, 514
4	[Ni(L) ₂ (H ₂ O) ₂]	Dark brown	70	160	52.72 (52.75)	4.82 (4.71)	8.84 (8.79)	9.74 (9.22)	–	2.82	278, 312, 376, 680
5	[Cu(L)(OAc)(H ₂ O)]	Dark brown	70	>300	46.64 (46.60)	4.12 (3.88)	6.79 (6.90)	15.38 (15.42)	–	2.10	278, 316, 377, 420, 682
6	[Zn(L) ₂]	Maroon	65	210	55.01 (55.32)	3.55 (3.62)	9.76 (9.22)	10.25 (10.76)	–	D ^a	278, 310, 370, 445, 512

^a Diamagnetic.

Download English Version:

<https://daneshyari.com/en/article/1230745>

Download Persian Version:

<https://daneshyari.com/article/1230745>

[Daneshyari.com](https://daneshyari.com)