



ORIGINAL ARTICLE

Thermal, photosynthesis and antibacterial studies of bioactive safrole derivative as precursor for natural flavor and fragrance



Suzan A. Khayyat ^{a,*}, Salha H. Al-Zahrani ^b

^a Chemistry Department, Faculty of Sciences, King Abdul Aziz University, Jeddah, Saudi Arabia

^b Microbiology Department, Faculty of Sciences, King Abdul Aziz University, Jeddah, Saudi Arabia

Received 27 May 2011; accepted 10 September 2011

Available online 22 September 2011

KEYWORDS

Safrole;
Epoxide;
Photoepoxidation;
Hydroperoxide;
Hydrogen peroxide;
3-Chloroperoxybenzoic acid;
Antibacterial agents;
Bacillus subtilis;
Escherichia coli;
Staphylococcus aureus

Abstract Safrole [5-allylbenzo[d][1,3]dioxole] was subjected to photochemical oxidation reaction with hydrogen peroxide in the presence of sodium lamp to give the corresponding epoxy derivative [5-oxiranylmethylbenzo[1,3]dioxole]. The thermal oxidation of safrole with 3-chloroperoxybenzoic acid at room temperature gave the same epoxide derivative in quantitative yield. Antibacterial studies were carried out on safrole and its photoproducts (safrole epoxide and safrole hydroperoxide). The results revealed that safrole hydroperoxide was the most effective than safrole epoxide than safrole against Gram-positive bacteria *Bacillus subtilis* ATCC6633, *Staphylococcus aureus* ATC C25923, and Gram negative bacteria *Escherichia coli* ATCC25422. This result proved that safrole derivatives are beneficial to human health, having the potential to be used for medical purposes.

© 2011 Production and hosting by Elsevier B.V. on behalf of King Saud University.

1. Introduction

Phytochemicals are present in a variety of plants utilized as important components of both human and animal diets. These include fruits, seeds, herbs and vegetables (Okwu, 2005).

Propenyl benzene are common aromatic compounds, these are widely used as starting compounds for the production of

various flavor and fragrances (Xu et al., 2007). Flavors and fragrances are widely used in the food beverage and cosmetic industries (Priefert et al., 2001). Most of them in the world market today are obtained by chemical synthesis. Less than 5% is extracted from plants and can therefore be classified as natural (Xu et al., 2007).

In recent years, according to the Food and Drug Administration (FDA) and European legislation products obtained by photo and biotechnological methods can also be considered natural, if the substrate for the process is of natural origin (Serra et al., 2005).

Medicinal and aromatic plants have demonstrated its contribution to the treatment of diseases such as HIV/AIDS, malaria, diabetes, sickle-cell anemia, mental disorders (Elujoba et al., 2005; Okigbo et al., 2005) and microbial infections (Okigbo and Mmeka, 2006).

* Corresponding author. Tel.: +966 504511318; fax: +966 26990781.

E-mail address: suzan122@hotmail.com (S.A. Khayyat).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

Natural phenyl terpenoides undergo oxidation on exposure to air. The oxidation process is enhanced thermally or by irradiation to form epoxide derivative, which acts as the precursor of flavors and fragrance (Elgendy and Khayyat, 2008a). This is the first step in the biosynthesis of important products (Hua et al., 2007; Yamada et al., 2007; Zang et al., 2006).

Venturello (1992) reported that oxidation of safrole (**1**) with tungsten phosphoammonium chloride complex ($(R_4N)_3PW_4O_{24}$) gives epoxy derivative **2**.

Taking into account important activities of plant phenylpropanoides, in the present work, we believed it to be relevant to examine thermal and photochemical oxidation of safrole, and study the biological activity of the oxidation product.

2. Materials and methods

2.1. Safrole [5-allylbenzo[d][1,3]dioxole] (**1**)

Safrole [5-allylbenzo[d][1,3]dioxole] (**1**) is the major component (80%) of the essential oil of sassafras (*Piper hispidinervum*) (*Piperaceae*) in its leaves. IR spectra were performed on a Perkin-Elmer 16 FPC FT-IR spectrophotometer as thin films. 1H -NMR and ^{13}C -NMR spectra were obtained in $CDCl_3$ solution with a Bruker AVANCE D.P.X. 600 MHz apparatus. GCMS were determined by Joel JMS 600H, GC Hewlett Packard, HP 6890 Series, with capillary column (30 m \times 0.32 mm \times 0.25 μ m) HP-5 cross linked 5% dimethyl polysiloxane. A sodium lamp (Phillips G/5812 SON) was used for photoirradiation reactions. Thin layer chromatography (TLC) and preparative layer chromatography (PLC): Polygram SIL G/W 254, Mecherey-Nagel. A rotatory evaporator (at 20 °C 15 torr) was used to remove the solvents.

2.2. Test organisms

Gram-positive, e.g. *Bacillus subtilis* ATCC6633 and Gram-negative, e.g. *Escherichia coli* ATCC25422, were obtained from the library of military Hospital in Riyadh. *Staphylococcus aureus* ATCC25923 was obtained from the laboratory of Jeddah King Fahad Hospital in Saudi Arabia. It was cultured on Mueller Hinton media (Oxoid CM 41) at 37 °C.

2.3. The methods

2.3.1. Photochemical oxidation of safrole (**1**) with hydrogen peroxide

A solution of 30% hydrogen peroxide H_2O_2 (2.5 ml) was carefully added in a dropwise manner over a period of 5 min to a solution of 5 mmol compound (**1**) in 25 ml of ethanol C_2H_5OH under stirring at 0 °C. The mixture was irradiated for 55 h using a sodium lamp in a nitrogen atmosphere. The mixture was then evaporated under reduced pressure at room temperature to give a resinous material. The residue was treated with 25 ml of chloroform. The extract was dried over anhydrous sodium sulfate Na_2SO_4 and evaporated under reduced pressure to give the crude product which was purified by column chromatography on silica gel adsorbent using petroleum ether (60–80 °C) and diethyl acetate (9:2) to isolate compound **2**.

2.3.2. Oxidation of safrole (**1**) using *m*-chloroperoxybenzoic acid

A solution of 10 mmol of 80% *m*-chloroperoxybenzoic acid was added cautiously dropwise over a period of 15 min to a solution of 5 mmol of compound **1** in 25 ml of chloroform under stirring at 0 °C. The mixture was then stirred at room temperature under nitrogen atmosphere. The progress of the reaction being monitored by thin layer chromatography (TLC) and peroxide test (using a 10% solution of KI). The mixture was carefully washed with a saturated aqueous solution of $NaHCO_3$ (3 \times 10 ml) and distilled water (3 \times 10 ml). The organic layer was separated, dried over anhydrous Na_2SO_4 and evaporated under reduced pressure at room temperature. The residue product was purified by column chromatography on silica gel adsorbent using petroleum ether (60–80 °C) and diethyl acetate (9:2) to isolate compound **2** as a viscous oily substance.

2.3.2.1. 5-Oxiranyl methylbenzo[1,3]dioxole (**2**). Colorless oil, $C_{10}H_{10}O_3$ (M 178.16). IR spectrum, ν , cm^{-1} : 3018, 2896, 1606.3, 1099. 1H -NMR spectrum, δ , ppm: 2.53 d (1H, 1'-H, $J = 5$ Hz), 2.75 dd (1H, 3'-H, $J = 4.9$ Hz), 2.77 d (1H, 1'-H, $J = 5$ Hz), 2.8 dd (1H, 3'-H, $J = 4.9$ Hz), 3.10 comp. pat. (1H, 2'-H), 5.91 s (2H, C^2H_2), 6.63 d (1H, 6-H, $J = 8$ Hz), 6.73 s (1H, 4'-H), 6.75 d (1H, 7-H, $J = 8$ Hz). ^{13}C -NMR spectrum, δ , ppm: 38.4 (C^1), 46.9 (C^3), 52.9 (C^2), 100.8 (C^2H_2), 108.1 (C^4), 109.8 (C^7), 121.7(C^6) 130.1 (C^5), 146.5 (C^1), 147.7(C^3). GC-MS data:retention time 12.83 min; m/z (I_{rel} %): 178.16(98)[M^+], 162 (5) [$M-O$] $^+$, 148 (30) [$M-CH_2O$] $^+$, 132(5)[C_9H_8O] $^+$, 105 (15) [C_8H_9] $^+$, 75 (7) [C_6H_3] $^+$.

2.4. Antimicrobial activity of safrole, safrole epoxide and safrol hydroperoxide

The antimicrobial activity of the above mentioned compounds was separately determined using the disk diffusion method in plates containing 15 ml of Muller–Hinton agar medium (Oxoid (CM 41), Hampshire, England) were seeded with a 24 h culture of the bacterial strains in nutrient broth, the turbidity of each bacterial suspension was adjusted to reach an optical comparison to that of a 0.5 McFarland standard, resulting in a suspension containing approximately 1–2 \times 10⁸ CFU/ml. Mueller–Hinton agar plates were inoculated by streaking the swab over the entire sterile agar surface.

Each pure compound was checked for antibacterial activity; disk of filter paper (3 mm in diameter) was soaked with 2.6 \times 10⁻⁵ mol of the compound and placed on the inoculated plate into duplicate plates of each pure compound and chloroform (solvent) as test control. The plates were allowed to stand at refrigerator temperature for 2 h for the compound to diffuse into the agar and then the cultures were incubated at 35 °C for 24 h. Antibacterial activities were determined by measuring the diameter of the inhibition zone formed around the disk for each compound.

3. Result and discussion

The essential oil of *P. hispidinervum* (C.DC.), (*Piperaceae*) contains high levels (83–93%) of safrole in leaves which can be easily extracted by hydrodistillation (Khayyat, 2011). Photochemical epoxidation of safrole (**1**) with hydrogen

Download English Version:

<https://daneshyari.com/en/article/1252814>

Download Persian Version:

<https://daneshyari.com/article/1252814>

[Daneshyari.com](https://daneshyari.com)