



Original article

Synthesis of quinazolinones from *o*-aminobenzamides and benzyl amines under metal-free conditions



Xin-Xin Qi^a, Zhen-Zhen Song^a, Jin-Long Gong^a, Zheng-Yu Fang^b, Xiao-Feng Wu^{a,c,*}

^a Department of Chemistry, Zhejiang Sci-Tech University, Xiasha Campus, Hangzhou 310018, China

^b Department of Anal-colorectal Surgery, the First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou 310006, China

^c Leibniz-Institut für Katalyse e.V. an der Universität Rostock, 18059 Rostock, Germany

ARTICLE INFO

Article history:

Received 27 May 2015

Received in revised form 2 July 2015

Accepted 7 July 2015

Available online 10 August 2015

Keywords:

Quinazolinones

Metal-free

Benzyl amine

H₂O₂

ABSTRACT

A convenient and transition-metal free protocol for quinazolinones synthesis with *o*-aminobenzamides and benzyl amines as substrates has been developed. Using H₂O₂ as the oxidant, various quinazolinones were obtained in moderate to good yields under metal- and additive-free conditions.

© 2015 Chinese Chemical Society and Institute of Materia Medica, Chinese Academy of Medical Sciences. Published by Elsevier B.V. All rights reserved.

1. Introduction

Quinazolinones represent a high value class of compounds in organic chemistry, which has a wide range of biological and pharmacological activities [1], including anti-bacterial [2], anti-cancer [3], anti-inflammatory [4], anti-microbial [5], anti-tubercular [6], anti-ulcer [7], and so on [8]. Additionally, they are important intermediates in natural products preparation [9] and applied as structural scaffold in drug discovery.

Owing to their diverse bioactivity and privileged sub-structure for drug design, the synthetic methodologies development becomes attractive and many synthetic routes have been developed [10a]. For example, copper-catalyzed Ullmann-type coupling reactions [10b], redox condensation between *o*-substituted nitrobenzenes with amines in the presence of iron and cobalt as catalysts [11], one-pot synthesis of quinazolinones from benzyl alcohols by iridium or ruthenium catalysts [12]. In these reactions, metal-salts turned out to be indispensable, and many metal wastes were generated. Besides, condensation between *o*-aminobenzamides and aldehydes with additional oxidants is also a typical method to prepare quinazolinones [13]. Recently, a procedure on

autoxidation of benzyl amines and applied in heterocycles synthesis was developed as well [14a]. Five examples of quinazolinones were prepared in moderate yields under 150 °C with 40% AcOH as the additive. Meanwhile, we reported a procedure for the transformation of benzyl amines to the corresponding imines [15]. Various imines were produced in good to excellent yields under metal-free conditions and with H₂O₂ as the green oxidant. From mechanistic point of view, the oxidation of benzyl amines to the corresponding benzaldehydes should be the first step and then followed by condensation to give the imines. Hence, we believe a procedure for quinazolinones preparation with *o*-aminobenzamides and benzylamines as the substrates without metal catalysts and using H₂O₂ as the green oxidant should be highly realizable. Under this context, we developed this interesting metal-free and additive-free procedure for the synthesis of quinazolinones from *o*-aminobenzamides and benzyl amines.

2. Experimental

A 15 mL tube was added 2-aminobenzamide (1 mmol), benzyl amine (1.5 mmol), and a stir bar. Then H₂O₂ (30 wt% in H₂O, 5 equiv.) was added by a syringe at room temperature under open air. The tube was closed and kept at 120 °C for 20 h. The conversion and yield were determined by GC and GC–MS using hexadecane (0.1 mmol) as the internal standard.

* Corresponding author at: Department of Chemistry, Zhejiang Sci-Tech University, Xiasha Campus, Hangzhou 310018, China.

E-mail address: xiao-feng.wu@catalysis.de (X.-F. Wu).

Table 1
Screening of the reaction conditions.^a

Entry	Oxidant (equiv.)	Solvent (2 mL)	Temp. (°C)	Yield (%) ^b
1	H ₂ O ₂ (5)	DMSO	100	25
2	H ₂ O ₂ (5)	DMF	100	0
3	H ₂ O ₂ (5)	DMA	100	0
4	H ₂ O ₂ (5)	dioxane	100	<5
5	H ₂ O ₂ (5)	THF	100	<5
6	H ₂ O ₂ (5)	H ₂ O	100	<5
7	H ₂ O ₂ (5)	C ₂ H ₅ OH	100	20
8	H ₂ O ₂ (5)	CH ₃ CN	100	20
9	H ₂ O ₂ (5)	CH ₃ NO ₂	100	0
10	H ₂ O ₂ (5)	DCE	100	35
11	H ₂ O ₂ (5)	–	100	50
12	H ₂ O ₂ (3)	–	100	30
13	H ₂ O ₂ (8)	–	100	<10
14	H ₂ O ₂ (5)	–	120	71
15	H ₂ O ₂ (5)	–	80	46

^a Reaction conditions: **1a** (1.0 mmol), **2a** (1.5 mmol), oxidant, solvent (2 mL), 20 h.

^b GC yield, with hexadecane as the internal standard.

3. Results and discussion

At the beginning, substrates **1a** and **2a** were chosen as the starting materials. The reaction was conducted with 5 equiv. of H₂O₂ (30 wt% in H₂O) at 100 °C in DMSO, fortunately, 25% yield of the target product **3aa** was obtained (Table 1, entry 1). Encouraged by this result, we continued with the studies of different solvents which did not improve the yield significantly (Table 1, entries 2–10). However, when we tried the reaction without solvent, the yield raised to 50% (Table 1, entry 11). Furthermore, the amount of H₂O₂ was examined, less and more H₂O₂ all decreased the yield (Table 1, entries 12 and 13). Notably, reaction temperature was found hold great influence for this transformation. 80 °C resulted in 46% yield (Table 1, entry 15), while 120 °C resulted in 71% yield (Table 1, entry 14). No better yield can be obtained under higher reaction temperature.

Inspired by this result, we next investigated the substrates scope. First, *o*-aminobenzamide and a variety of *para*-substituted benzyl amines were studied (Table 2, entries 1–10). The substrates with electron-donating group, such as methyl, *tert*-butyl group smoothly afford the quinazolinones in 59% and 69% yield (Table 2, entries 2 and 3). However, as an electron-rich group, *para*-methoxybenzylamine decreased the yield of the desired product to

Table 2
Synthesis of quinazolinones: substrates testing.^a

Entry	1	2	3	Yield 3 (%) ^b
1				71
2				59
3				69
4				44
5				50

Download English Version:

<https://daneshyari.com/en/article/1254564>

Download Persian Version:

<https://daneshyari.com/article/1254564>

[Daneshyari.com](https://daneshyari.com)