



International Conference of Sabaragamuwa University of Sri Lanka 2015 (ICSUSL 2015)

Assessment of behavioral patterns of food intakes on micro-sleepiness, with busy lifestyles

Rumesh Liyanage*, S.B.Nawaratne, Indira Wickramasinghe, K.K.D.S.Ranaweera

Department of Food Science and Technology, Faculty of Applied Sciences, University of Sri Jayawardenepura, Gangodawila, Nugegoda, Sri Lanka.

Abstract

The effect of food habits and attitudes on micro-sleepiness was attempted to study simultaneously with development of a food product to combat. Data were collected from Sri Lanka Police Traffic Division and from 250 respondents using a pre-tested questionnaire representing drivers (especially highway drivers), private and public sector workers (shift based) and cramming students (university and school). Questionnaires were directed to respondents to fill independently and analyzed statistically. Results exposed that 76.84, 96.39 and 80.93% out of total consumed rice for all three meals, which leads to obtain a higher glycemic meal. Having two hyper glycemic meals before 14.00h was identified as a cause for micro-sleepiness. Peak level of road accidents were witnessed at 14.00 - 20.00h (38.2%) and intensity of micro-sleepiness was also falling at the same period (37.36%). While from 14.00 to 16.00h was the peak time, 16.00 to 18.00h was the least; again it reappearing to some extent from 18.00 to 20.00h. Even though respondents expressed that peak hours of micro-sleepiness is 14.00-16.00h, according to police reports, peak hours fall in between 18.00-20.00h. Out of the interviewers, 69.27% strongly required to evade micro-sleepiness and keen to spend LKR 10-20 on a commercial product. Since age-old practices connected to suppress micro-sleepiness were time consuming, recent respondents (51.64%) like to have a quick answer via a product like rapid drink. Finally it was concluded that morning and noon food habits may reason for micro-sleepiness and dinner may affect for both, natural as well as micro sleepiness due to heavy glycemic load of food. According to the study, micro-sleepiness can be categorized in to three zones such as low-risk zone (08.00-10.00h and 18.00-20.00h), manageable zone (10.00-12.00h), and high-risk zone (14.00-16.00h).

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of International Conference of Sabaragamuwa University of Sri Lanka 2015 (ICSUSL 2015).

Keywords: Micro-sleepiness, Food habits, Road accidents, Glycemic Load.

* Corresponding author *E-mail address:* rumeshprasanga@gmail.com.

1. Introduction

Micro sleepiness is a temporary biological disorder, which has been a major cause of road accidents leading to physical injuries, deaths, disabilities and significant economic losses. Traffic accidents due to human errors cause many deaths and injuries all around the world. Especially the sleepiness simply feels after taking meals because body should gain more energy for food digestion. Therefore, food habits of morning and noon may cause for micro-sleepiness while dinner may cause for both, natural as well as micro sleepiness due to heavy glycaemic load of food. Micro-sleepiness may last about 1-30 seconds. The dietary habits of a person may strongly affect for this biological phenomenon which causes for half or full eye shutting unintentionally and waking up with unconscious mood. And also the micro-sleepiness is one of dangerous factors for drivers as which cause one in four fatal road accidents on highways.

Moreover, considerable fraction of Sri Lankan population does not consume a balanced diet¹. Total mean carbohydrate, protein and fat intakes of an adult are approximately 304.4g, 44.6g and 35 g and total energy generation from which 71.2, 10.8 and 18.9% respectively². Sri Lankans ingest numbers of starch sources and consume them for lunch or dinner by limiting themselves to three meals per day¹. Almost 65% consumed well beyond the upper level of the references and this is principally due to the average person's meal containing three-quarters of rice with lesser amount of vegetable curry (15g), piece of meat or fish (15g) and some starchy curry as potato or dhal¹. Incorporation of these types of higher glycaemic loads with stresses and tiredness may eventually the brain leads towards the micro sleepiness. The overall objective of this study was to evaluate the effect of food habits and attitudes of public on micro-sleepiness and to make a preventive measure via developing an effective food product.

2. Methodology

2.1. Preliminary Survey

Preliminary survey was carried out to gather information on food habits and attitudes of micro-sleepiness with a pre-designed questionnaires' from 250 respondents. Respondents were selected representing drivers (especially highway drivers), private and public sector workers (shift based) and cramming students (university and school). Questionnaires were directed to fill independently and personally. Finally collected data were analyzed statistically.

2.2. Data collection pertaining to Road accidents

Statistics on road accidents in Sri Lanka were collected from statistical unit of Sri Lanka Police Traffic Division³ for the last ten years. And collected information was analyzed statistically.

2.3. Data analysis

Collected data were analyzed through MS Excel and Excel-stat statistical programs with a view to identify peak hours of accident occurring under Sri Lankan context.

3. Results and Discussion

The occurrence of micro-sleepiness of respondents were shown in Fig 1 (a) and the number of road accidents due to sleepiness in Sri Lanka is shown in fig 1 (b). As per the graph 1 (b), large number of road accidents were occurred in between 14.00h to 20.00h (38.2%), also the highest number of road accidents happen in between 16.00h to 18.00h (13.43%) for last nine years (Sri Lanka Police Statistical Unit, 2015). Figure 1 (a) clearly indicates that peak intensity of micro sleepiness is occurring in between 14.00-16.00h (37.36%). And also found, it was gradually increasing from 10.00-16.00h and in declining trend from 16.00 to 18.00h. However, it is again reappearing in a slight incremental pattern from 18.00h to 20.00h. According to literature, micro-sleep is lasted for a short time and it is facilitated by high glycaemic load particularly from both breakfast and lunch. This finding was further validated by the statistics of

Download English Version:

<https://daneshyari.com/en/article/1266205>

Download Persian Version:

<https://daneshyari.com/article/1266205>

[Daneshyari.com](https://daneshyari.com)