



International Conference of Sabaragamuwa University of Sri Lanka 2015 (ICSUSL 2015)

## The study on thyroid status among newborns in Jaffna District in Sri Lanka

T. Yoganathan<sup>1</sup>, V. Arasaratnam<sup>2</sup>, M. Hettiarachchi<sup>3</sup>, C. Liyanage<sup>3</sup>

<sup>1</sup>Nuclear Medicine Unit, Faculty of Medicine, University of Jaffna, Sri Lanka

<sup>2</sup>Department of Biochemistry, Faculty of Medicine, University of Jaffna, Sri Lanka

<sup>3</sup>Nuclear Medicine Unit, Faculty of Medicine, University of Ruhuna, Sri Lanka

### Abstract

Iodine is an essential nutrient for the production of thyroid hormones triiodothyronine (T<sub>3</sub>) and thyroxine (T<sub>4</sub>). A woman needs more iodine during pregnancy to maintain normal metabolism as well as to meet the requirements of T<sub>4</sub> and iodide transfer to the fetus. Objective of this study was to assess the thyroid status among newborns in Jaffna District, because similar studies conducted nationally, but not in Jaffna and no data is available about the thyroid status of the neonates. Randomly selected 477 newborns in six Medical Officers of Health (MOH) Divisions out of twelve were studied. Blood spots were taken from the neonates within the one week of delivery on specified filter paper and thyroid stimulating hormone (TSH) was assayed by using enzyme-linked immunosorbent assay (ELISA)/Radioimmunoassay (RIA) technique depending on availability of the kits in the laboratory. Among the total newborns, 239 were males (50.5%). Mean birth weight (BW) of them was 3031.5(±432.6) g, while the mean length was 51.1 (±2.1) cm. BW of males ranged from 1.7 to 5.0 Kg and of females from 1.5 to 4.35 Kg. Length of the newborns ranged from 45.0 to 58.0 cm for males and from 44.0 to 57.0 cm for females. The low birth weight (LBW), normal birth weight (NBW) and higher birth weight (HBW) were 11.3 (n=54), 88.5 (n=422) and 0.2 % (n=1) respectively. Mean neonatal blood spot TSH concentration was 9.8 (±2.1) mIU/L, and ranged from 1.00 to 53.46 mIU/L. Neonatal TSH level of the entire blood spot was categorized as > 20 mIU/L and < 20 mIU/L and the blood spot TSH > 20 mIU/L was considered as positive for congenital hypothyroidism. Among the newborns, 18% (n=86) of them were identified as positive with 10 % males and 8% females. Only one newborn was diagnosed as being congenitally hypothyroid (serum TSH >9.8 mIU/L and free T<sub>4</sub> < 10 pmol/L) with very high blood spot TSH value of 360.91 mIU/L. Further, a higher prevalence (37.7%) of neonates with blood spot TSH >5 mIU/L was observed in this study.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of International Conference of Sabaragamuwa University of Sri Lanka 2015 (ICSUSL 2015).

**Keywords:** Thyroid hormone; congenital hypothyroidism; newborns; birth weight

## 1. Introduction

Iodine is an essential nutrient for the production of the thyroid hormones, triiodothyronine ( $T_3$ ) and thyroxine ( $T_4$ ). A woman needs more iodine during pregnancy to maintain normal metabolism as well as to meet the requirements of  $T_4$  and iodide transfer to the fetus. An insufficient supply of thyroid hormones to the developing brain of the fetus can result in congenital anomalies and intellectual impairment<sup>1</sup>. Further, symptoms and signs of congenital hypothyroidism (CH) are often non-specific unless tested biochemically, and CH will be frequently overlooked, resulting in irreparable neurological damage caused by thyroid hormone deficiency during this crucial period of brain development. For early detection and implementation of thyroid hormone therapy, systematic screening programs for neonatal thyroid function were introduced in many countries in the early 1970s<sup>2,3</sup>. The initial screening method was measurement of  $T_4$  by heel-prick blood spots sample. This has been superseded by measurement of thyroid stimulating hormone (TSH) level in most programmes around the world including in Sri Lanka. The major inconvenience in measurement of serum TSH level is that it is unable to detect central (hypothalamic or pituitary) hypothyroidism, a rare disorder occurring in approximately 1 in 20 000 neonates, which can be picked up only by doing  $T_4$  test. Serum TSH assay can detect the subclinical or transient primary hypothyroidism that can be missed by  $T_4$  assay in screening programs for CH<sup>4</sup>. The incidence of CH has been reported as 1 in 1500 to 1 in 2000 live births in Southern part of Sri Lanka between 2011 and 2012<sup>5</sup>. Decline in incidence may be due to the improved knowledge of pregnant mothers on iodine nutrition by health education in antenatal clinics. Only few studies have been published on the thyroid status of the newborns in Sri Lanka by assessing the neonatal TSH level. Further, after the implementation of the universal salt iodization program in Sri Lanka there have not been any published data on thyroid function among the newborns. Also, iodine supplementation during pregnancy is not a routine practice in Sri Lanka. Policy makers might be thinking that salt iodization is adequate to provide optimum iodine nutrition to the population. Therefore, this study was carried out to assess the thyroid status among newborns in Jaffna District. Although some similar studies have been conducted nationally, there are no data available about the thyroid status of the neonates in Jaffna district.

## 2. Methodology

Four hundred and seventy seven newborns of the randomly selected pregnant mothers in six (Jaffna, Uduvil, Nallur, Kopay, Karaveddy and Kays) Medical Officers of Health (MOH) Divisions out of twelve were included for this study during the period 2012/2013. Newborns were selected randomly with probability proportion to the number of deliveries occurred in the previous year of this study conducted. Ethical approval was obtained from Ethical Review Committee of the Faculty of Medicine, University of Jaffna. Results were presented by using descriptive statistics as mean ( $\pm$ standard deviation), percentage and frequency. Simple linear regression analysis was used to test the correlations between blood spot TSH and BW of the newborn. Weight of the newborn was measured by digital infant scale to the precision of 10 g with wearing nappies or only light clothing. First the “zero” level of scale was adjusted. Next the newborn was put on the weighing pan of the digital infant scale. The weighing scale was checked every morning prior to the use. The recumbent length of the infants was measured by using portable infantometer (SECA 417, Germany) with a precision of 0.1 cm. Further, examination was carried out and a heel prick blood spot (by filter paper) was collected from the neonate prior to the discharge. If the blood spot were not obtained from the newborn before the discharge from the hospital, spots were collected through home visits within a week of delivery. The cut-off values for TSH were decided depending on the age of the newborn at the time of blood spot collection and it was 40.0 mIU/L upto 48 hrs (i.e., Day 2) and 20.0 mIU/L after 48 hrs of life (i.e., Day 3 onwards). These values were based on the analysis of a pilot study done at Southern Province of Sri Lanka (Hettiarachchi & Amarasena, 2014). The blood spot TSH analyses were done using the IMMUCHEM™ NEONATAL TSH-MW ELISA (bulk kit –20 plates) and radioimmunoassay (RIA) kits (500 tubes kit) provided by MP Biochemicals, USA on availability of the kits in the laboratory. Once a positive case was found a repeat test was performed to confirm results. For that, the parents were contacted immediately through telephone in order to get a serum sample for the confirmation of the disease. Both serum TSH and free  $T_4$  were determined in these newborns using respective enzyme-linked immunosorbent assay (ELISA) kits provided by the MP Biochemicals, USA. CH was confirmed if serum TSH >9.8 mIU/L and free  $T_4$  < 10 pmol/L. Data were presented as descriptive statistics mean  $\pm$  SD, range, frequency and percentage. The coefficients of variation of the intra-assays was 5.5%, whereas inter-assay

Download English Version:

<https://daneshyari.com/en/article/1266219>

Download Persian Version:

<https://daneshyari.com/article/1266219>

[Daneshyari.com](https://daneshyari.com)