



International 58th Meat Industry Conference “Meat Safety and Quality: Where it goes?”

Microbiological and chemical evaluation of dried smoked meat product

Dragana Plavsic^{a,*}, Djordje Okanovic^a, Jasmina Gubic^a, Zvonko Njezic^a

^aUniversity of Novi Sad, Institute of Food Tehcnology in Novi Sad, Bulevar cara Lazara 1, Novi Sad, Serbia

Abstract

The aim of this study was to determine the chemical and microbiological quality of an artisanal dried smoked meat product, manufactured in a domestic household. At the end of production, the average moisture content was 37.83%, salt 5.07%, while water activity was 0.89. Yeast counts were approximately 1.0×10^7 cfu/g while mould counts were 1.2×10^6 cfu/g. Two commonly present *Penicillium* species were identified: *P. aurantiogriseum* and *P. commune*. Our results suggest that the artisan manufacturers of these types of meat product should take steps to ensure a higher quality of their products.

© 2015 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of scientific committee of The 58th International Meat Industry Conference (MeatCon2015)

Keywords: dried smoked meat product; a_w ; yeasts and moulds

1. Introduction

Dried smoked meat products are produced and consumed in different countries throughout the world¹. The production of smoked meat products combined with the use of salting, drying and smoking methods has a long tradition². Dried smoked meat products are characterized by low water activity (a_w) and high salt concentrations⁴. The primary objective of meat salting process in dry-curing is salt penetration into the muscle tissue, where salt, as

* Corresponding author. Tel.: +381 21 485 3822
E-mail address: dragana.plavsic@fins.uns.ac.rs

humectants, reduces a_w providing initial preservation effect during the first stage of production². However, if the production process is poorly managed, uncontrolled fungi (yeast and mould) growth on the surface of the dried smoked meat products can cause significant quality problems. As some moulds are mycotoxigenic, their growth on the dried smoked meat products could also pose a serious health risk⁵. Many studies have shown that xerotolerant and xerophilic fungi, species of *Aspergillus*, *Eurotium* and *Penicillium* are associated with dried smoked meat products in different parts of the world^{6,7,8} and develop in the reduced moisture content of the substrate. “Storage” moulds prefer a lower moisture content of the substrate (13-18%, $<0.75 a_w$) (dried fruit, milk in powder, grains and bakery products) and higher temperatures, so therefore, are the most frequently isolated from stored products⁹. The moulds are frequently referred to as cause for food spoilage leading to quality reduction and a major economic loss for the producers^{10,11}. Identification of moulds associated with a_w , moisture and salt contents typical for smoked dried meat products is a starting point to understand importance of the production of safe and durable products.

The aim of this study was to examine the chemical and microbiological quality of a domestically-produced dried smoked meat product.

2. Material and methods



Fig. 1. Dried smoked meat product.

A dried smoked meat product, manufactured in a domestic setting by an artisan producer was selected (Fig. 1).

Moisture and NaCl contents were quantified according to the ISO recommended standards 1442:1997¹² and 1841-1:1996¹³, respectively. a_w of the samples was measured using a water activity measurement device (TESTO 650, Germany) with an accuracy of ± 0.001 at 25°C.

The total numbers of yeasts and moulds were determined according to the SRPS ISO 21527-2¹⁴. Inoculated agar plates were incubated at 25°C. The results were read after five and seven days. Counts were performed in triplicate.

Monocultivation of moulds was conducted by transferring conidia and hypha fragments from mould colonies onto Czapek yeast (autolisate) extract agar (CYA) (NaNO₃ 3 g, K₂HPO₄ 1g, KCl 0.5 g, MgSO₄·7H₂O 0.5 g, FeSO₄·7H₂O 0.01g, yeast extract 5 g, sucrose 30 g, solution of microelements 1ml (ZnSO₄·7H₂O 1 g, CuSO₄·7H₂O 0.5 g, distilled water 100 ml), agar 20 g, distilled water 1000 ml). The criteria described by Samson et al.^{10,11} and Pitt & Hocking¹⁵ were used for species identification.

All data are presented as mean values. Analysis of variance and Duncan's multiple range tests were performed using the software package Statistica 10 for Windows, Stat Soft, Tulsa, Oklahoma, USA, 2009. Differences were considered significant at $p < 0.05$.

3. Results and discussion

Results of the chemical analyses of the dried smoked meat product are presented in Table 1. The a_w , moisture and NaCl content after drying period showed the usual trends observed in this type of product. After the drying period,

Download English Version:

<https://daneshyari.com/en/article/1266330>

Download Persian Version:

<https://daneshyari.com/article/1266330>

[Daneshyari.com](https://daneshyari.com)