

The “periodic table” of benzotriazoles: Uranium(VI) complexes☆



Sokratis T. Tsantis^a, Maria Mouzakitis^a, Aikaterini Savvidou^b, Catherine P. Raptopoulou^b, Vassilis Psycharis^{b,*}, Spyros P. Perlepes^{a,c,**}

^a Department of Chemistry, University of Patras, 26504 Patras, Greece

^b Institute of Nanoscience and Nanotechnology, NCSR “Demokritos”, 15310 Aghia Paraskevi Attikis, Greece

^c Institute of Chemical Engineering Sciences, Foundation for Research and Technology-Hellas (Forth/ICE-HT), Platani, P.O. Box 1414, 26504 Patras, Greece

ARTICLE INFO

Article history:

Received 22 May 2015

Received in revised form 26 June 2015

Accepted 29 June 2015

Available online 6 July 2015

Keywords:

Benzotriazoles

Crystal structures

Raman spectra

Uranyl complexes

ABSTRACT

The first employment of benzotriazoles in uranium(VI) chemistry has led to uranyl complexes with interesting structures and spectroscopic properties; the reactions of uranyl nitrate with 5,6-dimethylbenzotriazole give two products depending on the reactants' molar ratio.

© 2015 Elsevier B.V. All rights reserved.

A popular class of ligands consists of azoles (e.g. pyrazoles and imidazoles), and one broad family of them are benzotriazole and its derivatives (Scheme 1). Over the last 30 years or so, we [1–8] and other groups [8–17] have been exploring the coordination chemistry of benzotriazoles. Research in this area ranges from efforts to model the corrosion inhibition of some metals (particularly Cu and its alloys) by benzotriazoles [18], to the synthesis of benzotriazolato coordination clusters [2,4–7,9,11,13,15] and polymers [1,3,12,14,16,17] with esthetically pleasing structures and exciting physical properties. Although the use of this family of ligands has been investigated widely in 3d- [1–11, 13–15,17] and 4f-metal [12,16] chemistry, their employment in the chemistry of 5f-metal ions has completely been ignored. We have been trying to create a “periodic table” of the benzotriazole-based ligands; we are now pleased to be able to fill in the empty 5f-metal position in this “periodic table” by describing the syntheses, structures and preliminary spectroscopic characterization of the first U(VI) complexes with benzotriazole-based ligation.

Interest in the coordination chemistry of the uranyl group ($trans\ O=U=O$) has recently increased for several reasons [19–22]. A basic understanding of this chemistry is very important for the selective complexation and separation of this ion from acid media during

reprocessing of irradiated advanced heavy water reactor (AHWR) nuclear fuel, nuclear plant effluents, seawater, and biological and environmental samples [23]. However, few UO_2^{2+} -simple azole complexes have been reported [24], and none is known with benzotriazoles as ligands.

Thus, we have decided to amalgamate the above mentioned research topics and to study reactions of uranyl nitrate with simple benzotriazole ligands; we report here our preliminary results in this area.

The reaction of $UO_2(NO_3)_2 \cdot 6H_2O$ and Mebta in a 1:2 molar ratio in MeCN at room temperature gave a yellow-green solution that upon storage at 20 °C produced yellowish green crystals of $[UO_2(NO_3)_2(Mebta)_2]$ (**1**) in 40% yield. When an analogous reaction and crystallization was carried out by using 5MebtaH and diMebtaH, crystals of $[UO_2(NO_3)_2(6MebtaH)_2]$ (**2**) and $[UO_2(NO_3)_2(diMebtaH)_2]$ (**3**), respectively, were isolated in moderate to good yields. It is interesting to note that 5MebtaH tautomerizes into 6MebtaH ($R=R'=H$ and $R'=Me$ in Scheme 1) in complex **2**, i.e. upon coordination. A second product was isolated from the $UO_2(NO_3)_2 \cdot 6H_2O/diMebtaH$ reaction system by employing an excess of the ligand. The reaction of $UO_2(NO_3)_2 \cdot 6H_2O$ and diMebtaH in a 1:4 molar ratio in MeCN gave a dark brownish red solution that upon storage at 20 °C gave brown-yellow crystals of $[UO_2(NO_3)_2(diMebtaH)_2] \cdot (diMebtaH) \cdot MeCN$ (**4** · diMebtaH · MeCN) in yields higher than 75%.


The structures of the four complexes were determined by single-crystal X-ray crystallography. The structures of the molecules $[UO_2(NO_3)_2 L_2]$ ($L=Mehta, 6MebtaH, diMebtaH$; Fig. 1) that are present in **1–3** and **4** · diMebtaH · MeCN present similarities and characteristic differences. Complex **2** contains two structurally similar, crystallographically independent $[UO_2(NO_3)_2(6MebtaH)_2]$ molecules. The U^{VI} atoms

☆ In memory of Paul Cordopatis: A great scientist, a fantastic mentor and a precious friend.

* Corresponding author.

** Correspondence to: S.P. Perlepes, Department of Chemistry, University of Patras, 26504 Patras, Greece.

E-mail addresses: v.psycharis@inn.demokritos.gr (V. Psycharis), perlepes@patras.upatras.gr (S.P. Perlepes).



Scheme 1. The general structural formula of benzotriazoles. The parent member of this family of ligands is 1H-benzo[d][1,2,3]triazole or simply benzotriazole (btaH; $R=R'=R''=H$). R, R' and R'' can be non-donor (Me, Cl, NO_2 ,...) or donor (OH, COOH, CH_2OH , CH_2COOH ,...) groups. The ligands used and discussed in this work are 1-methylbenzotriazole (Mebta; $R=\text{Me}$, $R'=R''=H$), 5-methylbenzotriazole (5MebtaH; $R=\text{Me}$, $R=R''=H$) and 5,6-dimethylbenzotriazole (diMebtaH; $R'=R''=\text{Me}$; $R=H$).

are at inversion centers in **1–3**, whereas no molecular symmetry elements are present in **4** · diMebtaH · MeCN. The gross molecular structure is common to all four complexes and show that each metal ion is surrounded by six oxygen and two nitrogen atoms in a hexagonal bipyramidal geometry. The two uranyl oxygen atoms occupy the axial positions. Four oxygen atoms of the two bidentate chelating nitrate groups, together with two nitrogen atoms of the neutral, planar monodentate benzotriazole ligands form the equatorial hexagonal plane. The U^{VI} and donor atoms in the equatorial plane are almost coplanar. The donor atom of Mebta, 6MebtaH and diMebtaH is the nitrogen atom of the position 3 of the azole ring (Scheme 1). The $\text{U}=\text{O}$ bond lengths [1.734(5)–1.754(6) Å] are unremarkable [23], while the $\text{U}-\text{O}_{\text{nitrate}}$ bond lengths are in the range 2.479(5)–2.529(5) Å with the nitrate $\text{N}-\text{O}$ bond lengths being asymmetric due to coordination. The nitrate $\text{N}-\text{O}_{\text{coordinated}}$ bond distances are in the range 1.251(10)–1.337(12) Å, with the $\text{N}-\text{O}_{\text{uncoordinated}}$ bond distances in the range 1.206(8)–1.217(8) Å. The $\text{U}-\text{N}_{\text{benzotriazole}}$ bond distances are between 2.507(5) and 2.551(7) Å.

The differences of the four complexes concern the relative orientations of the benzotriazole ligands. In compounds **1**, **2** and **3**, the benzene rings of the ligands are at opposite sides of the molecules (Fig. 1a, b, c, d),

whereas in the case of **4** (the molecule lacks a center of symmetry) the benzene rings of the two *trans* diMebtaH ligands are on the same side of the molecule (Fig. 1e). Another remarkable difference concerns the twist of the planes of the organic ligands relative to the planes of the coordinated nitrate groups. The dihedral angles between these planes are 86.9(3)° for **1**, 84.2(4)° for the U1-containing molecule of **2**, 68.7(3)° for the U2-containing molecule of **2**, 55.7(2)° for **3** and 74.8(2)°/84.4(2)° for **4**.

H bonds and $\pi-\pi$ stacking interactions build the crystal structures of the four complexes. The most interesting crystal structure is that of **2** (Fig. 2), in which a 3D network is formed. The crystal structures of complexes **1**, **3** and **4** · diMebtaH · MeCN are shown in Figs. S1–S3 and full descriptions are provided in their legends.

The IR spectra of the complexes (Figs. S4–S7) show the existence of a strong band at 944 (**1**), 948 (**2**), 950 (**3**) and 934 (**4** · diMebtaH · MeCN) cm^{-1} (not present in the spectra of the free ligands) assigned to the IR-active antisymmetric vibration of the $\{\text{O}=\text{U}=\text{O}\}^{2+}$ group (ν_3) [25], which is red-shifted compared to the corresponding band for the aquo uranyl complexes ($\sim 965 \text{ cm}^{-1}$); this vibration does not appear in the Raman spectra of **1–3**, consistent with a centrosymmetric uranyl group in these complexes. The Raman-active uranyl symmetric stretch (ν_1) is observed at 855–865 cm^{-1} in the Raman spectra (Fig. 3) and it is absent from the IR spectra of **1–3**. The non-centrosymmetric nature of the UO_2^{2+} group and the lattice diMebtaH's interaction with one uranyl oxo-oxygen atom in **4** · diMebtaH · MeCN are also corroborated using IR and Raman spectroscopies; the ν_3 and ν_1 vibrations appear in both the IR (934, 860 cm^{-1}) and Raman (930, 863 cm^{-1}) spectra (Fig. S8) of this complex [26].

The ^1H -NMR spectra (Figs. S9–S11) of the uranyl complexes **1**, **2** and **4** · diMebtaH · MeCN in d_6 -DMSO reveal that these compounds decompose in solution to release the benzotriazole ligand; the spectra of the complexes are identical with those of the corresponding free ligands Mebta, 5MebtaH and diMebtaH, respectively. For example, the spectrum of **1** shows a sharp singlet at δ 4.30, a triplet at δ 7.39, a triplet at δ 7.55, a doublet at δ 7.83 and a doublet at δ 8.01 ppm, with relative




Fig. 1. Partially labeled plots of all the crystallographically independent molecules that are present in the structures of complexes **1** (a), **2** (b and c), **3** (d) and **4** (e). H atoms have been omitted. Symmetry codes ('): $1-x, 1-y, -z$; (*): $1-x, 1-y, 1-z$; and (**, #): $-x, 1-y, -z$. Symmetry-related uranyl oxido atoms and nitrate ligands have not been labeled.

Download English Version:

<https://daneshyari.com/en/article/1301424>

Download Persian Version:

<https://daneshyari.com/article/1301424>

[Daneshyari.com](https://daneshyari.com)