

Note

## Cymantrene-derived monodentate phosphites: New ligands for Rh-catalyzed enantioselective hydrogenation

Sergey E. Lyubimov <sup>a,\*</sup>, Vadim A. Davankov <sup>a</sup>, Nikolay M. Loim <sup>a</sup>, Lyudmila N. Popova <sup>a</sup>, Pavel V. Petrovskii <sup>a</sup>, Petr M. Valetskii <sup>a</sup>, Konstantin N. Gavrilov <sup>b</sup>

<sup>a</sup> Institute of Organoelement Compounds, Russian Academy of Sciences, 28 Vavilov street, 119991 Moscow, Russian Federation

<sup>b</sup> Department of Chemistry, Ryazan State University, 46 Svoboda street, 390000 Ryazan, Russian Federation

Received 29 June 2006; received in revised form 8 August 2006; accepted 10 August 2006

Available online 22 August 2006

### Abstract

Novel chiral monodentate phosphite ligands bearing cymantrene fragment have been prepared. Phosphite **4** with PPh<sub>3</sub>, instead of CO-ligand, in cymantrene moiety provided higher enantioselectivity than its unsubstituted analogue **3** in hydrogenation of (*Z*)-methyl-2-acet-amido-3-phenylacrylate.

© 2006 Elsevier B.V. All rights reserved.

**Keywords:** Phosphite ligands; Cymantrene; Asymmetric hydrogenation

### 1. Introduction

Bidentate derivatives of ferrocene, cymantrene, and some tricarbonyl complexes of chromium and rhenium have been actively investigated and demonstrated from good to excellent enantioselectivity in asymmetric hydrogenation, allylation, hydrosilylation and conjugate addition (for example, see [1–8]). Despite the encouraging performance of many bidentate ligands, especially phosphines, recent breakthroughs have shown that the use of bidentate ligands is not essential to obtain high enantioselectivity. Chiral monodentate phosphines [9], phosphonites [10], phosphites [11] and phosphoramidites [12] have been reported to be excellent ligands in the rhodium-catalyzed asymmetric hydrogenation. Undoubted leaders among modern P-monodentate ligands are phosphite and phosphoramidite derivatives of BINOL [11–13, and references cited therein]. They may have either only axial chirality or contain additional stereogenic centers. The latter group has been well developed during the last years; for example,

ferrocenylethyamine-derived phosphoramidites showed up to 99.5% ee in hydrogenation of enamides and dehydro-amino acids [12]. But ligands containing only axial chirality are more attractive because of their synthetic availability and low cost [14].


Herein, we report synthesis of BINOL-based monodentate phosphites derived from accessible cymantrenylcarbinol and its derivative obtained by changing a CO-ligand for PPh<sub>3</sub>. Due to the lack of additional stereocentres, correct correlation between the ligands structure and their efficiency in the Rh-catalyzed asymmetric hydrogenation can be made.

### 2. Results and discussion

Cymantrenylcarbinol was easily prepared from formyl-cymantrene [15]. To increase the electron-donor and steric demands of **1** the photochemical substitution of a CO ligand by PPh<sub>3</sub> was carried out (Scheme 1). Compound **2** showed two multiple signals for Cp-protons at  $\delta$  4.16 (2H) and  $\delta$  4.52 (2H) in the <sup>1</sup>H NMR spectrum in acetone-*d*<sub>6</sub> but only first of them possessed a PH-coupling of 3.2 Hz. Analysis of the NOESY spectrum of **2** and synthe-

\* Corresponding author. Tel./fax: +7 495 135 6471.

E-mail address: [lssp452@mail.ru](mailto:lssp452@mail.ru) (S.E. Lyubimov).



Scheme 1. Synthesis of cymantrenylcarbinols.

sis of its 2,5-D labeled analogue starting from 2,5-D labeled alcohol **1** [15] revealed that a PH-coupling exists between the P-atom of PPh<sub>3</sub> and protons at 3,4-positions of the Cp-ring. Selectivity of the observed coupling is connected with the hindered conformational rotation of the Mn(CO)<sub>2</sub>PPh<sub>3</sub> moiety.

Novel BINOL-based ligands were readily synthesized by direct phosphorylation of the corresponding cymantrene alcohols **1** and **2** (Scheme 2).

They represent crystalline compounds stable in prolonged storage in dark place.


To examine the catalytic behavior of the both cymantrene-based ligands **3** and **4** in asymmetric hydrogenation cationic rhodium(I) complexes **5** and **6** were obtained (Scheme 3).

Significantly larger  $\nu(\text{CO})$  frequency for the ligand **3** and its Rh(I) complex **5** (Table 1) prove higher  $\pi$ -acidity of phosphite **3** in comparison to **4** having a more efficient electron-donating ligand P(Ph)<sub>3</sub> than CO in the cymantrene part. We supposed that the different electron-donating properties of the ligands should have influence on the catalytic behavior of the complexes.

Complexes **5** and **6** were first used in the hydrogenation of (*Z*)-methyl 2-acetamido-3-phenylacrylate **7** (Scheme 4).

Both complexes showed excellent chemical yields. In the case of **5** poor enantioselectivity were obtained in all solvents used (Table 2, entries 1 and 2).


Contrary, the use of the complex **6** dramatically improved the enantioselectivity and gave product **8** in 91% ee (Table 2, entry 3). The result indicated that the bulky electron-donating ligand **4** tended to increase enantiomeric excess for the product **8**. However, the nature of the solvents showed no less important effect, changing the solvent from CH<sub>2</sub>Cl<sub>2</sub> to EtOAc reduced the ee value of the product **8** from 91% to 27%. Closer effect was published by Hua and van den Berg for bulky phosphites and phosphoramidites [16,17].

Scheme 2. Synthesis of ligands **3** and **4**.

Scheme 3. Complexation of the ligand with Rh(I).

Table 1  
Selected spectroscopic data for compounds **3–6**

Compound	IR, $\nu(\text{CO})$ , $\text{cm}^{-1}$ (CHCl <sub>3</sub> )	<sup>31</sup> P NMR, $\delta_{\text{P}}$ (CDCl <sub>3</sub> )
<b>3</b>	2028, 1944	137.7
<b>4</b>	1940, 1872	142.9, 91.1
<b>5</b>	2024, 1940	125.3 ( <sup>1</sup> J <sub>(P,Rh)</sub> , 241.9 Hz)
<b>6</b>	1932, 1864	125.4 ( <sup>1</sup> J <sub>(P,Rh)</sub> , 243.5 Hz), 91.85

Scheme 4. Hydrogenation of (*Z*)-methyl 2-acetamido-3-phenylacrylate.

Influence of ligand structure on the enantioselectivity and chemical yield was also examined in hydrogenation of less sterical hindered substrate – dimethyl itaconate (**9**) (Scheme 5).

In this case, moderate enantioselectivities were obtained for both complexes **5** and **6**. The use of complex **6** with bulky phosphite **4** led to a completely different behavior during hydrogenation of **9**. In this case not only conversion was lower, but the enantioselectivity was decreased due to increased steric demands of ligand **4** as compared with **3** (Table 3). Solvent effect is also important, complex **6** showed only 30% conversion in CH<sub>2</sub>Cl<sub>2</sub> (Table 3, entry 3), however EtOAc as solvent afforded product **10** with excellent chemical yield, but lower ees (Table 3, entry 4).

In summary, we have synthesized novel and easily prepared cymantrene-derived monodentate phosphites. It has been found that the more sterically hindered and better electron-donating phosphite **4** with a P(Ph)<sub>3</sub> group as ligand in cymantrene was a better asymmetric catalyst in hydrogenation of a dehydroamino acid ester (up to 91% ee), but less effective in hydrogenation of dimethyl itaconate. Further modification of cymantrene part, for example, substitution of second CO ligand by P(Ph)<sub>3</sub>, may improve enantioselectivity. Such experiments are in progress.

Download English Version:

<https://daneshyari.com/en/article/1328427>

Download Persian Version:

<https://daneshyari.com/article/1328427>

[Daneshyari.com](https://daneshyari.com)