

Journal Pre-proof

Correlates of violent suicide attempts in patients with bipolar disorder

Gianluca Rosso, Umberto Albert, Stefano Bramante, Elena Aragno, Federica Quarato, Gabriele Di Salvo, Giuseppe Maina



PII: S0010-440X(19)30059-8

DOI: <https://doi.org/10.1016/j.comppsy.2019.152136>

Reference: YCOMP 152136

To appear in:

Please cite this article as: Rosso G, Albert U, Bramante S, Aragno E, Quarato F, Di Salvo G, Maina G, Correlates of violent suicide attempts in patients with bipolar disorder, *Comprehensive Psychiatry* (2019), doi: <https://doi.org/10.1016/j.comppsy.2019.152136>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier.

Title: Correlates of violent suicide attempts in patients with bipolar disorder.

Author names and affiliations: Gianluca Rosso^a, Umberto Albert^c, Stefano Bramante^{a,b}, Elena Aragno^{a,b}, Federica Quarato^{a,b}, Gabriele Di Salvo^{a,b}, Giuseppe Maina^{a,b*}

- a. Psychiatric Unit, San Luigi Gonzaga Hospital, Regione Gonzole 10, 10043, Orbassano, Turin, Italy.
- b. Department of Neurosciences “Rita Levi Montalcini”, University of Turin, Via Cherasco 15, 10126, Turin, Italy.
- c. Department of Medicine, Surgery and Health Sciences, Psychiatric Section, University of Trieste, Via Guglielmo de Pastrovic 4 -34128, Trieste, Italy.

***Corresponding author**

Giuseppe Maina

Psychiatric Unit, San Luigi Gonzaga Hospital, Regione Gonzole 10, 10043, Orbassano, Turin, Italy.

Telephone number: +39 011 9026517

Fax number: +39 011 9026669

E-mail address: giuseppe.maina@unito.it

Highlights:

- Cross-sectional study to analyze violent suicide attempts in bipolar disorder;
- 30.6% of attempters used a violent suicide method;
- More men in the group of violent suicide attempters;
- Association between obesity and violent suicide.

Abstract

BACKGROUND: Suicide is one of the leading causes of death in bipolar disorder (BD); violent suicide attempts are associated with the highest level of lethality. We aimed to evaluate factors related to the risk of violent

Download English Version:

<https://daneshyari.com/en/article/13423515>

Download Persian Version:

<https://daneshyari.com/article/13423515>

[Daneshyari.com](https://daneshyari.com)