



Flavonoids from *Casearia sylvestris* Swartz variety *lingua* (Salicaceae)



Paula Carolina Pires Bueno^a, Fernando Passareli^a, Naira Buzzo Anhesine^a,
Roseli Buzanelli Torres^b, Alberto José Cavaleiro^{a,*}

^a Institute of Chemistry, Department of Organic Chemistry, São Paulo State University – UNESP, Araraquara, SP, Brazil

^b Agronomic Institute of Campinas, Herbarium IAC, Campinas, SP, Brazil

ARTICLE INFO

Article history:

Received 3 February 2016

Received in revised form 3 June 2016

Accepted 4 June 2016

Keywords:

Casearia sylvestris variety *lingua*

Salicaceae

Glycosylated flavonoids

1. Subject and source

Casearia sylvestris Sw. (Salicaceae) is the most widely studied species of the genus *Casearia* Jacq. due to its use in folk medicine and its unique biological properties (Santos et al., 2010). *C. sylvestris* is highly adaptable, its range extending from Mexico to South America. Based on the external morphological differences of *C. sylvestris* specimens, Sleumer (1980) proposed two varieties of this species: *C. sylvestris* var. *sylvestris*, which inhabits humid and dense forests; and *C. sylvestris* var. *lingua*, which commonly occurs in open and xeric habitats (Cavallari et al., 2010). Most previous studies of the biological activities of *C. sylvestris* have employed *C. sylvestris* var. *sylvestris*. To date, *C. sylvestris* var. *lingua* has not been the primary focus of phytochemical investigations.

In the present work, leaves of *Casearia sylvestris* var. *lingua* (Salicaceae) were collected in August 2012 in the city of Araraquara (São Paulo State, Brazil; 21°49.300'S, 48°11.460'W). A voucher specimen (IAC 55839) was deposited at the Herbarium of the Agronomic Institute of Campinas. After collection, the leaves were immediately dried at 40 °C in an oven with air circulation and were then crushed with liquid nitrogen in an analytical mill. The resulting powder was stored at room temperature until used in the phytochemical procedures. For comparison, leaves of *Casearia sylvestris* var. *sylvestris* (IAC 55840) were also collected from the same field and were processed and deposited under the same conditions described above for *C. sylvestris* var. *lingua*.

* Corresponding author. Institute of Chemistry, Department of Organic Chemistry, São Paulo State University – UNESP, CEP 14801-970, Araraquara, SP, Brazil.

E-mail address: albjcava@iq.unesp.br (A.J. Cavaleiro).

2. Previous work

Previous chemical investigations of the genus *Casearia* have led to the identification of more than 285 compounds, including terpenoids, steroids, phenylpropanoids, and flavonoids. These include 152 clerodane diterpenes, which are considered taxonomic markers for this genus (Xia et al., 2015). Phytochemical investigations of *C. sylvestris* revealed the presence of approximately 41 clerodane diterpenes including casearins, (Itokawa et al., 1990; Morita et al., 1991; Carvalho et al., 1998; Santos et al., 2007, 2010; Wang et al., 2009a), casearvestrins (Oberlies et al., 2002), and caseariasides (Wang et al., 2009b), among others. These compounds were found in the leaves, stem, stem bark, roots, and seeds of the plant (Carvalho et al., 2009). In addition to clerodane diterpenes, other compounds observed in this species include non-clerodane diterpenoids, sesquiterpenoids, phenylpropanoids, and phenolic compounds (Xia et al., 2015). Four ellagic acid derivatives were identified in *C. sylvestris* aqueous extracts (Da Silva et al., 2008a), and compounds found in alcoholic extracts include two gallic acid derivatives (Da Silva et al., 2008b), tyrosol (Wang et al., 2009c), and rutin (Silva et al., 2006).

3. Present study

Powdered leaves (27 g) of *C. sylvestris* var. *lingua* were successively extracted three times, at room temperature, using 175 mL of a 50:30:20 (% v/v) mixture of water/ethanol/isopropanol (Bueno et al., 2015). After evaporation of the alcohol, the extracts were lyophilized to obtain 5.6 g of crude dry extract (CDE). A portion (3 g) of this extract was fractionated using a medium-pressure chromatography system (Puriflash 4100, Interchim) equipped with two C-18 flash columns (Puriflash C18HQ, 15 μ m, 35 g, Interchim). Elution was performed using a gradient of water (A) and methanol (B): 5% of B from 0 to 8 min, 5–25% of B to 35 min, 25–100% of B to 65 min, and 100% of B to 110 min. The flow rate was 8 mL/min and the UV detection wavelength was 254 nm. Fractions 2, 3, and 4 were combined and lyophilized to yield 2.3 g of dry material (denoted CDE-F1). The isolated CDE-F1 fraction was submitted to solid phase extraction (SPE) with polyamide, and the resulting methanolic fraction was dried to give 113 mg of dry matter. Finally, the methanolic fraction was dissolved in water/acetonitrile (80:20, v/v) at a concentration of 19 mg/mL, filtered through a 0.22 μ m nylon filter (Millipore), and fractionated by preparative HPLC using a Shimadzu LC-8A chromatograph equipped with an SCL-10Avp controller and an SPD-M10Avp diode array detector. The compounds were separated on a Luna C-18 chromatographic column (5 μ m, 150 \times 21.5 mm, Phenomenex) using isocratic elution with an 80:20 (v/v) mixture of water/acetonitrile containing 0.1% (v/v) formic acid. The injection volume and flow rate were 0.4 mL and 10 mL/min, respectively. Under these conditions, 15 compounds were isolated and putative identification was achieved for ten of them (Table 1) using high resolution mass spectrometry (HRMS) and MS/MS in negative mode (micrOTOF-Q II, Bruker Daltonics). For five of the substances (1–5), structural elucidation was carried out by means of ^1H , ^{13}C , and 2D NMR analyses (Bruker Advance III HD 600, 14.7T), in order to confirm the putative identities and the glycosylation

Table 1
Identification of the main compounds in *C. sylvestris* var. *lingua* by HRMS and MS/MS in negative mode.

Peak ^a (#)	Compound name	RRt (min)	UV max (nm)	Negative ionization [M-H] ⁻ (m/z)		Exact mass (experimental)	Exact mass (calculated)	Error (ppm)
				HRMS	MS/MS			
1	(+)-Catechin	2.6	279	289.0729	30 eV: 289 \rightarrow 245; 221; 205; 203; 187; 179; 165	290.0802	290.0790	4.137
A	Isorhamnetin-3-O-trihexoside	9.4	254 (265 sh);	769.2203	55 eV: 769 \rightarrow 314 (315)	770.2276	770.2270	0.779
B ₁	Quercetin-3-O-dihexoside		353	579.1389	40 eV: 579 \rightarrow 300 (301)	580.1462	580.1428	5.861
B ₂	Kaempferol-3-O-dihexoside	9.6	256 (265 sh);	593.1537	50 eV: 593 \rightarrow 284 (285)	594.1610	594.1584	4.376
B ₃	Quercetin-3-O-hexoside		353	463.0907	30 eV: 463 \rightarrow 300 (301)	464.0980	464.0955	5.387
2	Quercetin-3-O-rutinoside (rutin)	10.0	256 (265 sh);	609.1486	45 eV: 609 \rightarrow 300 (301)	610.1559	610.1534	4.097
			353					
3	Isorhamnetin-3-O-neohesperidoside	10.6	254 (265 sh);	623.1646	45 eV: 623 \rightarrow 314 (315)	624.1719	624.1690	4.646
			354					
4	Isorhamnetin-3-O-rutinoside (narcissin)	11.5	255 (267 sh);	623.1665	50 eV: 623 \rightarrow 314 (315)	624.1738	624.1690	7.690
			354					
C	Quercetin-3-O-hexoside	12.0	256 (267 sh);	433.0811	35 eV: 433 \rightarrow 300 (301)	434.0884	434.0849	8.063
D ₁	Isorhamnetin-3-O-hexoside		354	477.1076	45 eV: 477 \rightarrow 314 (315)	478.1149	478.1111	7.948
D ₂	Isorhamnetin-3-O-[rhamnopyranosyl-3-hydroxy-3-methylglutaryl]-hexoside	12.4	254 (267 sh);	767.2065	30 eV: 767 \rightarrow 623 ([M-H-144] ⁻); 314 (315)	768.2138	768.2113	1.953
			352					
D ₃	Kaempferol-3-O-dihexoside			563.1448	35 eV: 563 \rightarrow 284 (285)	564.1521	564.1479	7.445
5	Isorhamnetin-3-O- α -L-rhamnopyranosyl-(1 \rightarrow 2)- α -L-arabinopyranoside	13.2	254 (265 sh);	593.1553	40 eV: 593 \rightarrow 314 (315)	594.1626	594.1585	6.901
			353					
E	Isorhamnetin-3-O-[3-hydroxy-3-methylglutaryl]-hexoside	13.7	254 (267 sh);	621.1502	55 eV: 621 \rightarrow 477 ([M-H-144] ⁻); 314 (315)	622.1575	622.1534	6.590
			351					
F	Isorhamnetin-3-O-pentoside	14.1	255 (265);	447.0975	40 eV: 447 \rightarrow 314 (315)	448.1048	448.1006	9.373
			351					

^a Compounds B₁, B₂, B₃, D₁, D₂, and D₃ were identified in mixtures. For compounds D₂ and E, the neutral loss of 144 corresponds to a 3-hydroxy-3-methylglutaryl moiety. Compounds 1–5 were confirmed by ^1H , ^{13}C , and 2D NMR spectroscopy.

Download English Version:

<https://daneshyari.com/en/article/1355124>

Download Persian Version:

<https://daneshyari.com/article/1355124>

[Daneshyari.com](https://daneshyari.com)