



## Sequence selective naked-eye detection of DNA harnessing extension of oligonucleotide-modified nucleotides



Daniela Verga, Moritz Welter, Andreas Marx\*

Department of Chemistry and Konstanz Research School Chemical Biology University of Konstanz, Universitätsstrasse 10, 78457 Konstanz, Germany

### ARTICLE INFO

#### Article history:

Received 7 December 2015

Revised 22 December 2015

Accepted 23 December 2015

Available online 24 December 2015

#### Keywords:

DNA polymerase

Oligonucleotide-modified nucleotide

Rolling circle amplification

Single-base resolution

Naked-eye detection

### ABSTRACT

DNA polymerases can efficiently and sequence selectively incorporate oligonucleotide (ODN)-modified nucleotides and the incorporated oligonucleotide strand can be employed as primer in rolling circle amplification (RCA). The effective amplification of the DNA primer by  $\Phi$ 29 DNA polymerase allows the sequence-selective hybridisation of the amplified strand with a G-quadruplex DNA sequence that has horse radish peroxidase-like activity. Based on these findings we develop a system that allows DNA detection with single-base resolution by naked eye.

© 2016 Elsevier Ltd. All rights reserved.

The intrinsic importance of nucleic acids as biological targets, due to their connection to gene mutation and expression, protein synthesis, heredity, and pathogens, prompted the development of signal amplification strategies to manufacture highly sensitive DNA biosensors for sequence-specific DNA detection.<sup>1</sup> The detection of low abundant target DNA plays essential roles in early-stage clinical diagnosis and medical treatments for some major diseases like cancer,<sup>2</sup> environmental monitoring<sup>3</sup> as well as forensic analysis.<sup>4</sup> Particularly, point-of-care testing (POCT) for nucleic acid detection is highly demanded where suitable facilities are not available and rapid answers are required.<sup>5</sup> During the past years, PCR-based technologies have been vastly exploited for this purpose.<sup>6</sup> However, they lack the essential features for device portability due to their need of rigid control of temperature cycling using advanced and expensive instrumentation and their request of highly trained analyst.<sup>6a,7</sup> In order to replace the laborious thermal cycling protocol, isothermal amplification methods have been developed. These methods include loop-mediated isothermal amplification (LAMP),<sup>8</sup> strand-displacement amplification (SDA),<sup>9</sup> rolling circle amplification (RCA),<sup>10</sup> and helicase-dependent amplification (HDA).<sup>11</sup>

So far, several naked-eye detection systems have been exploited for nucleic acids and pathogen identification, such as loop-mediated isothermal amplification devices,<sup>12</sup> solid-phase PCR microplate chip assay,<sup>13</sup> urease/DNAzyme coupled system,<sup>14</sup> rolling


circle amplification coupled to a DNAzyme,<sup>15</sup> and DNA probe-modified gold nanoparticles.<sup>16</sup>

Here, we describe a detection system that connects the sequence-selective incorporation of an oligonucleotide (ODN)-modified nucleotide to a colour generation that can be detected by the naked eye. This approach takes advantage from our previous findings that demonstrate the ability of DNA polymerases to incorporate triphosphates that are modified with bulky structured<sup>17</sup> and unstructured<sup>18</sup> single-stranded oligonucleotides, and to use those ODN-modified nucleotides as primers for rolling circle amplification.<sup>18</sup> In here, the elongated DNA strand is captured with a complementary DNA strand that harbours a DNAzyme able to generate a colour signal. Thereby the sequence-selective incorporation of an ODN-modified nucleotide is linked to a signal that is detectable by the naked eye.

The conceptual framework of our approach is depicted in [Figure 1](#). The approach is based on performing a primer extension reaction using an immobilized primer. Hybridization of the primer to a complementary DNA template induces the sequence selective incorporation of the ODN-modified nucleotide. After washing, the immobilized ODN serves as a primer in rolling circle amplification using a complementary circular template resulting in an extended DNA single strand of repeated sequences. The extended single strand product captures the complementary 3'-flanking region of a G-quadruplex-containing DNA. The G-quadruplex-containing DNA has horse radish peroxidase-like activity and thus catalyses the hemin-mediated oxidation of ABTS<sup>2-</sup> by H<sub>2</sub>O<sub>2</sub> to the corresponding radical ABTS<sup>•-</sup> which results in a green colour.<sup>19</sup>

\* Corresponding author.

E-mail address: [andreas.marx@uni-konstanz.de](mailto:andreas.marx@uni-konstanz.de) (A. Marx).




**Figure 1.** Schematic concept of the assay: Primers are immobilized on beads. The hybridisation of the primer to the target DNA template allows the sequence selective incorporation of the ODN-modified nucleotide ( $dT^*TP$ ). After washing steps, the isothermal amplification is performed in the presence of a circular template that binds to its complementary ODN strand and enables the extension of the complementary primer by multiple copies. For signal generation, the beads are incubated with the G-quadruplex DNAzyme sequence, and after washing steps the addition of hemin,  $ABTS^{2-}$ , and  $H_2O_2$  will induce the signal generation only in cases where the ODN-modified nucleotide was incorporated.

The proficiency of DNA polymerases to process  $dT^*TP$  in primer extension experiments were carried out as previously reported.<sup>18</sup> We evaluated the acceptance of  $dT^*TP$  in DNA polymerase promoted primer extension reactions catalysed by the Klenow Fragment of *Escherichia coli* DNA polymerase I ( $KF_{exo^-}$ ) (Fig. S2) and by *KlenTaq* DNA polymerase (Fig. S3). In accordance with previous results, both enzymes process the modified  $dT^*TP$  and incorporate the modified nucleotide in the nascent DNA strand. These results prompted us to explore this system for DNA detection at single-base resolution and facilitating naked-eye detection following the strategy of Figure 1. We chose to conduct experiments on the sequence context of the B type Raf kinase (*BRAF*) gene<sup>20</sup> to be able to compare the results with those previously obtained by incorporation of a nucleotide modified with a G-quadruplex-derived DNAzyme.<sup>17</sup>

For the development of a DNA detection system, a biotinylated primer probe (**P**) was immobilized on streptavidin-coated beads. Next, single incorporation of an ODN-modified nucleotide was performed by usage of  $dT^*TP$  and a template coding for the insertion of a thymidine (Fig. 2A). Therefore, the beads bearing the immobilized primer were incubated at 37 °C for 30 min in the presence of  $KF_{exo^-}$ , template  $T_A$  and  $dT^*TP$  (for details see Supporting information). After incubation, the beads were washed and subsequently incubated with a G-quadruplex DNA strand (**G4**), harbouring a 3'-flanking sequence that binds to the ODN-functionalized nucleotide. The beads were washed and subsequently hemin,  $ABTS^{2-}$ , and  $H_2O_2$  were added to the reaction mixture. The results are depicted in Figure 2B. The chromogenic reaction proceeds over time and stationary read out was observed after 1 h as it has been confirmed by UV measurements of  $ABTS^{2-}$  absorption at 420 nm in function of the time (Fig. 2C). As control a reaction in the absence of  $dT^*TP$  was performed, and no significant colour change was detected. We observed a detection limit at 50 nM (Fig. 2B) that is comparable to the value obtained with the previous developed assay that comprise the direct incorporation of a G-quadruplex-modified nucleotide.<sup>17</sup>

Next, we evaluated the ability of the system to discriminate between the presence of two DNA strands differing only at one nucleotide that is localised at the incorporation site. These experiments were conducted as described before either in the presence of



**Figure 2.** (A) Same as Figure 1 without RCA. (B) Detection limit identification by processing of  $dT^*TP$  with  $KF_{exo^-}$  DNA polymerase at 37 °C in the presence of different concentrations of template ( $T_A$ ). Signal identification via peroxidase-like activity of the complex hemin/hybridised G-quadruplex DNA strand. Concentration template  $T_A$ : test tube I: 400 nM; test tube II: 100 nM; test tube III: 50 nM; test tube IV: 10 nM. Control experiment: test tube C: 400 nM template  $T_A$ , experiment performed without  $dT^*TP$ . The pictures were taken after 1 h. (C)  $ABTS^{2-}$  absorption at 420 nm recorded in function of the time after addition of hemin,  $ABTS^{2-}$ , and  $H_2O_2$  in *caco*. KTD buffer 10 mM to the beads. The time points were taken after: 5, 15, 30, 45, 60, and 90 min. Concentration canonical template ( $T_A$ ): 400 nM (I, 12 pmol, black line), 100 nM (II, 3 pmol, red line), 50 nM (III, 1.5 pmol, blue line), 10 nM (IV, 0.3 pmol, magenta line), and control (C, green line). (D) Partial DNA sequence of primer and template. Discrimination experiments: template  $T_A$ /template  $T_T$  (sequences details see Supporting information). Template  $T_A$  concentrations: test tubes Ia (400 nM), IIa (100 nM), and IIIa (50 nM); template  $T_T$ : test tubes Ib (400 nM), IIb (100 nM), and IIIb (50 nM). The pictures were taken after 1 h.

template  $T_A$ , calling for the insertion of a thymidine analogue, or template  $T_T$ , calling for the incorporation of an adenosine analogue (Fig. 2D). As expected, little colour change was detected in the presence of template  $T_T$  while in the presence of the canonical template  $T_A$  a significant colour change was observed.

Next, we exploited the ability of DNA polymerases to utilise the incorporated ODN-modified nucleotide as a primer in RCA. Thereby the primer should be extended by multiple repeats of a sequence that binds to the G-quadruplex DNA strand (**G4**) that has DNAzyme activity, via its 3'-flanking sequence. This should in turn result in signal amplification. For this purpose, we performed primer extension reactions with natural dNTPs in the presence of DNA polymerase and a circular template (**cT**, see Supporting information) that is complementary to the ODN of the ODN-modified nucleotide. The optimal settings for these experiments that include RCA were obtained by optimising the conditions of the single nucleotide incorporation reaction and by exploring

Download English Version:

<https://daneshyari.com/en/article/1370069>

Download Persian Version:

<https://daneshyari.com/article/1370069>

[Daneshyari.com](https://daneshyari.com)