

A FRET-based assay for screening SIRT5 specific modulators



Yan Li^a, Wenfei Huang^{b,c}, Ling You^{b,c}, Ting Xie^{b,c}, Bin He^{b,c,*}

^a College of Basic Medical Science, Guiyang Medical University, Guiyang, Guizhou 550004, China

^b College of Pharmacy, Guiyang Medical University, Guiyang, Guizhou 550004, China

^c Engineering Research Center for the Development and Application of Ethnic Medicine and TCM, Ministry of Education, Guiyang Medical University, Guiyang, Guizhou 550004, China

ARTICLE INFO

Article history:

Received 10 December 2014

Revised 14 February 2015

Accepted 7 March 2015

Available online 14 March 2015

Keywords:

Sirtuin

Deacetylation

Desuccinylation

Fluorogenic assay

FRET assay

ABSTRACT

A fluorogenic assay for SIRT5 has been developed to screen their small molecule modulators based on the recent discovery that SIRT5 is a demalonylase and desuccinylase. However, this assay uses a fluorogenic peptide containing 7-amino-4-methylcoumarin (AMC), which becomes the cause of false positive hits from the screening. To overcome this, we have developed an alternative method called a FRET-based assay, which will be reliable and useful for screening SIRT5 modulators in a high-throughput format since no AMC group present in this assay.

© 2015 Elsevier Ltd. All rights reserved.

As Class III of histone deacetylases, sirtuins are originally known as a class of enzymes to remove posttranslational modification of acetyl group from lysine residue.^{1,2} Different from Class I/II/IV of histone deacetylases, sirtuins carry out the deacetylation by using nicotinamide adenine dinucleotide (NAD) as a cofactor instead of Zinc (Fig. 1).² There are seven human sirtuins (SIRT1–7) that present diversity in cellular localization and function (Fig. 1).³ Sirtuins involve many important biological functions, including life span, transcription, genome stability, metabolism and protein secretion.^{4–6} Therefore, sirtuins as important drug targets have been extensively investigated. Small molecules that can modulate sirtuin activity have been shown to have potential for treating many human diseases, such as obesity, diabetes, inflammation, cancer, cardiovascular, neurodegenerative diseases and other ageing-related diseases.^{7,8} For examples, SIRT1 inhibitors can induce cancer cell growth arrest and thus promote apoptosis.^{9,10} SIRT2 inhibitors show beneficial effects in Parkinson's disease.¹¹ On the other hand, sirtuin activators have the potentials for treating diabetes¹² and promoting longevity.¹³ However, there are still some controversies on the effects of sirtuin activators.^{14,15}

To discover small molecules that can regulate sirtuin activity, high-throughput assays have been developed by the detection of either deacetylated peptide formation or nicotinamide formation using a coupled enzymatic assay. For the detection of nicotinamide formation, nicotinamidase and glutamate dehydrogenase as

coupling enzymes have been used not only in sirtuin activity assay but also in other assays for nicotinamide-forming enzymes.¹⁶ On the other hand, the detection of deacetylated peptide formation become the most common strategy for sirtuins' high-throughput assays. Among these, one is called a fluorogenic assay that couples the deacetylation of sirtuins to the trypsin-catalyzed amide bond hydrolysis to release a fluorescent small molecule, 7-amino-4-methylcoumarin (AMC, Fig. 2).¹⁷ However, the use of this substrate containing AMC molecule has previously become the cause of false positive hits and thus bring the controversial on the effects of sirtuin activators (resveratrol and its analogues).^{14,15} Not only AMC itself, but also other large aromatic groups would probably cause similar issues due to a hydrophobic binding pocket for aromatic groups.¹⁸ The other method is a fluorescence resonance energy transfer (FRET)-based assay where a donor dye and a quencher dye are connected to an acetyl peptide substrate.¹⁹ A FRET effect will occur in this acetylated peptide, which will cause the fluorescence of a donor dye quenched by a quencher dye. The deacetylation of sirtuins followed by trypsin digestion disrupts the FRET signal and thus releases the fluorescence from a donor dye (Fig. 3).¹⁹

All fluorogenic or FRET-based assays have been developed previously only for SIRT1, SIRT2 and SIRT3 so far, three sirtuins with high deacetylase activity. Until recent discoveries that SIRT5 is a demalonylase and desuccinylase (Fig. 1)²⁰ while SIRT6 is a defatty-acylase (removing long chain fatty acyl groups) (Fig. 1)⁴, a fluorogenic assay for SIRT5 (Fig. 2) as well as for SIRT6 (Fig. 2) have been developed using a fluorogenic AMC-succinyl peptide

* Corresponding author. Tel.: +86 851 690 8899; fax: +86 851 690 8468.

E-mail address: binhe@gmc.edu.cn (B. He).




Figure 1. The deacetylation reactions catalyzed by different sirtuins.

and AMC-myristoyl peptide, respectively.^{21,22} As we discuss above, AMC molecule became an issue of accuracy during the screening.^{14,15} Additionally, AMC molecule has to be installed at the C-terminal of a fluorogenic peptide, which will cause less binding affinity to sirtuin. Therefore, an alternative assay for these sirtuins still needs to be further developed.

Here we report that an alternative method called a FRET-based assay for SIRT5 has been developed, which can be used to screen for SIRT5 inhibitors. We also demonstrate that combined with a secondary screen of SIRT1/2/3, we can identify inhibitors that are selective or not for SIRT5.

The design of a FRET-based assay of SIRT5 is a succinyl peptide containing a pair of donor dye and quencher dye at the end of each terminal, respectively (Fig. 3). Compared to other FRET pairs, a FRET pair of 4-(dimethylaminoazo)benzene-4-carboxylic acid/5-((2-Aminoethyl)aminonaphthalene-1-sulfonic acid, DABCYL/EDANS exhibits better sensitivity at E_x/E_m (maximum excitation and emission) = 340 nm/490 nm.²³ Therefore, we chose DABCYL/EDANS as a pair of donor dye and quencher dye (structures shown in Fig. S1, in the Supporting information). The synthetic route to

this FRET peptide is following the standard procedure of solid phase peptide synthesis shown in the Supporting information (SI). Because one of the known succinylated proteins is glutamate dehydrogenase,²⁰ we made a FRET peptide based on the sequence of the glutamate dehydrogenase containing the succinyl lysine residue, (DABCYL)ISGASE(SuK)DIVHSE(EDANS)G (Fig. 3), where SuK stands for succinyl lysine. For controls, we also synthesized the corresponding peptides with acetyl lysine and free lysine, (DABCYL)ISGASE(AcK)DIVHSE(EDANS)G and (DABCYL)ISGASEKDIDIVHSE(EDANS)G.

With these FRET peptides in hand, we first used (DABCYL)ISGASEKDIDIVHSE(EDANS)G to check whether it could be efficiently digested by trypsin to disrupt the FRET signal and then give the fluorescence. The results showed that 6.25 U trypsin could digest the peptide very efficiently and give about 15 times higher fluorescence compared to the control without trypsin in a one-hour reaction containing 10 μ M (DABCYL)ISGASEKDIDIVHSE(EDANS)G (Fig. S2, SI).

We then determined the kinetic constants of SIRT5 on the (DABCYL)ISGASE(SuK)DIVHSE(EDANS)G peptide versus (DABCYL)ISGASE(AcK)DIVHSE(EDANS)G peptide. Using an HPLC assay, we measured the initial rate velocities as a function of substrate concentration and fit the data to the Michaelis–Menten equation to give the K_m and k_{cat} values as shown in Table 1. The catalytic efficiency (k_{cat}/K_m) for the succinyl peptide was much higher than that for the acetyl peptide (Table 1), which is consistent with the results from the non-fluorophore containing peptide.²⁰ This result demonstrate that (DABCYL)ISGASE(SuK)DIVHSE(EDANS)G peptide is a substrate for SIRT5, a desuccinylase.

Encouraged by this, we next test the (DABCYL)ISGASE(SuK)DIVHSE(EDANS)G peptide by coupled the SIRT5 enzymatic reaction with trypsin digestion. In brief, the assay was carried out in one-hour incubation with 1 μ M SIRT5 and 10 μ M peptide followed by one-hour incubation with 6.25U trypsin. As shown in Figure 4, the fluorescence was increased over 10-fold compared with the control without SIRT5. In contrast, no fluorescence increase was observed when SIRT1, 2 or 3 was used instead of SIRT5 (Fig. 4). The fluorescence also show a dose-dependent manner against the concentrations of SIRT5 (Fig. S3, SI). These results demonstrate that the (DABCYL)ISGASE(SuK)DIVHSE(EDANS)G peptide is a suitable FRET substrate for SIRT5 activity assay.

We next tested whether this FRET-based assay could be utilized to pick up compounds that can modulate SIRT5 activity. We chose several compounds known to be sirtuin inhibitors, including nicotinamide,²⁴ suramin,²⁵ splitomicin,²⁶ AGK2¹¹ and sirtinol²⁷ (structures shown in Fig. S4, SI). Suramin was reported to inhibit SIRT5's deacetylase activity with IC₅₀ value of 22 μ M.²⁵ We measured its IC₅₀ value for the desuccinylase activity of SIRT5 using an HPLC-based assay and obtained a similar value of 25 μ M.²⁸ Other compounds are not efficient at inhibiting SIRT5, with IC₅₀ values >100 μ M.²⁸ Using the concentration of 30 μ M, we screened




Figure 2. Fluorogenic assays for different sirtuins.




Figure 3. A FRET-based assay of SIRT5 using a FRET pair of DABCYL and EDANS.

Download English Version:

<https://daneshyari.com/en/article/1371018>

Download Persian Version:

<https://daneshyari.com/article/1371018>

[Daneshyari.com](https://daneshyari.com)