



Discovery of a novel selective PPAR γ modulator from (–)-Cercosporamide derivatives

Akihiro Furukawa^{a,*}, Tsuyoshi Arita^a, Susumu Satoh^b, Kenji Wakabayashi^c, Shinko Hayashi^a, Yumi Matsui^a, Kazushi Araki^a, Masanori Kuroha^a, Jun Ohsumi^a

^aShinagawa R&D Center, Daiichi Sankyo Co., Ltd, 1-2-58, Hiromachi, Shinagawa-ku, Tokyo, Japan

^bDaiichi Sankyo RD Associe Co., Ltd, 3-10-2, Kitashinagawa, Shinagawa-ku, Tokyo, Japan

^cKasai R&D Center, Daiichi Sankyo Co., Ltd, 1-16-13, Kitakasai, Edogawa-ku, Tokyo, Japan

ARTICLE INFO

Article history:

Received 4 December 2009

Revised 15 February 2010

Accepted 17 February 2010

Available online 20 February 2010

Keywords:

Cercosporamide

Diabetes

PPAR

Modulator

ABSTRACT

In an investigation of (–)-Cercosporamide derivatives with a plasma glucose-lowering effect, we found that *N*-benzylcarboxamide derivative **4** was a partial agonist of PPAR γ . A SAR study of the substituents on carboxamide nitrogen afforded the *N*-(1-naphthyl)methylcarboxamide derivative **23** as the most potent selective PPAR γ modulator. An X-ray crystallography study revealed that compound **23** bounded to the PPAR γ ligand binding domain in a unique way without any interaction with helix12. Compound **23** displayed a potent plasma glucose-lowering effect in db/db mice without the undesirable increase in body fluid and heart weight that is typically observed when PPAR γ full agonists are administrated.

© 2010 Elsevier Ltd. All rights reserved.

Type 2 diabetes is one of the most serious health problems in the world. The World Health Organization (WHO) estimates that more than 180 million people worldwide have diabetes. This number is likely to more than double by 2030.¹ Although many therapeutic agents have already been used in clinical situations, it is still difficult to tightly control plasma glucose and prevent diabetic complications.^{2,3} So there is great need for novel pharmacotherapy which is able to achieve tight glycemic control singly or to be used with existing agents.

We have previously reported that (–)-Cercosporamide had a plasma glucose-lowering effect in KK/Ta mice, but it was accompanied by a severe decrease in food intake and a loss of body weight.⁴ In that report it was also shown that *N,O*-acetal type derivatives **1** and **2** (Fig. 1) lowered the plasma glucose in KK/Ta mice without notably affecting the food consumption or body weight. Although they showed an obvious and promising effect on hyperglycemia, the mechanism of action had not yet been made clear.

We designed benzylated compounds **3** and **4** to enhance the desirable effect because the fact that compound **2** was superior to **1** indicated that the phenyl ring near the carbamoyl group was important for the plasma glucose-lowering effect. The preparation of these compounds was accomplished in the following manner (Scheme 1). For the preparation of compound **3**, the carbamoyl nitrogen and 3-hydroxyl group of (–)-Cercosporamide were re-

acted with 2,2-dimethoxypropane to form an acetonide. The carbamoyl nitrogen and 1-hydroxyl group were benzylated by benzylbromide using sodium hydroxide as a base. The benzyl group on the 1-hydroxy group was selectively deprotected by Pd/C catalyzed hydrogenolysis to give compound **3**. For the preparation of compound **4**, the 3-hydroxyl group of (–)-Cercosporamide was selectively methylated by iodomethane in the presence of potassium carbonate. The carbamoyl nitrogen was benzylated to compound **4** by reductive amination with benzaldehyde, triethylsilane and trifluoroacetic acid in toluene.⁵

The plasma glucose-lowering effects of these compounds were tested in hyperglycemic KK/Ta mice. The test compounds were mixed with their diet in a ratio of 0.1% (about 100 mg/kg/day if the food intake did not change). This mixture was fed to the mice





Figure 1. Previously reported (–)-Cercosporamide derivatives.

* Corresponding author. Tel.: +81 3 3492 3131; fax: +81 3 5436 8563.

E-mail address: furukawa.akihiro.zy@daiichisankyo.co.jp (A. Furukawa).



Scheme 1. Reagents and conditions: (a) 2,2-dimethoxypropane, TsOH, acetone, reflux, 8 h, 80%; (b) BnBr, NaH, DMF 0–25 °C, 2 h, 79%; (c) Pd/C, EtOAc/EtOH, 6 h, 41%; (d) MeI, K₂CO₃, DMF, 25 °C, 24 h, 66%; (e) benzaldehyde, Et₃SiH, TFA, toluene, reflux, 8 h, 74%.

Table 1
Plasma glucose-lowering effect in KK/Ta mice

Compound number	Plasma glucose correction ^a (%)	Body Weight on 7th day ^b (%)	Food intake on 7th day ^b (%)
1	56	102	78
3	55	103	74
4	69	101	100

^a The values are the % change in the plasma glucose concentration of the drug-treated mice relative to the vehicle controls. All the values are the means of 4 or 6 mice.


^b The values are the % degree of the drug-treated mice relative to the vehicle controls.

for a week. The plasma glucose, body weight, and food intake data are summarized in Table 1. Compound 4 showed the most potent glucose-lowering effect and did not affect the body weight or food intake at all. On the other hand, compounds 1 and 3 slightly induced the decrease in food intake.

We were strongly interested in the mechanism of action of promising compound 4. We thoroughly investigated the interaction between compound 4 and various targets which were related to diabetes. Finally, we found that compound 4 partially activated a peroxisome proliferator-activated receptor gamma (PPAR γ (EC₅₀: 0.22 μ M, E_{\max} : 44%). PPAR γ agonists promote adipocyte differentiation and improve insulin resistance.⁶ The PPAR γ full agonists Pioglitazone and Rosiglitazone have been clinically used and show beneficial effects in patients with type 2 diabetes. However, the use of these drugs has been limited because of their adverse effects, such as weight gain, edema, and anemia.⁷ Recently, it was proposed that modulating PPAR γ activity, rather than activating it fully, was effective in improving insulin resistance without undesirable adverse effects.⁸ A number of compounds, such as Fmoc-L-Leucine,⁹ FK614,¹⁰ 5-substituted 2-benzoylaminobenzoic acids,¹¹ indole and azaindole derivatives,¹² T2384,¹³ INT-131,¹⁴ and MBX-102,¹⁵ are reported to modulate PPAR γ transactivation and improve insulin resistance in hyperglycemic rodent models. In some cases, it was observed that the adverse effects which accompany PPAR γ activation were attenuated.

Accordingly, we embarked on a SAR study to acquire a more potent PPAR γ modulator than compound 4. First, we incorporated various substituents on the phenyl ring. Next, we changed the phenyl ring to a heteroaryl ring or a benzo-fused aryl ring. All these compounds were synthesized in the same way as compound 4 with the corresponding aromatic aldehyde in toluene or acetonitrile. We tested the PPAR γ transactivation activities of these compounds. Then, for any compounds which showed potent transcriptional activities (EC₅₀ < 1 μ M), we tested the potency of their effect as partial antagonists. Their structures and transcriptional activities are summarized in Table 2. In the various substituents on the phenyl ring, only chloride substitution on the *ortho*- or *para*-position (compounds 11 and 13) sustained transactivation activity. In the heteroaryl or benzo-fused compounds, 2-furyl compound 16 and 1-naphthyl compound 23 had the potent PPAR γ transactivation activity equal to that of compound 4. As well compound 23 acted as a potent partial antagonist in comparison to compounds 4 and 16. Then we tested the binding affinities to

Table 2
PPAR γ transactivation



Compound number	Ar	Transactivation			
		EC ₅₀ ^a (μ M)	E_{\max} ^b (%)	IC ₅₀ ^c (μ M)	I_{\max} ^d (%)
4	Phenyl	0.22	44	ND ^e	
5	<i>o</i> -MeO-phenyl	ND ^e			
6	<i>m</i> -MeO-phenyl	3.8	36		
7	<i>p</i> -MeO-phenyl	1.7	43		
8	<i>o</i> -NO ₂ -phenyl	1.6	42		
9	<i>m</i> -NO ₂ -phenyl	2.4	34		
10	<i>p</i> -NO ₂ -phenyl	6.0	46		
11	<i>o</i> -Cl-phenyl	0.28	19	ND ^e	
12	<i>m</i> -Cl-phenyl	6.7	56		
13	<i>p</i> -Cl-phenyl	0.77	50	Negative	
14	2-Pyridinyl	Negative			
15	3-Pyridinyl	11	51		
16	2-Furyl	0.56	34	13	70
17	3-Furyl	3.2	54		
18	2-Thiophenyl	1.2	40		
19	3-Thiophenyl	4.4	45		
20	2-Benzofuryl	2.3	93		
21	2-Benzothiophenyl	3.8	77		
22	3-Benzothiophenyl	ND ^e			
23	1-Naphthyl	0.18	47	0.62	45
24	2-Naphthyl	ND ^e			
Pioglitazone		0.088	118	Negative	
Rosiglitazone		0.011	106	Negative	

^a EC₅₀ was defined as the compound concentration at which 50% of a given compound's intrinsic maximal response had been reached.

^b The transcriptional activity in the presence of an in-house potent PPAR γ full agonist (3.3 nM) was defined as 100%, while that in the vehicle alone was defined as zero. The maximum transcriptional activity in the presence of the test compound was defined as E_{\max} (%).

^c IC₅₀ was defined as the compound concentration at which 50% of a given compound's intrinsic maximal inhibition had been reached.

^d The maximum inhibition of the test compound against the transcriptional activity in the presence of an in-house potent PPAR γ full agonist (3.3 nM) was defined as I_{\max} (%).

^e ND means 'not determined' because the E_{\max} or I_{\max} was too low.

Download English Version:

<https://daneshyari.com/en/article/1374532>

Download Persian Version:

<https://daneshyari.com/article/1374532>

[Daneshyari.com](https://daneshyari.com)