



Tuning the pH sensitivities of orthoester based compounds for drug delivery applications by simple chemical modification

Hélène Bruyère, Andrew D. Westwell, Arwyn T. Jones *

Welsh School of Pharmacy, Cardiff University, Redwood Building, Cardiff CF10 3NB, Wales, UK

ARTICLE INFO

Article history:

Received 16 December 2009

Revised 8 February 2010

Accepted 8 February 2010

Available online 13 February 2010

Keywords:

Orthoester

pH sensitive linkers

Drug delivery

ABSTRACT

Orthoesters are acid-sensitive moieties that allow substantial structural diversity for biological applications including drug delivery. Here, the pH-sensitivity of a range of novel orthoester based compounds was compared in the range 7.5–4.5 that is characteristic of the increased acidification during endocytosis. We find that simple modifications close to the orthoester had major effects on both the rate and extent of hydrolysis, suggesting this could be exploited for activating drug delivery systems on endocytic pathways.

© 2010 Elsevier Ltd. All rights reserved.

Polymer based architectures have been extensively studied as backbones for drug delivery systems.^{1,2} The strategy is often based on their conjugation to a therapeutic cargo that is then delivered to the inside of cells by endocytosis.^{3–5} There is often a necessity for the therapeutic to be released from the polymer backbone for it to then reach the cytosol and possibly other intracellular targets. An approach to this design is to use the considerable drop in pH that molecules encounter as they are trafficked through endocytic pathways.⁶ These are used by cells to internalize molecules that will then encounter a drop from extracellular pH 7.3–7.5 to 4.5–5.0 in lysosomes.⁷ These organelles and to a lesser extent late endosomes constitute a strong hydrolytic environment, thus there may be a requirement to release the therapeutic early during endocytosis (pH 5–6) if it is labile to these conditions.

Several pH sensitive linkers have been studied in this context, including hydrazones,⁸ *cis*-aconityl,⁹ vinyl ethers,¹⁰ and acetals.¹¹ Also included here are orthoesters, and to date these are shown to be one of the most acid-labile functionalities available.¹² As noted, drug delivery systems may need to be activated at a specific pH, considerably higher than that encountered in lysosomes. A slightly acidic pH has also been observed in tumours¹³ and inflammatory tissues¹⁴ thus there may also be an interest in designing pH specific systems for mediating effects in these environments. Orthoesters are of particular interest in attempting to achieve this goal because they can be modified with respect to structural patterns or stereochemical, stereoelectronic and steric factors.^{15–17} There are numerous reports describing the behaviour of these in


acidic conditions and examples include poly(orthoesters) and other orthoester- or diorthoester-lipid conjugates, designed for delivery of biologically active cargo through endocytic pathways.^{18–21} Pure chemical and mechanistic studies tend to be performed in organic or mixed organic/aqueous solvents and rarely consider the efficiency of the system in an aqueous environment or with respects to time and temperature. Whereas those interested in drug delivery rarely focus on characterizing a range of different structures.

Specifically towards the design of a polymer-peptide based drug delivery system that is sensitive to acidic pHs found on endocytic pathways,²² we designed, synthesized and measured the rates of hydrolysis of different and novel model compounds containing an orthoester linker. For proof of concept towards the design of polymer-peptide conjugates, selected structures were linked to diethylene glycol as a template polymer for polyethylene glycol (PEG) and phenylalanine as a peptide substitute. Our studies highlight that simple modifications on the orthoester backbone have a major influence on hydrolysis profiles that is then transferable to more complete drug delivery systems. This may allow a high degree of flexibility with respects to pH activation that may, depending on the biological application, be a requirement.

The synthetic template is based on a previous publication²¹ and is outlined in [Scheme 1](#). The amino groups of commercially available amino-diols **1**, **4**, **5** and **6** were protected as their trifluoroacetate derivatives using ethyltrifluoroacetate. Reaction of these protected amino-diols **2** and **7–9** with trimethyl orthoformate under *p*-toluene sulfonic acid (PTSA) catalysis provided the orthoester building blocks (**3** and **10–12**) primed for diethylene glycol and phenylalanine functionalisation. Transesterification using diethyl-

* Corresponding author. Tel.: +44 0 2920 876431.

E-mail address: jonesat@cardiff.ac.uk (A.T. Jones).




Scheme 1. Synthesis of orthoester model compounds **14** and **18–20**. Reagents and conditions: (i) ethyltrifluoroacetate, THF, room temperature; (ii) trimethylorthoformate, PTSA, CH₂Cl₂, room temperature; (iii) diethylene glycol monomethyl ether, PPTS, toluene, reflux; (iv) 1 M NaOH, THF, room temperature; (v) *N*-Ac-L-phenylalanine, EDCl, DMAP, CH₂Cl₂, THF, room temperature; (vi) *N*-Ac-L-phenylalanine, EDCl, CH₂Cl₂, THF, room temperature.

ene glycol monomethyl ether gave the intermediate orthoesters **13** and **15–17**. The trifluoroacetyl protecting group was then removed by the action of aqueous NaOH in THF, and pseudo-peptidic coupling with *N*-acetyl phenylalanine provided the necessary orthoesters **14** and **18–20**. The specified conditions for synthesizing compound **18** were not suitable for generating the other model compounds unless DMAP was omitted as a catalyst. Due to acid-sensitivity and the fact that no starting material was detectable by ¹H NMR after work-up and concentration, the intermediate compounds were not purified. Indeed, attempts at purification after *trans*-etherification did not lead to significant improvements in yield of the final compound. Similarly, purification of the final compounds (**14** and **18–20**) by flash chromatography on silica gels was difficult thus neutral aluminum oxide was used as an alternative. Of note for NMR analysis is that several isomers were identified at different steps as the orthoester cycle is formed. From this, four different orthoester model compounds were prepared (**14**, **18**, **19**, **20**).

Two different hydrolysis pathways have been described for orthoesters depending on whether the first protonation occurs on an exocyclic or endocyclic oxygen atom.²³ The stages are detailed in Scheme 2 using a six-membered ring orthoester as an example. It is expected that any stereoelectronic or stereochemical/conformational effect on a model molecule would influence the nature of the hydrolysis products, their stability and their endocyclic or exocyclic pathway ratio.^{16,24}

In order to directly measure the hydrolysis ratio between the parent orthoester and the hydrolysis products by NMR, we used deuterated solvents to prepare all buffers. These were prepared from a mixture of 0.5 M citric acid and 0.5 M Na₂HPO₄ solutions in deuterium oxide. Two similar ¹H NMR methods were then used to evaluate the stability of the orthoester-containing products under a range of pH values from 7.5 to 4.5. Initially, hydrolysis studies were carried out for 90 min at 37 °C in single tubes inside the NMR spectrometer that was previously equilibrated to this temperature.



Scheme 2. Orthoester hydrolytic pathways.

Alternatively the reactions were performed in a 37 °C water bath and aliquots were removed at different time periods for NMR analysis. To stop further hydrolysis these samples were quenched with 0.5 M Na₂HPO₄ in deuterium oxide (pH 8.5). We confirmed that this completely inhibited any further hydrolysis irrespective of the pH of the initial solution and experiments performed in the laboratory and on the NMR instrument yielded identical data (not shown). Some experiments were performed at room temperature whereby the same reaction tubes were periodically placed on the NMR instrument.

The extent of hydrolysis was determined by relative integration of specific orthoester and newly formed hydrolysis peaks (Fig. 1). Several peaks were visible in each specific area because of the creation of different diastereoisomers as mentioned above. The pH of the hydrolysis mixtures did not change during the reaction

Download English Version:

<https://daneshyari.com/en/article/1374557>

Download Persian Version:

<https://daneshyari.com/article/1374557>

[Daneshyari.com](https://daneshyari.com)