

Synthesis and biological activity of novel peptide mimetics as melanocortin receptor agonists

Xue-Wei Liu,^{a,*} Jimei Ma,^a Anny-Odile Colson,^b Doreen Cross Doersen^b
and Frank H. Ebetino^b

^a*Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences,
Nanyang Technological University, Singapore 637371, Singapore*

^b*Procter & Gamble Pharmaceuticals, Health Care Research Center, 8700 Mason Montgomery Road, Mason, OH 45040, USA*

Received 16 October 2007; revised 25 November 2007; accepted 28 November 2007
Available online 4 December 2007

Abstract—A series of novel peptidomimetic analogs was prepared containing cyclohexyl, phenyl, or heterocyclic groups to ostensibly orient the guanidine or mimic of an arginine in a putative melanocortin receptor ligand pharmacophore. Some binding affinity at the melanocortin receptors MC₃ and MC₄ was noted. In silico docking also indicated that the relative positions of the hydrogen-bonding sites and hydrophobic regions of the compounds are reasonably well matched to the receptor-binding site. This may present a lead entry into a selective series of MC₄R agonists.

© 2007 Elsevier Ltd. All rights reserved.

Melanocortin receptors (MCR) are members of the G-protein coupled receptor subfamily consisting of MC_{1,2,3,4,5}.^{1,2} The MC₁ receptor has been found in melanocytes where it controls skin pigmentation while recent studies indicate that MC₄R is involved in the control of feeding behavior. As a result, the MC₄ receptor has received increased attention as a receptor target from both academia and industry. Since this subtype is localized in the central nervous system, transport of exogenous ligands across the blood-brain barrier remains one of the challenges in this field.^{3,4}

Melanocyte stimulating hormone, α -MSH is a natural ligand for four of the melanocortin receptor subtypes, namely, the MC₁, MC₃, MC₄, and MC₅ receptors. This peptide hormone is 13 amino acids in length and binds with high affinity to the MC₁, MC₃, MC₄, and MC₅ receptors. Other ligands designed earlier shared a common core sequence motif: His-Phe-Arg-Trp.⁵ A number of linear and cyclic peptide and peptidomimetics containing this sequence have now been reported. Most show a similar selectivity profile to the natural MSH.^{6–8} Interestingly, replacement of L-Phe with D-Phe was re-

ported to increase the affinities for the MC receptors.^{9,10} Also, replacement of the His residue was shown to increase selectivity for the MC₄ receptor.^{11,12} It is only recently that subtype selective, orally active, small (MW < 500) ligands were reported. Workers at Merck synthesized a new class of cyclohexyl substituted piperidines that presumably mimicked this HFRW sequence (Fig. 1).¹³ Compound A is a potent (EC₅₀ = 2.1 nM), selective (1184-fold vs. MC₃R, 350-fold vs. MC₅R), small-molecule full agonist of the human MC₄ receptor. In an attempt to better understand this finding and related theories on the minimum pharmacophore required for MC₄ activity, we designed a related series of cyclohexyl-, phenyl-, and heterocyclic-containing linear compounds. These compounds representing the piperidine ring-opened analogs of compound A were thus designed to test the criticality of the conformational restriction aspects of this drug lead. In this letter, we report the synthesis and biological activity of novel analogs based on this concept.

The novel cyclohexyl-directed triazole analog was synthesized according to the route presented in Scheme 1. The commercially available cyclohexyl methyl ketone **1** was brominated with benzyltrimethylammonium tribromide yielding α -bromo ketone **2** in high yield (98%). Overnight reaction of **2** with sodium triazole in DMF at 50 °C yielded the α -triazole ketone **3** (91%) which,

Keywords: Melanocortin receptor; Peptidomimetics; MC₄R agonist; Homology model.

*Corresponding author. Tel.: +65 6316 8901; fax: +65 6791 1961; e-mail: xuewei@ntu.edu.sg





Figure 1. Constrained (A) and linear (B) conformations of the ligand.



Scheme 1. Reagents and conditions: (a) $\text{PhCH}_2\text{N}(\text{CH}_3)_3\text{Br}_3$, CH_2Cl_2 , 5°C –rt, 4 h, 98%; (b) Sodium triazole, DMF, 5°C , 91%; (c) diethyl phosphonoacetonitrile, LiCl, DBU, CH_3CN , rt, 90%; (d) H_2 , Ni, NH_4OH , MeOH, 40 psi, 4 h, 98%; (e) Boc-D-Phe(*p*-F)-OH, EDC, HOBt, NMM, DMF, rt, overnight, 84%; (f) TFA, CH_2Cl_2 , 97%.

via a facile Horner–Wadsworth–Emmons reaction with diethyl phosphonoacetonitrile in the presence of LiCl and DBU, provided α -, β -unsaturated nitrile **4** in 90% yield. The primary amine **5** was obtained in almost quantitative yield by hydrogenation of **4** on a Parr shaker at 40 psi. This amine was coupled with Boc-D-Phe(*p*-F)-OH in the presence of HOBt, EDC, and NMM. The crude product after reaction workup was treated with trifluoroacetic acid and purified by reverse-phase HPLC to generate the final compound **6** as trifluoroacetate salt in a yield of 81% for two steps.

Scheme 2 illustrates the synthesis of dipeptidomimetics **12** from the commercially available *N*-*t*-Boc-L-phenylalanine **7**. The aldehyde **7** was coupled to diethyl phosphonoacetonitrile via a Horner–Wadsworth–Emmons reaction in 89% yield. The resulting α -, β -unsaturated nitrile **8** was reduced to the amine **9** (92%). Rh/alumina was an optimal catalyst for this step to hydrogenate the double bond nitrile and the phenyl ring in one step. The free primary amine **9** was converted to the di-Cbz protected guanidine derivative **10** by guanylation with *N,N*-di-Cbz-*S*-methylpseudourea in the presence of HgCl_2 and Et_3N . After deprotection with 50% TFA in CH_2Cl_2 , **10** was coupled to Boc-D-Phe(*p*-F)-OH to generate **11**, followed by removal of Boc with 50% TFA in CH_2Cl_2 and catalytic hydrogenation to remove the Cbz group. The dipeptidomimetics **12** was obtained after HPLC purification.

Phenyl analog **16** of cyclohexyl compound **12** was prepared as shown in **Scheme 3** starting from α -, β -unsaturated nitrile **8**. Catalyzed by Raney nickel, it was reduced to saturated amine **13** with the phenyl ring intact. The final target **16** was obtained from primary amine **13** via **14**, and **15**, following the procedures used to prepare **12**.

Schemes 4 and **5** outline the methodology for preparing dipeptides D-Phe(*p*-F)-Arg with terminal modifications. Boc-D-Phe(*p*-F)-OH **17** was coupled with the methyl ester of Arg **18** under typical EDC/HOBt coupling conditions, resulting in dipeptide **19**. The terminal methyl ester group was converted to the corresponding acid **20** with LiOH in THF and water (3:1) followed by acidification. EDC/HOBt coupling of the acid **20** and piperidine gave **21**, which was transformed to dipeptide **22** after removal of the Boc group with TFA in CH_2CH_2 and hydrogenolytic removal of the nitro group in structure **21**.

As shown in **Scheme 5**, methyl ester **19** was reduced to alcohol **23** with LiBH_4 in THF at room temperature in a reasonable yield (77%). In the presence of DEAD and Ph_3P , this alcohol reacted with piperidine in a Mitsunobu reaction in a moderate yield (41%), generating dipeptidomimetics **24**. Treatment of **24** with TFA in CH_2CH_2 and hydrogenolysis of the nitro group, followed by HPLC purification, provided the tertiary amine analog **25**.

Download English Version:

<https://daneshyari.com/en/article/1377373>

Download Persian Version:

<https://daneshyari.com/article/1377373>

[Daneshyari.com](https://daneshyari.com)