

Short communication

Cancer chemopreventive activity of sulforamate derivatives

Robert M. Moriarty, Rajesh Naithani *, Jerome Kosmeder, Om Prakash

4500, Department of Chemistry, SES Building, 845, W. Taylor Street, University of Illinois at Chicago, Chicago 60607, IL, USA

Received 9 May 2005; received in revised form 30 September 2005; accepted 6 October 2005

Available online 21 November 2005

Abstract

Chemoprevention can be defined as an intervention in the carcinogenic process by use of natural or synthetic substances. Induction of Phase II enzyme is an important mechanism of chemoprevention. In the present studies we have synthesized several derivatives of (+)(–) 4-methylsulfinyl-1-(*S*-methylthiocarbamyl)-butane (sulforamate) and evaluated their effectiveness as monofunctional inducer of the NAD(P)H Quinone oxidoreductase [quinone reductase (QR)] a phase II enzyme in cultured Hepa1c1c7 murine hepatoma cells. The cytotoxicity of some of the derivatives was strongly reduced in comparison to [(–)-1-isothiocyanato-4(R)-(methylsulfinyl)butane] (sulforaphane). However, the induction potential of these compounds was comparable to sulforaphane. On the basis of these results sulforamate derivatives can be regarded as simple, inexpensive and readily available chemopreventive agents.

© 2005 Elsevier SAS. All rights reserved.

Keywords: Chemoprevention; Phase II enzymes; Sulforamate; Quinone oxidoreductase (QR); Monofunctional inducer

The term “cancer chemoprevention” coined by Sporn [1,2] and coworkers in 1976 has been defined as a strategy for reducing cancer mortality by the prevention, delay or reversal of cancer by ingestion of dietary or pharmaceutical agents capable of mediating the process of carcinogenesis [3]. In recent years cancer prevention by natural products has received considerable attention. Much emphasis has been placed on dietary intake of certain fruits and vegetables that have shown to reduce the risk of developing cancer. Numerous epidemiological studies have shown an association between reduced cancer risk and increased intake of green yellow fresh vegetables [4–6]. Particularly noteworthy is the group of food plant generally falls in the family Cruciferae and genus brassica, i.e. cauliflower, cabbage, brussels sprouts and broccoli [7].

It has already been established that three dietary components brassinin (1, indole based dithiocarbamate), cyclobrassinin (2) spirobrassinin (3) sulforaphane (4, aliphatic isothiocyanate) and 1,2-dithiol-3-thione (5) found in the Cruciferous plants are capable of mediating chemopreventive activity in


animal models [8–10]. Feeding of vegetables induces enzymes of xenobiotic metabolism and thereby accelerates the metabolic disposal of the xenobiotics. The human body's first lines of defense against cancer are the Phase I and Phase II enzymes. The metabolic activation and deactivation of carcinogens is catalyzed by xenobiotic Phase I and II enzymes. By measurement of induction of Phase II detoxifying enzymes, such as quinone reductase (QR), glutathione *S*-transferase, UDP-glucuronosyl transferase and epoxide hydrolase, compounds may be screened for anticarcinogenic activity^{10a}. These enzymes can convert electrophiles (ultimate carcinogens) to less reactive compounds, which can in turn conjugate to sugars, glutathione and sulfates resulting in less toxic and more excretable compounds. Phase I enzymes are promoted by cytochromes P-450 that are mediated by the cytosolic Ah receptor (Aryl hydrocarbon). When ligands bind to the Ah receptors the complex is transported to the nucleus and binds to the xenobiotic responsive elements (XRE) upstream for P-450 genes^{10b}. Phase I enzymes typically carry out oxidation and reduction reactions that make carcinogens more water soluble, however they are capable of activating compounds to electrophilic species, which can damage DNA. These two families of enzyme help

* Corresponding author. Tel.: +1 312 996 9585; fax: +1 312 996 0431.
E-mail address: naithani@uic.edu (R. Naithani).

body protect itself from all types of carcinogens that routinely enter the human body through the diet and the environment.


A key component in the understanding the initial events of carcinogenesis is the reorganization of the fact that many of the carcinogens are not chemically reactive per se but undergo metabolic activation to form electrophilic reactants [10c].

Amount of carcinogens available represents a balance between activating and detoxifying reactions of Phase I and Phase II enzymes, respectively [11–13]. This balance under normal circumstances is genetically controlled but gets modulated by variety of factors like age, hormone, and exposure to drugs.



On the basis of structural similarities of brassinin to compounds like indole-3-carbinol and indole-3-acetonitrile, it was found to induce drug-metabolizing enzymes in cell culture. It was found that Brassinin also induced the activity of P-450 isoenzymes. This particular induction is a risk factor in the activation of carcinogens. Infact indole-3-carbinol was found to induce liver carcinogenesis in rainbow trout when applied to the post initiation stage [14]. Although indole-3-carbinol and related compound demonstrate significant chemopreventive activity, induction of Phase I metabolizing enzymes, is considered as a risk factor in the activation of carcinogens. On the basis of this potential cancer risk of the indole based compounds and relative toxicity of the isothiocyanates a hybrid molecule was conceptualized. Isothiocyanate was replaced by the dithiocarbamate moiety of brassinin thus exchanging the indole ring with methyl sulfinyl butane side chain.

In a program directed towards discovery of the novel chemo preventive agents and to evaluate the potential of test agents to enhance the activity of the QR we currently report the synthesis and activity of six novel sulforamate derivatives. These compounds are aliphatic analogs of Brassinin and show structural similarity to sulforaphane. The methyl group of the



Scheme 1.

dithiocarbamate has been replaced by different groups (aliphatic chain, substituted benzyl) Scheme 1.

We anticipated that during metabolism the ability of these moieties to act as a good leaving group would have effect on the induction potential. Compound **10f** (having *para*-fluoro benzyl group instead of methyl in sulforamate) **10b** and **10c** (having propyl and butyl group, respectively, in place of methyl in sulforamate) have shown excellent activity. The CD (concentration required to double QR induction) and CQ (concentration required to quadruple QR induction) of these compounds are comparable to sulforaphane. In addition to this the reduced toxicity of these compounds make it far better chemopreventive agent than sulforaphane. The chemopreventive index (CI) of these compounds is three to four times that of Sulforaphane (Table 1).

Synthesis of **10** (Scheme 1) was carried out by addition of sodium thiomethoxide to N-(4-bromobutyl)phthalimide that in turn was obtained by treating phthalimide with 1,4, dibromobutane. Subsequent oxidation with hydrogen peroxide resulted in the formation of sulfoxide **8**, which upon addition of hydrazines afforded the amines (**9**).

The amine (**9**) was later treated with triethylamine, carbon disulfide and then iodomethane to yield **10** in good yields.

Table 1

Compound number	Concentration required to double QR induction (CD) (μg/M)	Concentration for 50% inhibition of cell viability (IC50) (μg/M)	CQ concentration required to quadruple QR induction (μg/M)	CI
Sulforaphane	0.06	1.7	0.55	28
10a	0.08	9.9	0.94	123
10b	0.07	7.9	0.24	112
10c	0.2	17.9	1.5	90
10d	0.07	1.5	0.6	21
10e	0.1	2.5	1.4	25
10f	0.05	6.1	0.5	122
10g	0.06	0.8	0.4	13

Download English Version:

<https://daneshyari.com/en/article/1397182>

Download Persian Version:

<https://daneshyari.com/article/1397182>

[Daneshyari.com](https://daneshyari.com)