

## Enzymatic synthesis of optically pure cyanohydrins in microchannels using a crude cell lysate

Kaspar Koch<sup>a</sup>, Rutger J.F. van den Berg<sup>a</sup>, Pieter J. Nieuwland<sup>a</sup>, Roel Wijtmans<sup>a</sup>,  
Marcel G. Wubbolts<sup>b</sup>, Hans E. Schoemaker<sup>b</sup>, Floris P.J.T. Rutjes<sup>a,\*</sup>, Jan C.M. van Hest<sup>a,\*</sup>

<sup>a</sup> Institute for Molecules and Materials, Radboud University Nijmegen, Toernooiveld 1, 6525 ED Nijmegen, The Netherlands

<sup>b</sup> DSM Research, PO Box 18, 6160 MD Geleen, The Netherlands

### Abstract

The synthesis of optically pure cyanohydrins using a crude cell lysate containing the enzyme hydroxynitrile lyase (HNL) was studied in a microreactor in an aqueous–organic biphasic system. Different aldehyde substrates were selected to be converted to their corresponding cyanohydrins. It was successfully demonstrated that this crude cell lysate could readily be applied as a biocatalyst in a microchannel without clogging the channels. The biocatalytic activity toward the different substrates could rapidly be investigated with only small quantities of enzyme needed, compared to batch scale screening. Furthermore, the optimal contact between two immiscible phases in a microreactor resulted in enzymatic reactions with a high initial reaction rate and enantioselectivity, comparable to a batchwise process in which optimized conditions were achieved by vigorous stirring. Thus, performing the selected enzymatic reaction in a microreactor is a facile and cost efficient screening method leading to results which can be directly translated to batchwise processes.

© 2007 Elsevier B.V. All rights reserved.

**Keywords:** Cyanohydrins; Hydroxynitrile lyase; Crude cell lysate; Microreactors

### 1. Introduction

During the last decades, there has been increased interest in using enzymes for asymmetric synthesis and kinetic resolution to obtain pure enantiomers. A family of enzymes that has been extensively studied are the hydroxynitrile lyases (HNLs). These enzymes catalyse enantioselective C–C-bond formation via the addition of HCN to aldehydes or ketones to yield the corresponding optically active cyanohydrins (Scheme 1), which are synthetically versatile building blocks for the synthesis of fine chemicals, pharmaceuticals and agrochemicals. Both (*R*)- and (*S*)-selective HNLs are widely present in nature, and several genes have been cloned and expressed in microorganisms such as *E. coli* and *P. pastoris* [1].


Traditionally, synthesis of enantiopure cyanohydrins catalysed by HNL was performed in aqueous media (single phase system). However, in several cases less satisfactory results were obtained with respect to enantiopurity and conversion. A significant advancement to overcome this problem was developed

by Griengl et al. [2]. They showed that by employing a vigorously stirred biphasic system, cyanohydrins could be prepared in excellent enantiomeric purity and high yield as opposed to the previously described single phase system. Furthermore, the same group concluded that fast formation of a stable emulsion in the reaction mixture was of utmost importance for a successful reaction under these biphasic conditions. In a traditional batch process, such a situation can only be achieved through vigorous stirring. We envisaged that these optimal mixing conditions might well be obtained in suitably designed microchannels.

Various classes of chemical reactions have already been performed in microchannels offering more control over selectivity and suppression of by-product formation due to the high surface-to-volume ratio [3,4]. Furthermore, microreactor technology enables rapid reaction optimisation and screening using relatively small quantities of chemicals, which is of importance for the industry to maintain their rapid development of new chemicals. However, in order for a screening tool to be useful, optimised reaction conditions which have been found using microreactor technology have to be translated to industrial scale processes, which is not trivial since physical properties such as mixing behaviour and mass transport can be markedly different and therefore hamper the comparison between the different scale processes.

\* Corresponding authors. Tel.: +31 24 365 3202; fax: +31 24 365 3393.

E-mail addresses: F.Rutjes@science.ru.nl (F.P.J.T. Rutjes), J.Vanhest@science.ru.nl (J.C.M. van Hest).

Scheme 1. General enzymatic synthesis of (*S*)-cyanohydrins.

Remarkably, the field of enzymatic synthesis in microreactors is still relatively unexplored. Only a few examples are known, in which either an immobilised enzyme or a purified enzyme is used [5–9], catalysing in most cases hydrolytic reactions. Enzymes that are applied in commercial syntheses, however, are typically crude cell lysates or partially purified preparations to reduce the cost price of the biocatalyst [10]. Due to the high surface-to-volume ratio and small channel dimensions, one can expect that clogging can readily occur when this crude cell lysate will be used; this study is the first report in which this issue will be addressed.

In view of the potential advantageous mixing situation in microreactors and the synthetic viability of C–C-bond formation via HNLs, we set out to investigate the possibility of synthesising cyanohydrins in a microchannel using a crude cell lysate in a biphasic system. Therefore, four commercially available aldehydes **2a–d** were selected (Scheme 2) to be converted into their corresponding cyanohydrins in the microreactor setup, using the crude cell lysate of a HNL.


## 2. Experimental

### 2.1. Microreactor (Fig. 1)


The microchannel structure was designed using the software program CleWIN. The actual microreactor was fabricated from borosilicate glass by Micronit Microfluidics BV, Enschede, The Netherlands (HF etched). Chip dimension: length 45 mm, width 15 mm, height 2.2 mm. Channel dimension: width 120  $\mu\text{m}$ , depth 20  $\mu\text{m}$ , total length 70 cm, total internal volume 1.3  $\mu\text{L}$ . A pillar structure was constructed in the middle of the microchannel to stabilise a biphasic laminar flow system, according to a literature procedure [11] (Fig. 1).

### 2.2. Microreactor setup

The microreactor setup that was designed for the experiments is schematically depicted in Scheme 3. In order to perform the experiments, the syringes (brand: SGE Analytical Science; type: 1MDF-LL-GT with Luer Lock) mounted on the syringe pump (brand: Harvard; type: PicoPlus) were connected using Luer adapters, NanoTight nuts and sleeves (brand: Upchurch Sci-



Scheme 2. Aldehydes subjected to the HNLs.

Fig. 1. Schematic representation of the microreactor; total internal volume of the channel is 1.3  $\mu\text{L}$ .

entific; type: P-659, F-331N, F-242) to the capillaries (brand: Polymicro; type: fused silica, i.d. 50–100  $\mu\text{m}$ , o.d. 375  $\mu\text{m}$ ).

These capillaries were connected using the standard procedure for the chipholder (brand: Micronit Microfluidics BV; type: standard chipholder), to the microreactor using ferrules (brand: Upchurch Scientific; type: N-123-03) to ensure a leak free fluidic connection. For optical inspection of the flow inside the microchannel an inverted brightfield reflected light microscope was used (brand: Zeiss; type: Axiovert).


### 2.3. Chemicals

Unless stated otherwise, all chemicals were obtained from commercial sources and used without further purification: MTBE (Acros, 99%), citric acid (Aldrich, 99.5%), 3-phenyl propionaldehyde (Janssen), benzaldehyde (Acros), 2-thiophene-carboxaldehyde (Acros, 98%), 4-methoxy benzaldehyde (Aldrich, 98%), KCN (Acros, p.a.), anisole (Acros, 99%). All liquid aldehydes were distilled before use.

The (*S*)-selective hydroxynitrile lyase, originating from the rubber tree *Hevea brasiliensis* ((*S*)-HbHNL) was applied. The gene encoding for (*S*)-HbHNL was cloned and efficiently expressed in the yeast strain *P. pastoris* as previously described [12]. The enzyme was used as crude cell lysate and was kindly provided by DSM, Geleen, The Netherlands.

### 2.4. General procedure for the enzyme-catalyzed synthesis of cyanohydrins in microchannels

Solution A (organic phase, MTBE) containing 0.23 M aldehyde and 0.18 M anisole (internal standard) and solution B (aqueous phase, 0.4 M citric acid buffer, pH 5.0) containing 0.23 M KCN and 10% (v/v) crude enzyme solution were prepared.



Scheme 3. Schematic representation of the microreactor setup.

Download English Version:

<https://daneshyari.com/en/article/153713>

Download Persian Version:

<https://daneshyari.com/article/153713>

[Daneshyari.com](https://daneshyari.com)