

HOSTED BY



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Journal of Radiation Research and Applied Sciences

journal homepage: <http://www.elsevier.com/locate/jrras>

CrossMark

Insulin, estradiol levels and body mass index in pre- and post-menopausal women with breast cancer

Alkhansa S. Mahmoud ^{a,b,*}, Mohammed A. Abdulrahman ^a,
Khalid H. Bakheit ^{c,d}

^a Sudan Atomic Energy Commission, Khartoum, Sudan

^b Sudan Academy of Sciences – Council of Cancer Researches, Khartoum, Sudan

^c Khartoum University – Faculty of Medicine, Khartoum, Sudan

^d Tabouk University – Faculty of Medicine, Saudi Arabia

ARTICLE INFO

Article history:

Received 7 April 2015

Received in revised form

4 June 2015

Accepted 21 July 2015

Available online 1 August 2015

Keywords:

Insulin

Estradiol

Body mass index

Menopausal

ABSTRACT

Breast cancer is the most common cancer among women where it is associated with considerable morbidity and mortality. The aim of this study was to investigate the association between insulin, estradiol levels and body mass index (BMI) as risk factors for breast cancer. Methods: 80 women newly diagnosed with breast cancer stage I–III invasive breast cancer, were selected randomly and divided in two groups: 40 pre-menopausal aged 26–46 years and 40 post-menopausal aged 52–90 years. Radioimmunoassay used for serum insulin levels measurement, ELISA was used for estradiol levels and BMI calculated by weight (kg)/height (m²). Results: Insulin levels in premenopausal (16.6 ± 10.5) and postmenopausal (17.9 ± 8.8); breast cancer patients showed increasing pattern from the normal levels (4.0–16.0 μIU/ml). While, the levels of estradiol in premenopausal (233 ± 173) and postmenopausal, (549 ± 468); estradiol level in postmenopausal was higher than normal level (50–300 ng/ml), its level showed significantly increase in postmenopausal breast cancer (P.Value = 0.001). Conclusion: Insulin levels increased in pre- and post-menopausal breast cancer patients while estradiol levels do not showed association with premenopausal breast cancer. High BMI, high insulin and estradiol levels in post-menopausal women may be considered as risk factors for breast cancer.

Copyright © 2015, The Egyptian Society of Radiation Sciences and Applications. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Breast cancer is one of the most common types of cancer that affects millions of women around the world (Parkin, Ferlay, &

Pisani, 2005). Insulin level is one of the established risk factors for breast cancer (Muti et al., 2002); it may be associated with common risk factors such as high body mass index (BMI). Insulin is an important metabolic hormone that affects body fat mass (Gisela, 2005). Estradiol is important in sexual

* Corresponding author. Sudan Atomic Energy Commission, Khartoum, Sudan.

E-mail address: alkhansa.salih@gmail.com (A.S. Mahmoud).

Peer review under responsibility of The Egyptian Society of Radiation Sciences and Applications.

<http://dx.doi.org/10.1016/j.jrras.2015.07.004>

1687-8507/Copyright © 2015, The Egyptian Society of Radiation Sciences and Applications. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

development and other body functions. It is produced mainly in the ovaries in premenopausal women, and mainly in fat tissue in postmenopausal women. Estradiol in the blood is linked to an increased risk of breast cancer in women after menopause (Key et al., 2011). Hormone balance that may promote breast cancer growth include elevated levels of circulating or tissue estrogens and insulin which have been a key role in the development and progression of breast cancer (Pollak, 2008); some studies in Western countries have found associations between higher BMI and a higher risk of death among women with breast cancer (Masaaki et al., 2012), although other studies have found no such association (Barnett et al., 2008).

2. Materials and methods

A cross sectional study of 80 Sudanese women newly diagnosed with breast cancer, they were not used any medication including hormone replacement therapy, they were randomly selected to participate in the study. Serum insulin level was measured by using radioimmunoassay (RIA) with polyclonal antibody and antigen I^{125} labeled (Beijin Isotope Nuclear Electronic Machine Co. LTD, China). The normal range of insulin level is from 4.0 to 16.0 μ IU/ml while estradiol levels were measured by ELISA kits with normal range (50–300 ng/ml) and BMI was calculated using the following equation [weight (kg)/height (m^2)]. The patients were classified as underweight ≤ 18.5 , normal weight in ranged 18.5–24, over weight in ranged 25–29.9 and obese ≥ 30 (kg/m^2).

2.1. Statistical analysis

Data analyzed by using the computer program SPSS version 20.0 for frequencies, percentage, mean and standard deviation. Linear correlations were performed by using the Pearson's Correlation Coefficient, r . $p \leq 0.05$ was considered significant.

3. Results

The mean levels and standard deviation of insulin in both pre- and post-menopausal breast cancer patients showed increasing pattern from the normal levels premenopausal (16.6 ± 10.5) and postmenopausal (17.9 ± 8.8), and there was no significant difference (P .Value = 0.5), which may indicate that there is a relationship between high insulin and breast cancer. While, the levels of estradiol in premenopausal (233 ± 173) and postmenopausal (549 ± 468), estradiol in postmenopausal was higher than normal levels (50–300 ng/ml), it's significantly increased in postmenopausal breast cancer (P .Value = 0.001) as shown in Fig. 1. Regarding insulin serum levels there was no significant correlation with BMI in pre- and post-menopausal breast cancer patients ($r = 0.0$, $P = 0.9$ and $r = 0.1$, $P = 0.4$ respectively), also estradiol serum levels and BMI had no significant correlation ($r = 0.01$, $P = 0.9$ and $r = 0.05$, $P = 0.7$ respectively). However, insulin level in premenopausal women increased in normal weight, over weight and obese; underweight women had normal insulin




Fig. 1 – Insulin and estradiol levels among study population.

levels, while estradiol levels based in the normal levels as shown in Table 1. Among postmenopausal, the mean levels of insulin were increased in normal weight and obese; while estradiol levels were increased in different BMI groups as shown in Table 2.

4. Discussion

This study demonstrated that BMI associated with insulin and estradiol levels at diagnosis. Our findings showed high insulin level in both pre- and post-menopausal women with breast cancer. In a relevant study, positive association was found between insulin level and breast cancer in both pre- and post-menopausal women (Nichols et al., 2009). Insulin has been shown to stimulate cell proliferation in normal breast tissue and in human breast cancer cells (Lawlor, Smith, & Ebrahim, 2004). In contrast another study showed that insulin level was strongly correlated with BMI in non-diabetic women newly diagnosed breast cancer (Chappell et al., 2001); the

Table 1 – The mean levels of Insulin and Estradiol among BMI groups in premenopausal women with breast cancer.

BMI (kg/m^2)	Number (%)	Mean \pm SD	
		Estradiol (50–300 ng/ml)	Insulin (4–16.0 μ IU/ml)
Normal weight	16 (40%)	265.1 \pm 243.1	17.4 \pm 12.1
Over weight	14 (35%)	195.5 \pm 151.1	18.5 \pm 9.3
Low weight	8 (20%)	244.9 \pm 142.1	10.2 \pm 1.9
Obese	2 (5%)	192 \pm 45	23.1 \pm 21

Download English Version:

<https://daneshyari.com/en/article/1570275>

Download Persian Version:

<https://daneshyari.com/article/1570275>

[Daneshyari.com](https://daneshyari.com)