



Full paper/Mémoire

Synthesis of 2,4,5-triaryl and 1,2,4,5-tetraaryl imidazoles using silica chloride as an efficient and recyclable catalyst under solvent-free conditions

Hemant V. Chavan^{a,*}, Dattatraya K. Narale^b^a Department of Chemistry, ASP College, Devrukh, Dist: Ratnagiri 415 804, Maharashtra, India^b Department of Chemistry, Gopal Krishna Gokhale College, Kolhapur 416001, Maharashtra, India

ARTICLE INFO

Article history:

Received 11 August 2013

Accepted after revision 5 December 2013

Available online 22 July 2014

Keywords:

Imidazole

Silica chloride

Solvent-free method

Reusability

ABSTRACT

An efficient solvent-free synthesis of various 2,4,5-triaryl imidazoles and 1,2,4,5-tetraaryl imidazoles has been developed using silica chloride as a heterogeneous catalyst. The present methodology offers several advantages, such as excellent yields, shorter reaction times, economic availability, and reusability of catalyst.

© 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.


1. Introduction

In recent years, considerable interest has been devoted to finding new methodologies for the synthesis of numerous organic compounds under solvent-free conditions. The toxicity and volatile nature of many organic solvents, particularly chlorinated hydrocarbons that are widely used in large amounts for organic reactions have posed a serious threat to the environment [1]. Thus, the design of solvent-free catalytic reactions has received tremendous attention in recent times in the area of green synthesis [2]. Imidazoles are common scaffolds in highly significant biomolecules, including biotin, the essential amino acid histidine, histamine, the pilocarpine alkaloids [3], and other alkaloids, which have been shown to exhibit interesting biological activities, such as antimicrobial, anti-ryptococcal, cytotoxic activities and inhibition of nitric oxide synthase [4]. Many of the substituted imidazoles are known as inhibitors of P38 MAP kinase [5], fungicides and herbicides

[6], plant growth regulators [7] and therapeutic agents [8]. Owing to the wide range of pharmacological and biological activities, the development of synthetic methods enabling facile access to this heterocycle is still desirable. Several methods have been reported for the synthesis of imidazoles. 2,4,5-Triaryl-1*H*-imidazoles are generally synthesized by three-component cyclocondensation of a 1,2-diketone, α -hydroxyketone or α -ketomoxime with an aldehyde and ammonium acetate, which comprises the use of microwaves [9], ionic liquids [10] europium triflate [11], oxalic acid [12], TBAB [13], CAN [14], refluxing in AcOH [15] and silica sulfuric acid [16]. On the other hand, the synthesis of 1,2,4,5-tetraaryl-1*H*-imidazoles are carried out by four-component condensation of a 1,2-diketone, α -hydroxyketone or α -ketomoxime with an aldehyde, a primary amine and ammonium acetate using microwaves [17], heteropolyacid [18], silica gel/NaHSO₄ [19] InCl₃·3H₂O [20] or HClO₄-SiO₂ [21]. In addition, they can also be accessed by hetero-Cope rearrangements [22] and condensation of a 1,2-diketone with an aryl nitrile and primary amine under microwave irradiation [23]. Despite their potential utility, these methods have a number of drawbacks, including poor yields, utilization of excessive glacial AcOH, prolonged

* Corresponding author.

E-mail address: hemantchavan.sus@rediffmail.com (H.V. Chavan).



Scheme 1. Silica chloride catalyzed synthesis of 2,4,5-triaryl-1H-imidazoles.

reaction times, high cost of most conventional room temperature ionic liquids, use of hazardous and expensive catalysts and use of toxic organic solvent that limit their utility.

Hence, the development of clean, safe, effective, economical, high-yielding and environmentally benign protocols is still desirable and much in demand.

Silica chloride ($\text{SiO}_2\text{-Cl}$) is gaining importance as a heterogeneous catalyst and is used in various organic transformations. It is a good solid acid in terms of convenience, cheapness, easy production, and insolubility in all organic solvents. Due to its insolubility in organic solvents, excellent *in situ* proton generation, and accepting different nucleophiles, it can be used for different purposes in organic chemistry. It has been used as a dehydrating agent [24], and some cyclocondensations yielding dihydropyrimidinone [25], 2-aminothiazoles [26], 6-methyl-3-propynylthio-1,2,4-triazin-5(2H)-one [27], pyranoquinolines, or furanoquinolines [28] are catalyzed by silica chloride. The use of silica chloride to expedite oxidation of alcohol [29], esterification and transesterification [30] transdithioacetalization of acetals [31], deprotection of oximes/hydrazones/semicarbazones [32] synthesis of nitriles and amides [33], synthesis of sulfoxide [34], synthesis of bisindolylmethanes [35], etc. has been well explored.

Considering the significance of the silica chloride and in continuation of our interest in developing new and rapid synthetic methods for the construction of biologically important structural scaffolds [36], we report herein a rapid, efficient, economic, environmentally benign and easy to scale-up method for the synthesis of 2,4,5-triaryl-1H-imidazoles using silica chloride as a heterogeneous and recyclable catalyst under solvent-free conditions (Scheme 1).

2. Results and discussion

Initially, to optimize the amount of catalyst and the reaction temperature, the reaction of benzil (**1**, 1 mmol), benzaldehyde (**2a**, 1 mmol), and NH_4OAc (10 mmol) was studied under solvent-free conditions in the presence of $\text{SiO}_2\text{-Cl}$ at different temperatures; the results are summarized in Table 1. They revealed that, when 250 mg/mmol $\text{SiO}_2\text{-Cl}$ at 80 °C were used, the reaction proceeded smoothly and gave product **3a** in highest yield (80%). Further increase in the amount of silica chloride above 250 mg/mmol did not lead to substantial change in the yield and the reaction time (Table 1). Then, we examined

the effect of solvents, like CH_3CN , toluene, EtOH, MeOH and THF for the above reaction. The results in Table 2 indicate that solvents affected the efficiency of the reaction and very poor yields were obtained after 1 h. The best results were obtained under solvent-free conditions (Table 2).

Under optimized reaction conditions, a mixture of benzil (1 mmol), an aromatic aldehyde (1 mmol), NH_4OAc (10 mmol) and $\text{SiO}_2\text{-Cl}$ (250 mg) was stirred at 80 °C for the appropriate time (Table 3). After completion of the reaction (TLC), CH_2Cl_2 (20 mL) was added, and the solid catalyst was recovered by filtration. The solvent was distilled off and the crude product was purified by column chromatography using (SiO_2 , hexane:AcOEt) as an eluent to afford pure 2,4,5-trisubstituted imidazoles in good to excellent yields. Moreover, the catalyst recovered from the reaction mixture was reused for successive reactions

Table 1
Synthesis of 2,4,5-triaryl-1H-imidazole (**3a**) under various conditions.

Entry	$\text{SiO}_2\text{-Cl}$ (mg/mmol) ^a	Temperature (°C)	Time (min)	Yield (%) ^b
1	–	80	180	00
2	50	80	120	45
3	100	80	120	58
4	150	80	90	65
5	200	80	60	72
6	200	100	60	75
7	250	60	60	55
8	250	80	30	80
9	250	100	30	80
10	300	80	30	79
11	300	100	30	80
12	300	120	30	78

^a Reaction conditions: benzil (1 mmol); benzaldehyde (1 mmol); ammonium acetate (10 mmol); neat.

^b Yields of isolated products.

Table 2
Effect of solvent for $\text{SiO}_2\text{-Cl}$ catalyzed synthesis of 2,4,5-triaryl imidazole (**3a**).

Entry	Solvent ^a	Temperature (°C)	Time (min)	Yield (%) ^b
1	CH_3CN	Reflux	60	20
2	Toluene	Reflux	60	40
3	EtOH	Reflux	60	30
4	MeOH	Reflux	60	35
5	THF	Reflux	60	25
6	none	80	30	80

^a Reaction conditions: benzil (1 mmol); benzaldehyde (1 mmol); ammonium acetate (10 mmol), solvent (5 mL).

^b Yields of isolated products.

Download English Version:

<https://daneshyari.com/en/article/170422>

Download Persian Version:

<https://daneshyari.com/article/170422>

[Daneshyari.com](https://daneshyari.com)