



# Ten limit cycles around a center-type singular point in a 3-d quadratic system with quadratic perturbation<sup>☆</sup>



Pei Yu<sup>a,b,\*</sup>, Maoan Han<sup>a</sup>

<sup>a</sup> Department of Mathematics, Shanghai Normal University, Shanghai, 200234, PR China

<sup>b</sup> Department of Applied Mathematics, Western University, London, Ontario N6A 5B7, Canada

## ARTICLE INFO

### Article history:

Received 14 October 2014

Received in revised form 9 December 2014

Accepted 9 December 2014

Available online 27 December 2014

### Keywords:

3-d quadratic system

Focus

Center

Limit cycle

Hopf bifurcation

Center manifold

Normal form

Focus value computation

## ABSTRACT

In this paper, we show that perturbing a simple 3-d quadratic system with a center-type singular point can yield at least 10 small-amplitude limit cycles around a singular point. This result improves the 7 limit cycles obtained recently in a simple 3-d quadratic system around a Hopf singular point. Compared with Bautin's result for quadratic planar vector fields, which can only have 3 small-amplitude limit cycles around an elementary center or focus, this result of 10 limit cycles is surprisingly high. The theory and methodology developed in this paper can be used to consider bifurcation of limit cycles in higher-dimensional systems.

© 2015 Elsevier Ltd. All rights reserved.

## 1. Introduction

The phenomenon of limit cycles can arise in almost all areas of science and engineering, from physics, biology, ecology, economy, mechanics, electronics, and even from financial and social systems. Limit cycle theory has been playing a very important role in the study of dynamical behavior of nonlinear systems. The study of limit cycles was initiated by Poincaré [1], and its further development was perhaps motivated by the well-known Hilbert's 16th problem [2], which was posed by Hilbert in 1900. The second part of this problem is to find the upper bound, called the Hilbert number  $H(n)$ , on the number of limit cycles that planar polynomial systems of degree  $n$  can have. Many results have been obtained in study of the Hilbert number; see for example [3–7]. However, the finiteness problem remains unsolved even for quadratic polynomial systems, that is, whether  $H(2) = 4$  is still open. For cubic polynomial systems, many results have been obtained on the lower bound of the Hilbert number. So far, the best result for cubic systems is  $H(3) \geq 13$  [8,9].

When we consider the limit cycles bifurcating from isolated fixed points, Hilbert's 16th problem becomes studying degenerate Hopf bifurcations. This local problem has been completely solved only for generic quadratic systems [10], showing existence of 3 limit cycles in the vicinity of a singular point which may be a focus or center. For cubic systems, regarding the case of focus, the best result obtained so far is 9 limit cycles; see for example [11] and references therein. In the case

<sup>☆</sup> This work was supported by the Natural Science and Engineering Research Council of Canada (No. R2686A02) and National Natural Science Foundation of China (No. 1271261 and No. 11431008).

\* Corresponding author.

E-mail addresses: [pyu@uwo.ca](mailto:pyu@uwo.ca) (P. Yu), [mahan@shnu.edu.cn](mailto:mahan@shnu.edu.cn) (M. Han).

of center, on the other hand, it has been shown that more limit cycles can bifurcate from a singular point. The best result obtained recently is 12 limit cycles; see more details in [12] where a literature review is given.

Higher-dimensional vector fields may not only more likely exhibit limit cycles, but also may co-exist complex dynamical behaviors such as chaos. But not many results have been obtained on limit cycles for higher-dimensional vector fields since the analysis for such systems is much more complex than planar systems. A very natural extension from the study of limit cycles in 2-d systems is to study 3-d systems with a Hopf singular point. A similar problem is to investigate the maximal number of limit cycles which may exist in the vicinity of such a singular point under proper perturbations. This is a very challenging problem. There are very few results in the literature. However, unlike 2-d systems, an interesting finding has revealed that even a quadratic 3-d system can exhibit infinitely many small limit cycles, which appear on an infinite family of algebraic invariant surfaces [13,14]. In this paper, we want to ask: What is an upper bound for the cyclicity of a singular point of a quadratic 3-d system restricted to a single center manifold? Although we cannot answer this open problem, we will try to provide a better lower bound in this paper and hope that this will help promote research in this direction.

Restricted to a single center manifold, a well-known 3-d competitive Lotka–Volterra model has been studied extensively over the last twenty years. The model is described by a 3-d differential system:

$$\dot{x}_i = x_i \left( b_i - \sum_{j=1}^3 a_{ij} x_j \right), \quad i = 1, 2, 3, \quad (1)$$

where the dot indicates differentiation with respect to time,  $t$ ,  $x_i$  represents the population of the  $i$ th species, and the coefficients take positive real values,  $b_i > 0$ ,  $a_{ij} > 0$ ,  $i, j = 1, 2, 3$ . This is a special case of general 3-d quadratic systems. This system has 8 equilibria and only one of them is a positive equilibrium (i.e., all components of the equilibrium take positive values). An interesting problem is to consider the limit cycles around the positive equilibrium. In the past two decades, several researchers have paid attention to this model and particularly studied bifurcation of limit cycles around the positive equilibrium (e.g., see [15–19]). So far, the best result is 4 limit cycles, obtained by Gyllenberg & Yan [19], using appropriate parameter values. These 4 limit cycles include 3 small-amplitude limit cycles, proved by using focus value computation, and one large limit cycle, shown by constructing a heteroclinic loop. Recently, Tian and Yu revisited this problem [20] and showed that this system, which may not be a competitive system, might have maximal 8 limit cycles, but it is very difficult to prove this using the existing methodology.

For general 3-d quadratic systems with a Hopf singular point, the best result obtained so far is 7 small-amplitude limit cycles bifurcating from the origin of a simple 3-d system [21]. In this paper, we shall consider a quadratic system similar to the one in [21] but with a center-type critical point at the origin. Then, by adding quadratic polynomial perturbations in the form of  $\varepsilon p_i(x_1, x_2, x_3)$  to each equation of the system, we obtain 10 small-amplitude limit cycles around the origin. In the next section, we formulate the 3-d quadratic system, and then in Section 3 we prove existence of 10 limit cycles.

## 2. Formulation of 3-d system

In this section, we present a 3-d quadratic system which may yield 10 limit cycles. To achieve this, we start from the following generic 3-d quadratic polynomial system:

$$\begin{aligned} \dot{x}_1 &= \alpha x_1 + x_2 + a_{11} x_1^2 + (2b_{11} + a_{12})x_1x_2 + a_{22} x_2^2 + a_{33} x_3^2 + a_{13} x_1x_3 + a_{23} x_2x_3, \\ \dot{x}_2 &= -x_1 + \alpha x_2 + b_{11} x_1^2 + (2a_{11} + b_{12})x_1x_2 + b_{22} x_2^2 + b_{33} x_3^2 + b_{13} x_1x_3 + b_{23} x_2x_3, \\ \dot{x}_3 &= -\beta x_3 + c_{11} x_1^2 + c_{12} x_1x_2 + c_{22} x_2^2 + c_{33} x_3^2 + c_{13} x_1x_3 + c_{23} x_2x_3, \end{aligned} \quad (2)$$

where  $\alpha, \beta > 0$ ,  $a_{ij}, b_{ij}$  and  $c_{ij}$  are parameters. Note that the first two equations have taken Bautin's system format, which can be obtained by a rotating transformation. System (2) has a Hopf singularity at the origin when  $\alpha = 0$ . To simplify the generic system (2), first note that without loss of generality, suppose  $b_{11} \neq 0$ ,  $c_{12} \neq 0$ , and then we can introduce proper scaling into system (2) so that  $b_{11} = c_{12} = 1$ . Then, we want that the focus values of the resulting system around the origin are all zero, i.e. the origin becomes a center-type point. There are many possibilities, but a complete classification is difficult. Here, we set  $\alpha = a_{33} = a_{13} = a_{23} = b_{33} = b_{13} = b_{23} = c_{11} = c_{22} = c_{13} = c_{23} = 0$ , and  $\beta = c_{33} = 1$ . According to Bautin's theory, setting  $a_{22} = -a_{11}$  in the resulting system yields all focus values to vanish, i.e. the origin becomes a center restricted to a center manifold under this condition. Finally, we add quadratic perturbations to system (2) under above conditions to obtain the following perturbed system:

$$\begin{aligned} \dot{x}_1 &= x_2 + a_{11} x_1^2 + (2 + a_{12})x_1x_2 - a_{11} x_2^2 + \varepsilon(a_{100} x_1 + a_{020} x_2 \\ &\quad + a_{110} x_1^2 + a_{120} x_1x_2 + a_{220} x_2^2 + a_{330} x_3^2 + a_{130} x_1x_3 + a_{230} x_2x_3), \\ \dot{x}_2 &= -x_1 + x_1^2 + 2a_{11} x_1x_2 - x_2^2 + \varepsilon(b_{100} x_1 + b_{020} x_2 + b_{110} x_1^2 \\ &\quad + b_{120} x_1x_2 + b_{220} x_2^2 + b_{330} x_3^2 + b_{130} x_1x_3 + b_{230} x_2x_3), \\ \dot{x}_3 &= -x_3 + x_1x_2 + x_3^2 + \varepsilon(c_{100} x_1 + c_{020} x_2 + c_{110} x_1^2 + c_{120} x_1x_2 \\ &\quad + c_{220} x_2^2 + c_{330} x_3^2 + c_{130} x_1x_3 + c_{230} x_2x_3), \end{aligned} \quad (3)$$

where  $0 < \varepsilon \ll 1$  is a perturbation parameter.

Download English Version:

<https://daneshyari.com/en/article/1707766>

Download Persian Version:

<https://daneshyari.com/article/1707766>

[Daneshyari.com](https://daneshyari.com)