



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data article

Data on the calcium-induced mobility shift of myristoylated and non-myristoylated forms of neurocalcin delta



Jeffrey Viviano^a, Anuradha Krishnan^a, Hao Wu^a,
Venkat Venkataraman^{a,b,*}

^a Graduate School of Biomedical Sciences, Rowan University, Stratford, NJ 08084, USA

^b School of Osteopathic Medicine, Rowan University, Stratford, NJ 08084, USA

ARTICLE INFO

Article history:

Received 19 January 2016

Received in revised form

21 February 2016

Accepted 3 March 2016

Available online 11 March 2016

ABSTRACT

This data article presents the differences observed between the myristoylated and non-myristoylated forms of the neuronal calcium sensor protein, neurocalcin delta (NCALD). Analysis of the myristoylated and non-myristoylated versions of the protein by mass spectrometry provided difference in mass values consistent with addition of myristoyl group. In the presence of calcium, mobility retardation was observed upon electrophoresis of the protein in native gels. The retardation was dose-dependent and was exhibited by both the myristoylated and non-myristoylated forms of the protein.

© 2016 Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Biology
More specific subject area	Electrophoretic Techniques
Type of data	Table, graph, figure

DOI of original article: <http://dx.doi.org/10.1016/j.ab.2015.11.005>

* Corresponding author.

E-mail address: vvenkat2007@gmail.com (V. Venkataraman).

<http://dx.doi.org/10.1016/j.dib.2016.03.021>

2352-3409/© 2016 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

How data was acquired	Mass spectroscopy: Bruker Microflex LRF MALDI-TOF Electrophoresis: Bio-Rad miniPROTEAN
Data format	Analyzed
Experimental factors	For mass spectrometry, the samples were co-crystallized in 1:1 mixture of sinapinic acid and matrix solution containing acetonitrile, water and trifluoroacetic acid. For electrophoresis, standard protocols were used.
Experimental features	Myristoylated and non-myristoylated forms of NCALD were analyzed
Data source location	Stratford, NJ 08012, USA
Data accessibility	Data is within this article

Value of the data

- The relevance of myristoylation to the biological functions of Neuronal Calcium Sensor (NCS) proteins remains to be elucidated.
- Data is presented on the myristoylated and non-myristoylated forms of NCALD assessed by calcium-induced mobility shift assay.
- The calcium-induced mobility shift assay may be useful in assessing the contribution of myristoylation and other post-translational modifications to response of all NCS proteins to calcium.

1. Data

Bacterially expressed NCALD was purified in its myristoylated and non-myristoylated forms independently. Mass spectrometric analyses was carried out to determine the mass of each form and to determine if the myristoylated form also contained the non-myristoylated form. A table of the mass values of the two forms is provided, along with those documented in an earlier report by other investigators. Based on the analyses, there is little non-myristoylated NCALD in the myristoylated preparations. The two preparations were subjected to electrophoresis in native gels in the presence of incremental concentrations of calcium. A dose-dependent mobility retardation is observed with both forms. However, the myristoylated form exhibits a greater amplitude and increased sensitivity to change in calcium concentrations.

2. Experimental design, materials and methods

NCALD was expressed in *E. coli* ER2566 as described in [1]. Briefly, cells grown overnight were inoculated (1% inoculum) into fresh LB medium and grown to an optical density of 0.6 at 600 nm. IPTG (1 mM final concentration) was then added for induction. For myristoylation, cells with yeast *N*-Myristoyl Transferase were used and myristic acid was supplemented. Cells without the transferase were used to generate the non-myristoylated form; the supplementation was also skipped. Cells were collected 2.5 h after induction, sonicated and the protein was purified on phenyl sepharose columns as described previously [2,3]. The purified protein was then washed with calcium-depleted Tris–Cl

Table 1
Mass Spectrometric Analyses of NCALD.

Description	Myr ⁻ Neurocalcin	Myr ⁺ Neurocalcin
Molar mass (g mol ⁻¹ ± SD) (MALDI-MS)	22,107.1 ± 0.8	22,325.8 ± 1.4
Previously reported molar mass (g mol ⁻¹ ± SD) (ESI-MS)	22,110 ± 2	22,325 ± 2

Download English Version:

<https://daneshyari.com/en/article/174803>

Download Persian Version:

<https://daneshyari.com/article/174803>

[Daneshyari.com](https://daneshyari.com)