



The STARK-B database as a resource for “STARK” widths and shifts data: State of advancement and program of development

Sylvie Sahal-Bréchet^{a,*}, Milan S. Dimitrijević^{a,b}, Nicolas Moreau^a, Nabil Ben Nessib^c

^a *Laboratoire d'Étude du Rayonnement et de la Matière en Astrophysique, Observatoire de Paris, UMR CNRS 8112, UPMC, 5 Place Jules Janssen, 92195 Meudon Cedex, France*

^b *Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia*

^c *Department of Physics and Astronomy, College of Science, King Saud University, Riyadh 11451, Saudi Arabia*

Available online 28 August 2013

Abstract

“Stark” broadening theories and calculations have been extensively developed for about 50 years and can now be applied to many needs, especially for accurate spectroscopic diagnostics and modeling. This requires the knowledge of numerous collisional line profiles. Nowadays, the access to such data via an online database becomes essential. STARK-B is a collaborative project between the Astronomical Observatory of Belgrade and the Laboratoire d'Étude du Rayonnement et de la matière en Astrophysique (LERMA). It is a database of calculated widths and shifts of isolated lines of atoms and ions due to electron and ion collisions (impacts). It is devoted to modeling and spectroscopic diagnostics of stellar atmospheres and envelopes, laboratory plasmas, laser equipments and technological plasmas. Hence, the domain of temperatures and densities covered by the tables is wide and depends on the ionization degree of the considered ion. STARK-B has been fully opened since September 2008 and is in free access.

The first stage of development was ended in autumn 2012, since all the existing data calculated with the impact semiclassical-perturbation method and code by Sahal-Bréchet, Dimitrijević and coworkers have now been implemented. We are now beginning the second stage of the development of STARK-B. The state of advancement of the database and our program of development are presented here, together with its context within VAMDC. VAMDC (Virtual Atomic and Molecular Data Center) is an international consortium which has built a secure, documented, flexible interoperable platform e-science permitting an automated exchange of atomic and molecular data.

© 2013 COSPAR. Published by Elsevier Ltd. All rights reserved.

Keywords: Databases; Atomic data; Atomic processes; Line: profiles; Stars: atmospheres

1. Introduction

Broadening of spectral lines due to perturbation of emitting or absorbing atoms, ions, or molecules, by their interactions with the surrounding particles in a plasma or gas is called pressure broadening. If perturbers are charged particles, electrons or ions, such broadening mechanism is called Stark broadening. Data on Stark broadening are of interest for plasma diagnostic and modeling, as well as for spectra

analysis and synthesis in many domains for research of different kinds of plasma in laboratory, inertial fusion investigations, for lasers and laser produced plasmas and especially in astrophysics, since thanks to space missions, large space observatories like Hubble, Chandra, Spitzer, Lyman, and to ground-based telescopes of the ten meters class, spectra with very high resolution could be obtained from X to radio wavelength ranges.

In stellar astronomy, Stark broadening data are of particular importance for white dwarfs, where Stark broadening is the dominant collisional line broadening mechanism (Popović et al., 1999b; Tankosić et al., 2003; Simić et al., 2006; Hamdi et al., 2008; Simić et al., 2009; Dimitrijević

* Corresponding author. Tel.: +33 15077442; fax: +33 1 45077100.

E-mail addresses: sylvie.sahal-brechot@obspm.fr (S. Sahal-Bréchet), mdimitrijevic@aob.bg.ac.rs (M.S. Dimitrijević), nicolas.moreau@obspm.fr (N. Moreau), nbennessib@ksu.edu.sa (N. Ben Nessib).

et al., 2011; Dufour et al., 2011; Larbi-Terzi and Sahal-Bréchet, 2012). Such data are also of significance for interpretation, analysis and synthesis of A and B type star spectra (see e.g. Lanz et al., 1988; Popović et al., 2001a,b, 1999a,b; Tankosić et al., 2003; Simić et al., 2005; Sahal-Bréchet et al., 2010). Modern codes for stellar atmosphere modeling, like e.g. PHOENIX (Baron et al., 1998; Hauschildt et al., 1999; Short et al., 1999) require the knowledge of atomic data for as much as possible transitions, especially for trace elements, so that the access to such atomic data, including Stark broadening ones, via online databases becomes very important.

The database STARK-B (Sahal-Bréchet et al., 2013), which contains Stark broadening parameters, namely calculated line widths and shifts of isolated lines of atoms and ions due to collisions with electrons, protons and different ions, first of all ionized helium as the most important after electrons and protons for stellar atmospheres, is not useful only for modeling and spectroscopic diagnostics of stellar atmospheres and envelopes, but also for laboratory plasmas, inertial fusion plasma, laser equipments and technological plasmas. This is a common project where participants are from the Paris Observatory and the Astronomical Observatory of Belgrade. The database STARK-B has been opened online since September 2008, and is in free access. The objective of the first phase of database development, was to implement all the existing data calculated with the impact semiclassical-perturbation method and code by Sahal-Bréchet, Dimitrijević and coworkers. We are now working on the second stage and the state of advancement of the database STARK-B will be presented here, as well as our program of development. We note also that STARK-B enters in Virtual Atomic and Molecular Data Center – VAMDC (Dubernet et al., 2010; Rixon et al., 2010), a secure, documented, flexible, interoperable platform based on e-science, which permits an optimized search and exchange of atomic and molecular data. The relation between STARK-B database and VAMDC was discussed in Sahal-Bréchet et al. (2012).

2. STARK-B database

Within the impact-complete collision-isolated line approximation, the profile of an isolated spectral line is the Lorentz one, characterized by a width at half maximum of intensity and a shift, so called Stark broadening parameters, depending on temperature T and density N of perturbers. For not too high densities, due to the impact approximation, widths and shifts are linearly proportional to the density of perturbers. But at high densities, due to the Debye screening effect which decreases the width and the shift, and which for such conditions can become significant, the dependence of Stark broadening parameters as a function of perturber density may decline from the linear one. This is taken into account in majority of our calculations, so that for many of atoms and ions the Stark broad-

ening parameters for a grid of perturber densities are implemented in STARK-B.

In the first stage of STARK-B development, we implemented in STARK-B Stark broadening parameters determined with the semiclassical-perturbation method (SCP) developed by Sahal-Bréchet (1969a,b) and updated in further papers by Sahal-Bréchet (1974) for more complex atoms, by Fleurier et al. (1977) for inclusion of Feshbach resonances in elastic cross-sections for the line widths for ion emitters, and by Mahmoudi et al. (2008) for the particular cases of very complex atoms. The numerical codes have been also updated by Dimitrijević and Sahal-Bréchet (1984) and in further papers. The accuracy is about 20% for the widths of simpler spectra but is worse for the shifts and very complex spectra, particularly when configuration mixing is present in description of energy levels. The results of Stark broadening parameters determination performed by Dimitrijević, Sahal-Bréchet, and co-workers using the semiclassical-perturbation method are contained in more than 130 publications and now have been implemented in the STARK-B database.

STARK-B is devoted for modeling and spectroscopic diagnostics of various plasmas in astrophysics, laboratory physics, technology and other topics, so that the range of temperatures and densities in the tables is wide and vary with the ionization degree of the considered ion. The temperatures in tables vary from several thousands Kelvins for neutral atoms to several millions for highly charged ions. Also the perturber densities vary from 10^{12} cm^{-3} to several 10^{22} cm^{-3} . The presentation of data, in particular the definition of configurations, terms and atomic energy levels is in accordance with the VAMDC standards, in order to allow interoperability with other atomic databases included in the Virtual Atomic and Molecular Data Center. We note also that the wavelengths in STARK-B are usually calculated from the energy levels that are used as input data for the determination of Stark broadening parameters. Consequently, the tabulated wavelengths are most often different from the measured ones, especially if the used energy levels are theoretically calculated. For the identification of the lines, the configurations, terms and levels are included in tables as well as the multiplet number from NIST database (Kramida et al., 2012).

If one opens the homepage, proposed menus are “Introduction”, “Data Description”, “Access to the Data”, “Updates” and “Contact”. In “Introduction” methods of Stark broadening parameter calculations and different approximations which are adopted are briefly explained. “Data Description” describes the data which are tabulated. “Access to the Data” provides a graphical interface enabling to the visitor to click on the needed element in the Mendeleev periodic table and then on the corresponding ionization degree. For elements in yellow cells, which symbols are additionally enhanced by boldface, Stark broadening data exist in the database, and for other cells, which have the color a little bit lighter than the background, there is no data. After choosing the needed

Download English Version:

<https://daneshyari.com/en/article/1764683>

Download Persian Version:

<https://daneshyari.com/article/1764683>

[Daneshyari.com](https://daneshyari.com)