

INTEGRAL observation of the Crab pulsar

T. Mineo^{a,*}, G. Cusumano^a, L. Foschini^b, A. Segreto^a, A. Bazzano^c, G. Di Cocco^b,
G. Malaguti^b, L. Natalucci^c

^a *INAF-IASF, Sezione di Palermo, via U. La Malfa 153, 90146 Palermo, Italy*

^b *INAF-IASF, Sezione di Bologna, via P. Gobetti 101, 40129 Bologna, Italy*

^c *INAF-IASF, via Fosso del Cavaliere 100, 00133 Roma, Italy*

Received 29 September 2004; received in revised form 6 December 2004; accepted 8 December 2004

Abstract

In this paper, we present the timing and spectral analysis of several observations of the Crab pulsar with INTEGRAL. All these observations, when summed together provide an high-statistics data set which can be used for accurate phase resolved spectroscopy of the pulsed emission over the energy range 3–350 keV. In particular, we present the light curves in several energy ranges and phase-resolved spectral analysis of the pulsed emission.

© 2005 COSPAR. Published by Elsevier Ltd. All rights reserved.

Keyword: Pulsar

1. Introduction

The Crab pulsar (PSR B0531 + 21) can be observed in almost every energy band, of the electromagnetic spectrum. Its pulse profile is characterized by a double peak structure with a phase separation of 0.4 that is almost aligned in absolute phase over all wavelengths (Rots et al., 2000; Tennant et al., 2001; Kuiper et al., 2003).

In the X-ray range, the spectral energy distribution of the Crab pulsar profile changes with phase: the first peak (P1), dominant at low X-ray energies, becomes smaller than the second one (P2) at soft γ -rays; moreover, in the same range, an enhancement with energy of the bridge between these peaks, usually called interpeak (Ip), is also observed.

A first detailed study of the phase-resolved spectra has been performed by Pravdo et al. (1997), in the 5–

200 keV energy interval, based on RXTE (PCA and HEXTE) data. They found a variation of the photon index as function of the pulse phase, with the inter-peak region systematically harder than the main peaks. Similar results have been reported by Massaro et al. (2000) in the energy range observed by BeppoSAX (0.1–300 keV). In particular, a photon index difference of 0.14 ± 0.03 and 0.31 ± 0.07 is observed between the first peak and the second peak, and between the first peak and the interpeak, respectively. Moreover, these authors found that the photon index relative to the same phase interval significantly increases with energy, and that the phase-resolved spectral distribution can be well-modeled by a single curved power law with a slope variable with $\log(E)$. Applying this model to three wider phase intervals, the first peak, the inter-peak and the second peak, a value of ~ 0.15 for the bending parameter has been measured in the three intervals (Massaro et al., 2001).

Kuiper et al. (2001) presented an exhaustive high-energy picture of the Crab pulsar from 0.1 keV up to 10 GeV by using the high-energy γ -ray data from the

* Corresponding author. Tel.: +39 91 6809478; fax: +39 91 6882258.
E-mail address: teresa.mineo@pa.iasf.cnr.it (T. Mineo).

CGRO satellite together with data obtained at X-ray energies from other observatories.

In this paper, we present the timing and spectral analysis of several observations of the Crab pulsar with JEM-X2, ISGRI and PICSIT on board INTEGRAL.

2. Observation and data reduction

The International Gamma-Ray Astrophysics Laboratory (INTEGRAL; Winkler et al., 2003) observed the Crab nebula and pulsar for calibration purposes several times from February 2003 (Rev. 39) to August 2003 (Rev. 103). In this paper, we present the analysis of the data obtained by JEM-X (Lund et al., 2003) and IBIS (Ubertini et al., 2003). JEM-X consists of two identical coded-aperture mask telescopes with a geometrical area of 500 cm² and microstrip gas chamber (MSGC) at the focal plane that operates in the energy range 3–35 keV. IBIS is a coded aperture telescope composed by two detection layers: ISGRI (Lebrun et al., 2003) and PICSIT (Di Cocco et al., 2003). ISGRI is a large CdTe γ -ray camera operating in the range 15 keV–1 MeV, with a geometrical area of 2621 cm² and an energy resolution of $\sim 8\%$ at 60 keV. PICSIT is composed by 64 \times 64 Caesium–Iodide (CsI) scintillation pixels working in the energy intervals 175 keV–10 MeV. The time resolution is 90 and 150 μ s for IBIS and JEM-X, respectively; the INTEGRAL absolute timing accuracy, as estimated by Kuiper et al. (2003) from Crab data, is about 40 μ s. PICSIT detector cannot be routinely configured in photon-by-photon mode due to the tight telemetry budget. For timing studies, observers can select the spectral–timing mode in which spectra are accumulated on board in four energy bins with an integration time of 1 ms.

Observations with a maximum off-axis angle of 2° were selected for the analysis. For JEM-X, we consid-

ered data from only one of the two units (JEM-X2); for PICSIT, we selected observation intervals with time resolution of 1 ms. Moreover, to increase the statistics of PICSIT light curve, data from Rev. 0041, with an off-axis of 9.6°, were also included. Table 1 summarizes the log of the observations used in this analysis together with the time exposures. Standard reduction procedures have been applied to data and photon list files have been generated with the standard pipeline (INTEGRAL OSA v3.0). In particular, we created for JEM-X2 and ISGRI a list of photons selecting only events falling in pixels illuminated by the Crab, increasing the signal-to-noise ratio. PICSIT data were accumulated in four light curves, one for each available energy channel.

JEM-X2 and ISGRI response matrices are also provided by the standard software; in particular, matrices provided with OSA v4.1 has been used for spectral analysis. No response matrix is available at the moment for PICSIT; data from this detector has not been included in the spectral analysis.

3. Timing and spectral analysis

Arrival times were converted to the Solar System Barycentre and folded with proper Crab ephemerides. Our data set spans several months and for each observation we used contemporary radio ephemeris provided by Jodrell Bank radio telescope (UK) that continuously monitors this pulsar.

The resulting phase histograms in six energy bands from 3 to 360 keV are shown in Fig. 1 with a phase resolution ranging from 0.01 (0.33 ms) to 0.03 (1.1 ms). The well-known double peaked structure is prominent in all the profiles with a high-statistical significance.

A detailed phase resolved spectral analysis was performed by selecting spectra in 43 phase intervals 0.01 wide between phase -0.1 and 0.46 . The off-pulse level,

Table 1
Observation log for the data used in this analysis

Revolution	Start–stop time (MJD)	Exposure (ks)		
		JEM-X2	ISGRI	PICSIT
39	52677.2–52679.8	149.3	142.3	–
40	52680.2–52681.6	122.6	56.0	106.5
41	52683.2–52685.8	–	–	204.0
42	52686.4–52688.2	75.4	61.6	–
43	52689.6–52691.7	–	30.7	–
44	52692.2–52694.4	–	9.8	–
45	52695.2–52696.7	–	2.1	–
102	52866.3–52868.1	–	37.9	–
103	52868.6–52868.8	–	21.6	–
	Total	347.3	362.0	310.5

Download English Version:

<https://daneshyari.com/en/article/1768309>

Download Persian Version:

<https://daneshyari.com/article/1768309>

[Daneshyari.com](https://daneshyari.com)