

Radioactivity levels in soil of salt field area, Kelambakkam, Tamilnadu, India

R. RAVISANKAR^{1*} A. RAJALAKSHMI¹ P. ESWARAN² V GAJENDIRAN.³
V. MEENAKSHISUNDRAM³

¹ Department of Physics, Sri Sivasubramania Nadar College of Engineering, Kalavakkam, Chennai- 603110, Tamilnadu, India;

² Department of Physics, SMK Fomra Institute of Technology, Thaiyur, Chennai- 603 103, Tamilnadu, India;

³ Radiological Safety Section, Health and Safety Division, Indira Gandhi Centre for Atomic Research (IGCAR), Kalpakkam- 603102, Tamilnadu, India)

Abstract Soil samples in and around salt field area have been analyzed for natural radioactivity concentration using gamma ray spectrometer. The activity concentration for ²³²Th, ²³⁸U and ⁴⁰K are ranged from 27.88 to 45.27 Bq/kg, from BDL to 13.30 Bq/kg, and from 135.54 to 381.28 Bq/kg, respectively. The measured activity concentrations for these radionuclides were compared with world average activity of soil. The average activity concentration of ²³²Th in the present study is 1.19 times higher than world median value while the activity of ²³⁸U and ⁴⁰K is found to be lower. The absorbed dose rates due to these radionuclides were calculated. The average absorbed gamma dose rate due to the presence of ²³²Th, ²³⁸U and ⁴⁰K in soil samples is 36.99 nGy·h⁻¹. These results indicate no radiological anomaly. The data presented in this study will serve as a base line survey for primordial radionuclides concentration in the study area.

Key words Natural radioactivity, Soil samples, Salt field, Gamma-ray spectrometer

CLC numbers P619.21+1, X125

1 Introduction

Mankind has all along lived in natural radiation environment due to various natural sources of terrestrial and extra terrestrial origin. The extra terrestrial sources of radiation are cosmic rays and other radioactive elements produced due to the interaction of cosmic rays with the atmosphere. The sources of terrestrial radiation are radionuclide present in the earth's crust or in the atmosphere. Most of the terrestrial radionuclides originate from ²³⁸U and ²³²Th series. They also originate from the singly occurring primordial radionuclide ⁴⁰K. This radionuclide may enter into our food chain and cause internal exposure. The distribution of natural radionuclides is an essential prerequisite for the evaluation and control of public exposures. Gamma radiation from these radionuclides represents the main external source of radiation to the human

body. Natural environmental radioactivity and the associated external exposure due to gamma radiation depend primarily on the geological and geographical conditions, and appear at different levels in the soils of each different geological region^[1]. The specific levels of terrestrial environmental radiation are related to the geological composition of each lithologically separated area, and to the content of thorium, uranium and potassium in the rock from which the soil originate in each area.

Investigations on terrestrial natural radiation have received particular attention world wide and lead to extensive surveys in many countries. They mainly serve as baseline data of natural radioactivity such that man-made contaminations can be detected and quantitatively determined. They can further be used to assess

* Corresponding author. E-mail: ravissn@gmail.com, ravi@ssnce.ac.in

Received date: 2007-07-16

public dose rates and to perform epidemiological studies. The results obtained in each country can be exploited to enrich the world's data bank which is needed for evaluating worldwide average values of radiometric and dosimetric quantities.

The gamma ray spectrometry method for the determination of naturally occurring radioactive materials (NORM) is of great interest in Earth sciences. There are a number of possible applications spanning from ore exploration to environmental radiation monitoring problems, most of them involving the determination of the U, Th and K in amount in soils, sediments and rocks ^[2].

Mother Nature has gifted mankind with lot of precious gifts. Common salt is one of them. In the globe, Tamilnadu is one of the ideal locations for producing salt. Kelambakkam salt field area is one of the leading producers of salt in global market. The climate, soil and availability of brine are a great asset for producing quality salts. This salt field consists of nearly 500 acres of saltpans in the town of Kelambakkam very near to the East coast of Tamilnadu and connected to the Information Technology highway ^[3].

Salt-field, a closed salt-production unit, is composed of a basin for the first pumping for water reserves, five-grade evaporation basins, crystallization basins and collector basin for effluent discharges. During high tide, salters let the water to the salt-field from the main sea channel through the conduit sluice gate. Here it flows from one evaporation basin to the other along little channels and through sluice gates with the aid of wind and hand pumps. Table salt begins to separate from its crystallization basins. This is the time for the salt to be harvested and stored. During low tide, salters let effluent discharges and rainwater into the channel and back into the sea through the drain sluice gate. This salt-production unit also includes the salt-pans house and the main channel for the inflow of seawater.

In the present work, the primordial radionuclides concentration in soil samples collected in and around the salt field area, Kelambakkam, Tamilnadu was measured using gamma ray spectrometer.

2 Sample collection

The soil profiles were sampled using an auto-

matic core driller. Cores extruded and sectioned at 10 cm diameter and 25 cm depth were used to take soil samples. The soil samples were collected at six sites, each covering 1m² area. After making 1m×1m area and removing the top layer of the soil, five pits were dug and from each pit the soil samples were collected, mixed and packed. This ensures uniform mixing of the soil at the particular site. The locations of the sampling sites were randomized to avoid any bias in results. The collected samples after coning, quartering and sieving were used for different analyses. These samples were weighed individually. Nearly 2 kg of the samples were air dried for 10 days and kept in an oven at 105°C. After homogenization samples were sieved through 100-mesh sieve. These fine samples were used for the radioactivity analysis. The salt samples were also collected for analysis. Sample S₇ was collected slightly away from the salt field area and was used for reference in the analysis.

3 Materials and methods

The selected known amount of samples were packed in plastic containers and aged for four weeks to allow the parent nuclides in the samples to reach equilibrium with their respective progeny ^[4]. All the soil samples were subjected to gamma ray spectral analysis with a counting time of 20,000 s. A 3"×3" NaI (TI) detector was employed with adequate lead shielding which reduced the background by a factor of about 95. The efficiency for various energies was arrived at using IAEA standard sources and required geometry. The system was calibrated both in terms of energy response and also for counting efficiency. The concentrations of various nuclides of interest are determined in Bq/kg using the count spectra. The peaks corresponding to 1.46 MeV (⁴⁰K), 1.76 MeV (²¹⁴Bi) and 2.614 MeV (²⁰⁸Tl) were considered in arriving at the activity levels.

At each sampling spot the ambient gamma dose was measured using a digital Environmental Radiation Dosimeter (ERDM), which comprises of a Geiger Muller (G.M) counter. Before starting the survey, the ERDM was calibrated regularly using a standard source. The ERDM readings were recorded at 1 m above ground level. A total of five readings were

Download English Version:

<https://daneshyari.com/en/article/1848471>

Download Persian Version:

<https://daneshyari.com/article/1848471>

[Daneshyari.com](https://daneshyari.com)