



Graviton as a pair of collinear gauge bosons



Stephan Stieberger^{a,*}, Tomasz R. Taylor^b

^a Max-Planck-Institut für Physik, Werner-Heisenberg-Institut, 80805 München, Germany

^b Department of Physics, Northeastern University, Boston, MA 02115, USA

ARTICLE INFO

Article history:

Received 23 September 2014

Accepted 24 October 2014

Available online 30 October 2014

Editor: L. Alvarez-Gaumé

ABSTRACT

We show that the mixed gravitational/gauge superstring amplitudes describing decays of massless closed strings – gravitons or dilatons – into a number of gauge bosons, can be written at the tree (disk) level as linear combinations of pure open string amplitudes in which the graviton (or dilaton) is replaced by a pair of collinear gauge bosons. Each of the constituent gauge bosons carry exactly one half of the original closed string momentum, while their ± 1 helicities add up to ± 2 for the graviton or to 0 for the dilaton.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

Quantization of gravitational waves yields gravitons: massless spin 2 particles with two polarized degrees of freedom (helicity $+2 \equiv ++$ and $-2 \equiv --$) in four dimensions. While the existence of gravitational waves is well established, the detection of individual gravitons may be impossible due to extremely low cross sections. Nevertheless, theoretical understanding of gravitons and their interactions is a prerequisite for constructing a viable theory of quantum gravity.

Superstring theory offers an interesting insight into gravitons. In this framework, they appear as zero modes of *closed* strings. On the other hand, it is known that zero modes of *open* strings give rise to spin 1 gauge bosons. With the closed string seen as a loop of two open strings connected at both ends, graviton appears to be a “bound state” of two vector bosons. This is also suggested by the form of graviton vertex operator: in type II superstring theory, it is a product of two spin 1 vertex operators (from the left- and right-moving sectors of world-sheet excitations). Helicity $++$ appears as a superposition of two helicity $+$ states while helicity $--$ comes as a superposition of two helicity $-$ states. In addition, the products $+-$ and $-+$ create two degrees of freedom of the scalar (complex) superstring dilaton.

In 1985, Kawai, Lewellen and Tye (KLT) [1] derived a formula which expresses any closed string tree amplitude in terms of a sum of the products of appropriate open string tree amplitudes. At the level of zero modes, KLT relations allow expressing the graviton and dilaton amplitudes in terms of products of gauge boson amplitudes. The existence of such relations means that, at least in the leading order of perturbation theory, the content of Einstein's

gravity is encoded in Yang–Mills (YM) theory. The quadratic form of KLT relations is perfectly consistent with the heuristic picture of a closed string as a loop of two open strings. In fact, string field theory suggests a similar description [2]. This does not help, however, in answering the question of whether the graviton can be considered as a pair of gauge bosons beyond the world-sheet, as an actual bound state in physical space-time. One alternative description has been developed in [3,4], by constructing closed superstring amplitudes through the “single-valued” projection of open superstring amplitudes. This projection yields linear relations between the functions encompassing effects of massive closed and open superstring excitations, to all orders in the inverse string tension α' . They reveal a deeper connection between gauge and gravity string amplitudes than what is implied by the KLT relations, but they do not provide new insight into their $\alpha' \rightarrow 0$ field theory limit.

In this Letter, we present a *linear* relation between the amplitude for the decay of one massless closed string state, i.e. a graviton or a dilaton, into an arbitrary number $N - 2$ of gauge bosons and a sum of purely open string amplitudes involving N gauge bosons. The sum involves so-called partial amplitudes associated to particular gauge group factors. The original closed string state is replaced by two vector bosons, each of them carrying exactly one half of its momentum, and its helicity is split in the same way as in string vertex operators. In the forthcoming publication [5], we will show that in all open and closed string amplitudes, gravitons and dilatons can be replaced by pairs of such collinear vectors bosons.

Although our derivation utilizes full-fledged Type II superstring theory, it is instructive to discuss the field theory limit (i.e. the zero slope $\alpha' = 0$ limit of Regge trajectories) of mixed

* Corresponding author.

E-mail address: stephan.stieberger@mpp.mpg.de (S. Stieberger).

gravitational/gauge interactions. This limit is described by Einstein–Yang–Mills (EYM) theory coupled to the dilaton.¹ All tree level amplitudes can be constructed by using the recursion relations derived by Britto, Cachazo, Feng and Witten (BCFW) [8], with the basic building blocks provided by the following three-point amplitudes:

$$\begin{aligned} A(1^{--}, 2^{--}, 3^{++}) &= \frac{\langle 12 \rangle^6}{\langle 23 \rangle^2 \langle 31 \rangle^2}, \\ A(1^{--}, 2^{+-}, 3^{-+}) &= \frac{\langle 12 \rangle^2 \langle 13 \rangle^2}{\langle 23 \rangle^2}, \\ A(1^{+}, 2^{-}, 3^{--}) &= \frac{\langle 23 \rangle^4}{\langle 12 \rangle^2}, \\ A(1^{-}, 2^{-}, 3^{+-}) &= \langle 12 \rangle^2, \end{aligned} \quad (1)$$

where we used superscripts to label helicity states, with $+-$ and $-+$ assigned to the dilaton and its complex conjugate, respectively. We are using standard notation of the helicity formalism, see [9]. The mass dimension (-1) gravitational coupling $\sqrt{\kappa}$ is implied by the above expressions. In addition, three gauge bosons interact with the well known Yang–Mills amplitude

$$A(1^{-}, 2^{-}, 3^{+}) = \frac{\langle 12 \rangle^3}{\langle 23 \rangle \langle 31 \rangle}, \quad (2)$$

where we omitted the (dimensionless) gauge coupling constant.

A good example of an amplitude involving both gravitational and gauge couplings is the amplitude for the graviton decay into three gauge bosons. In this case,

$$A(1^{+}, 2^{+}, 3^{-}; q^{--}) = \frac{(3q)^4}{\langle 12 \rangle \langle 23 \rangle \langle 31 \rangle}, \quad (3)$$

which can be obtained either by using BCFW recursion relations or by a straightforward Feynman diagram calculation. In this Letter, we focus on the amplitudes similar to (3), describing gravitons and dilatons decaying into an arbitrary number of gauge bosons. In string theory, these are disk amplitudes with one closed string vertex insertion on the world-sheet and a number of open strings attached at the boundary.

In order to compute the amplitudes, it is convenient to use the “doubling trick,” to convert disk correlators to the standard holomorphic ones by extending the fields to the entire complex plane [10]. Furthermore, the integration over positions of world-sheet symmetric closed string states (such as graviton or dilaton) can be extended from the half-plane covering the disk to the full complex plane. Open string vertices representing $N - 2$ gauge bosons with momenta p_i , $i = 1, \dots, N - 2$ (in an arbitrary helicity configuration) are inserted on the real axis at x_i , while a single closed string vertex operator, which represents the graviton or dilaton with momentum q , is inserted at complex z . All momenta are restricted to four dimensions, with $p_i^2 = q^2 = 0$ (although the following derivation is independent on the space–time dimension and can be adjusted to massive states). The amplitudes involve integrals of the form

$$\begin{aligned} F_N &= V_{\text{CKG}}^{-1} \delta^{(4)} \left(\sum_{i=1}^{N-2} p_i + q \right) \\ &\times \int \prod_{i=1}^{N-2} dx_i \prod_{1 \leq r < s \leq N-2} |x_r - x_s|^{2\alpha' p_r p_s} (x_r - x_s)^{n_{rs}} \end{aligned}$$

¹ For early work on EYM scattering amplitudes, see Ref. [6]; for more recent work, see [7].

$$\times \int_{\mathbb{C}} d^2 z (z - \bar{z})^n \prod_{i=1}^{N-2} |x_i - z|^{2\alpha' p_i q} (x_i - z)^{n_i} (x_i - \bar{z})^{\bar{n}_i}, \quad (4)$$

where we included the momentum-conserving delta function and divided by the volume V_{CKG} of the conformal Killing group. The powers n_{rs} , n_i , $\bar{n}_i$, n are some integer numbers. To be specific, we focus on the amplitude associated to one particular Chan–Paton factor (partial amplitude), $\text{Tr}(T^1 T^2 \dots T^{N-2})$, with the integral over ordered $x_1 < x_2 < \dots < x_{N-2}$.

The techniques for evaluating generic disk integrals involving both open and closed strings have been developed in [11]. For the concrete case (4), we write the complex integral as an integral over holomorphic and anti-holomorphic coordinates, by following the method proposed in [1]. After writing $z = z_1 + iz_2$, the integrand becomes an analytic function of z_2 with $2(N - 2)$ branch points at $\pm i(x_i - z_1)$. We then deform the z_2 -integral along the real axis $\text{Im}(z_2) = 0$ to the pure imaginary axis $\text{Re}(z_2) = 0$. In this way, the variables

$$\xi = z_1 + iz_2 \equiv z, \quad \eta = z_1 - iz_2 \equiv \bar{z} \quad (5)$$

become real. After changing the integration variables $(z_1, z_2) \rightarrow (\xi, \eta)$ (with the Jacobian $\det \frac{\partial(z_1, z_2)}{\partial(\xi, \eta)} = \frac{1}{2}$), Eq. (4) becomes an integral over N real positions x_i, ξ, η

$$\begin{aligned} F_N &= V_{\text{CKG}}^{-1} \delta^{(4)} \left(\sum_{i=1}^N k_i \right) \int \prod_{i=1}^{N-2} dx_i \int_{-\infty}^{\infty} d\xi \int_{-\infty}^{\infty} d\eta \\ &\times \prod_{1 \leq r < s \leq N-2} |x_r - x_s|^{2\alpha' k_r k_s} (x_r - x_s)^{n_{rs}} \\ &\times \frac{i}{2} (\xi - \eta)^n \prod_{i=1}^{N-2} \Pi(x_i, \xi, \eta) |x_i - \xi|^{2\alpha' k_i k_{N-1}} |x_i - \eta|^{2\alpha' k_i k_N} \\ &\times (x_i - \xi)^{n_i} (x_i - \eta)^{\bar{n}_i}, \end{aligned} \quad (6)$$

with the open string momenta $k_r = p_r$, $r = 1, \dots, N - 2$ and the closed string momentum split in half:

$$k_{N-1} = k_N = \frac{1}{2} q. \quad (7)$$

Eq. (6) resembles a generic open string integral involving N open strings with external momenta k_i supplemented by the extra phase factors

$$\Pi(x_i, \xi, \eta) = e^{2\pi i \alpha' k_i k_N \theta[(x_i - \xi)(x_i - \eta)]}, \quad (8)$$

where θ denotes the Heaviside step function. These monodromy factors (8) account for the correct branch of the integrand, making the integral well defined. Note that the phases, which are independent on the integers $n_{rs}, n_i, \bar{n}_i, n$ do not depend on the particular values of integration variables, but only on the ordering of ξ and η with respect to the original $N - 2$ vertex positions. In this way, the original integral becomes a weighted (by phase factors) sum of integrals, each of them having the same form as the integrals appearing in N -point (partial) open string amplitudes, with the vertices inserted at x_l , $l = 1, \dots, N$, where we identified $x_{N-1} \equiv \xi$ and $x_N \equiv \eta$. Note that the order of the original $N - 2$ positions remains unchanged. Since the graviton as well as dilaton vertices factorize into two gauge bosons inserted at $z = \xi = x_{N-1}$ and $\bar{z} = \eta = x_N$, we conclude that the amplitude $A(1, 2, \dots, N - 2; q)$ describing graviton (or dilaton) decays into $N - 2$ gauge bosons can be written as a weighted sum of pure open string amplitudes with the graviton (or dilaton) replaced by a pair of collinear gauge bosons, each of them carrying exactly one-half of its momentum, cf. Eq. (7).

Download English Version:

<https://daneshyari.com/en/article/1850471>

Download Persian Version:

<https://daneshyari.com/article/1850471>

[Daneshyari.com](https://daneshyari.com)