



# Asymptotically flat charged rotating dilaton black holes in higher dimensions

A. Sheykhi<sup>a,b,\*</sup>, M. Allahverdizadeh<sup>a</sup>, Y. Bahrampour<sup>c</sup>, M. Rahnema<sup>a</sup>

<sup>a</sup> Department of Physics, Shahid Bahonar University, PO Box 76175, Kerman, Iran

<sup>b</sup> Research Institute for Astronomy and Astrophysics of Maragha (RIAAM), Maragha, Iran

<sup>c</sup> Department of Mathematics, Shahid Bahonar University, Kerman, Iran

## ARTICLE INFO

### Article history:

Received 1 June 2008

Received in revised form 24 June 2008

Accepted 26 June 2008

Available online 5 July 2008

Editor: M. Cvetič

## ABSTRACT

We find a class of asymptotically flat slowly rotating charged black hole solutions of Einstein–Maxwell–dilaton theory with arbitrary dilaton coupling constant in higher dimensions. Our solution is the correct one generalizing the four-dimensional case of Horne and Horowitz [J.H. Horne, G.T. Horowitz, Phys. Rev. D 46 (1992) 1340]. In the absence of a dilaton field, our solution reduces to the higher-dimensional slowly rotating Kerr–Newman black hole solution. The angular momentum and the gyromagnetic ratio of these rotating dilaton black holes are computed. It is shown that the dilaton field modifies the gyromagnetic ratio of the black holes.

© 2008 Elsevier B.V. All rights reserved.

## 1. Introduction

General Relativity in higher dimensions has been the subject of increasing attention in recent years. There are several motivations for studying higher-dimensional Einstein's theory, and in particular its black hole solutions. First of all, string theory contains gravity and requires more than four dimensions. In fact, the first successful statistical counting of black hole entropy in string theory was performed for a five-dimensional black hole [1]. This example provides the best laboratory for the microscopic string theory of black holes. Besides, the production of higher-dimensional black holes in future colliders becomes a conceivable possibility in scenarios involving large extra dimensions and TeV-scale gravity. Furthermore as mathematical objects, black hole spacetimes are among the most important Lorentzian Ricci-flat manifolds in any dimension. While the non-rotating black hole solution to the higher-dimensional Einstein–Maxwell gravity was found several decades ago [2], the counterpart of the Kerr–Newman solution in higher dimensions, that is the charged generalization of the Myers–Perry solution [3] in higher-dimensional Einstein–Maxwell theory, still remains to be found analytically. Indeed, the case of charged rotating black holes in higher dimensions has been discussed in the framework of supergravity theories and string theory [4–6]. Recently, charged rotating black hole solutions in higher dimensions with a single rotation parameter

in the limit of slow rotation has been constructed in [7] (see also [8,9]).

On the other hand, a scalar field called the dilaton appears in the low energy limit of string theory. The presence of the dilaton field has important consequences on the causal structure and the thermodynamic properties of black holes. Thus much interest has been focused on the study of the dilaton black holes in recent years. While exact static dilaton black hole solutions of Einstein–Maxwell–dilaton (EMd) gravity have been constructed by many authors (see e.g. [10–13]), exact rotating dilaton black hole solutions have been obtained only for some limited values of the dilaton coupling constant [14–16]. For general dilaton coupling constant, the properties of charged rotating dilaton black holes only with infinitesimally small charge [17] or small angular momentum in four [18–20] and five dimensions have been investigated [21]. Recently, one of us has constructed a class of charged slowly rotating dilaton black hole solutions in arbitrary dimensions [22]. However, in contrast to the four-dimensional Horne and Horowitz solution, these solutions [22] have unusual asymptotics. They are neither asymptotically flat nor (A)dS. Besides, they are ill-defined for the string case where  $\alpha = 1$ . The purpose of the present Letter is to generalize the four-dimensional Horne and Horowitz solution with sensible asymptotics, to arbitrary dimensions. These asymptotically flat solutions describe an electrically charged, slowly rotating dilaton black hole with an arbitrary value of the dilaton coupling constant in various dimensions. It is worth noting that in this Letter, we restrict ourselves to the rotation in one plane, so our black hole has only one angular momentum parameter. We also investigate the effects of the dilaton field on the angular momentum and the gyromagnetic ratio of these rotating dilaton black holes.

\* Corresponding author at: Department of Physics, Shahid Bahonar University, PO Box 76175, Kerman, Iran.

E-mail addresses: sheykhi@mail.uk.ac.ir (A. Sheykhi), bahram@mail.uk.ac.ir (Y. Bahrampour), Majid.Rahnema@mail.uk.ac.ir (M. Rahnema).

## 2. Field equations and solutions

We consider the  $n$ -dimensional ( $n \geq 4$ ) theory in which gravity is coupled to dilaton and Maxwell field with an action

$$S = \frac{1}{16\pi} \int_{\mathcal{M}} d^n x \sqrt{-g} \left( R - \frac{4}{n-2} \partial_\mu \Phi \partial^\mu \Phi - e^{-\frac{4\alpha\Phi}{n-2}} F_{\mu\nu} F^{\mu\nu} \right) - \frac{1}{8\pi} \int_{\partial\mathcal{M}} d^{n-1} x \sqrt{-h} \Theta(h), \quad (1)$$

where  $R$  is the scalar curvature,  $\Phi$  is the dilaton field,  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  is the electromagnetic field tensor, and  $A_\mu$  is the electromagnetic potential.  $\alpha$  is an arbitrary constant governing the strength of the coupling between the dilaton and the Maxwell field. The last term in Eq. (1) is the Gibbons–Hawking boundary term which is chosen such that the variational principle is well-defined. The manifold  $\mathcal{M}$  has metric  $g_{\mu\nu}$  and covariant derivative  $\nabla_\mu$ .  $\Theta$  is the trace of the extrinsic curvature  $\Theta^{ab}$  of any boundary  $\partial\mathcal{M}$  of the manifold  $\mathcal{M}$ , with induced metric  $h_{ab}$ . The equations of motion can be obtained by varying the action (1) with respect to the gravitational field  $g_{\mu\nu}$ , the dilaton field  $\Phi$  and the gauge field  $A_\mu$  which yields the following field equations

$$R_{\mu\nu} = \frac{4}{n-2} \partial_\mu \Phi \partial_\nu \Phi + 2e^{-\frac{4\alpha\Phi}{n-2}} \left( F_{\mu\eta} F_\nu^\eta - \frac{1}{2(n-2)} g_{\mu\nu} F_{\lambda\eta} F^{\lambda\eta} \right), \quad (2)$$

$$\nabla^2 \Phi = -\frac{\alpha}{2} e^{-\frac{4\alpha\Phi}{n-2}} F_{\lambda\eta} F^{\lambda\eta}, \quad (3)$$

$$\partial_\mu (\sqrt{-g} e^{-\frac{4\alpha\Phi}{n-2}} F^{\mu\nu}) = 0. \quad (4)$$

We would like to find  $n$ -dimensional rotating solutions of the above field equations. For small rotation, we can solve Eqs. (2)–(4) to first order in the angular momentum parameter  $a$ . Inspection of the  $n$ -dimensional Kerr solutions shows that the only term in the metric that changes to the first order of the angular momentum parameter  $a$  is  $g_{t\phi}$ . Similarly, the dilaton field does not change to  $O(a)$  and  $A_\phi$  is the only component of the vector potential that changes. Therefore, for infinitesimal angular momentum we assume the metric being of the following form

$$ds^2 = -U(r) dt^2 + \frac{dr^2}{W(r)} - 2af(r) \sin^2 \theta dt d\phi + r^2 R^2(r) (d\theta^2 + \sin^2 \theta d\phi^2 + \cos^2 \theta d\Omega_{n-4}^2), \quad (5)$$

where  $d\Omega_{n-4}^2$  denotes the metric of a unit  $(n-4)$ -sphere. The functions  $U(r)$ ,  $W(r)$ ,  $R(r)$  and  $f(r)$  should be determined. In the particular case  $a = 0$ , this metric reduces to the static and spherically symmetric cases. For small  $a$ , we can expect to have solutions with  $U(r)$  and  $W(r)$  still functions of  $r$  alone. The  $t$  component of the Maxwell equations can be integrated immediately to give

$$F_{tr} = \sqrt{\frac{U(r)}{W(r)}} \frac{Q e^{\frac{4\alpha\Phi}{n-2}}}{(rR)^{n-2}}, \quad (6)$$

where  $Q$ , an integration constant, is the electric charge of the black hole. In general, in the presence of rotation, there is also a vector potential in the form

$$A_\phi = -aQC(r) \sin^2 \theta. \quad (7)$$

Asymptotically flat static ( $a = 0$ ) black hole solutions of the above field equations was found in [23]. Here we are looking for the asymptotically flat solutions in the case  $a \neq 0$ . Our strategy for obtaining the solution is the perturbative method suggested in [18]. Inserting the metric (5), the Maxwell fields (6) and (7) into the

field equations (2)–(4), one can show that the static part of the metric leads to the following solutions [23]

$$U(r) = \left[ 1 - \left( \frac{r_+}{r} \right)^{n-3} \right] \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^{1-\gamma(n-3)}, \quad (8)$$

$$W(r) = \left[ 1 - \left( \frac{r_+}{r} \right)^{n-3} \right] \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^{1-\gamma}, \quad (9)$$

$$R(r) = \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^{\gamma/2}, \quad (10)$$

$$\Phi(r) = \frac{n-2}{4} \sqrt{\gamma(2+3\gamma-n\gamma)} \ln \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right], \quad (11)$$

while the rotating part of the metric admits a solution

$$f(r) = (n-3) \left( \frac{r_+}{r} \right)^{n-3} \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^{\frac{n-3-\alpha^2}{n-3+\alpha^2}} + \frac{(\alpha^2 - n + 1)(n-3)^2}{\alpha^2 + n - 3} r_-^{n-3} r^2 \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^\gamma \times \int \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^{\gamma(2-n)} \frac{dr}{r^n}, \quad (12)$$

$$C(r) = \frac{1}{r^{n-3}}. \quad (13)$$

We can also perform the integration and express the solution in terms of hypergeometric function

$$f(r) = (n-3) \left( \frac{r_+}{r} \right)^{n-3} \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^{\frac{n-3-\alpha^2}{n-3+\alpha^2}} + \frac{(\alpha^2 - n + 1)(n-3)^2}{(1-n)(\alpha^2 + n - 3)} \left( \frac{r_-}{r} \right)^{n-3} \left[ 1 - \left( \frac{r_-}{r} \right)^{n-3} \right]^\gamma \times {}_2F_1 \left( \left[ (n-2)\gamma, \frac{n-1}{n-3} \right], \left[ \frac{2n-4}{n-3} \right], \left( \frac{r_-}{r} \right)^{n-3} \right). \quad (14)$$

Here  $r_+$  and  $r_-$  are the event horizon and Cauchy horizon of the black hole, respectively. The constant  $\gamma$  is

$$\gamma = \frac{2\alpha^2}{(n-3)(n-3+\alpha^2)}. \quad (15)$$

The charge  $Q$  is related to  $r_+$  and  $r_-$  by

$$Q^2 = \frac{(n-2)(n-3)^2}{2(n-3+\alpha^2)} r_+^{n-3} r_-^{n-3}, \quad (16)$$

and the physical mass of the black hole is obtained as follows [24]

$$M = \frac{\Omega_{n-2}}{16\pi} \left[ (n-2)r_+^{n-3} + \frac{n-2-p(n-4)}{p+1} r_-^{n-3} \right], \quad (17)$$

where  $\Omega_{n-2}$  denotes the area of the unit  $(n-2)$ -sphere and the constant  $p$  is

$$p = \frac{(2-n)\gamma}{(n-2)\gamma - 2}. \quad (18)$$

The metric corresponding to (8)–(14) is asymptotically flat. In the special case  $n = 4$ , the static part of our solution reduces to

$$U(r) = W(r) = \left( 1 - \frac{r_+}{r} \right) \left( 1 - \frac{r_-}{r} \right)^{\frac{1-\alpha^2}{1+\alpha^2}}, \quad (19)$$

$$R(r) = \left( 1 - \frac{r_-}{r} \right)^{\frac{\alpha^2}{1+\alpha^2}}, \quad (20)$$

$$\Phi(r) = \frac{\alpha}{(1+\alpha^2)} \ln \left( 1 - \frac{r_-}{r} \right), \quad (21)$$

Download English Version:

<https://daneshyari.com/en/article/1852329>

Download Persian Version:

<https://daneshyari.com/article/1852329>

[Daneshyari.com](https://daneshyari.com)