

available at www.sciencedirect.comjournal homepage: <http://www.rpor.eu/>

Craniospinal radiotherapy in children: Electron- or photon-based technique of spinal irradiation

Marzanna Chojnacka^{a,*}, Anna Skowrońska-Gardas^a, Marzena Morawska-Kaczyńska^b, Anna Zygmuntowicz-Piętko^b, Katarzyna Pędziwiatr^a, Anna Semaniak^b

^a Department of Radiotherapy, Maria Skłodowska Curie Memorial Cancer Centre – Institute, Wawelska 15, 00-973 Warsaw, Poland

^b Department of Medical Physics, Maria Skłodowska Curie Memorial Cancer Centre – Institute, Wawelska 15, 00-973 Warsaw, Poland

ARTICLE INFO

Keywords:

Craniospinal irradiation

Supine position

Late toxicity

ABSTRACT

Background: The prone position and electron-based technique for craniospinal irradiation (CSI) have been standard in our department for many years. But this immobilization is difficult for the anaesthesiologist to gain airway access. The increasing number of children treated under anaesthesia led us to reconsider our technique.

Aim: The purpose of this study is to report our new photon-based technique for CSI which could be applied in both the supine and the prone position and to compare this technique with our electron-based technique.

Materials and methods: Between November 2007 and May 2008, 11 children with brain tumours were treated in the prone position with CSI. For 9 patients two treatment plans were created: the first one using photons and the second one using electron beams for spinal irradiation. We prepared seven 3D-conformal photon plans and four forward planned segmented field plans. We compared 20 treatment plans in terms of target dose homogeneity and sparing of organs at risk.

Results: In segmented field plans better dose homogeneity in the thecal sac volume was achieved than in electron-based plans. Regarding doses in organs at risk, in photon-based plans we obtained a lower dose in the thyroid but a higher one in the heart and liver.

Conclusions: Our technique can be applied in both the supine and prone position and it seems to be more feasible and precise than the electron technique. However, more homogeneous target coverage and higher precision of dose delivery for photons are obtained at the cost of slightly higher doses to the heart and liver.

© 2009 Wielkopolskie Centrum Onkologii. Published by Elsevier Urban & Partner Sp. z o.o.

All rights reserved.

1. Background

The entire craniospinal axis is the standard radiation volume for the treatment of primary brain tumours with a risk of leptomeningeal spread.¹ Many of these tumours are found in young children who may require anaesthesia during their therapy. The prone position during CSI was standard for many

years. But this therapeutic immobilization is difficult for the anaesthesiologist to gain airway access. Some authors report that the supine position is simple and together with airway access provides other advantages. It is more comfortable and more reproducible.^{2–5} But this position constrains application to only the photon-based technique for spine irradiation. In young children, especially, severe toxicity is of our interest. Adverse effects connected with neuraxis irradiation include

* Corresponding author. Tel.: +48 225709186; fax: +48 225709186.

E-mail address: marzanna.ch1@wp.pl (M. Chojnacka).

neuroendocrine and neurocognitive dysfunctions, but in addition, the exit dose of the photon beam for spinal irradiation is potentially associated with late cardiac, pulmonary and bone damage.^{2,3,6–9} Electrons are more conformal and potentially spinal irradiation with electrons is more protective for structures outside the target volume on the beam path.^{3,10}

The CSI technique in our department has evolved over the last decades. High-energy electrons for spine irradiation have been used since 1986 and this 2D technique and treatment results in 158 children with medulloblastoma were published previously.¹¹ In the next report we presented the outcome and patterns of failure for 95 patients with medulloblastoma treated with 3D-conformal radiotherapy.¹² New equipment installed in our department (accelerators with CBCT and TPS with forward IMRT planning) together with an increasing number of young children treated under anaesthesia led us to reconsider our technique of craniospinal irradiation. Therefore we propose a new, photon-based technique with characteristic features: two isocentres, intrafraction moving beam junction, and oblique beams in the thyroid region between lateral cranial and posterior spinal beams.

2. Aim

The aim of this study is to report our new photon-based technique for CSI which could be applied in both the supine and the prone position and to compare this technique with the old electron-based technique. The treatment plans were compared in terms of dose homogeneity in the target volume and sparing of organs at risk.

3. Materials and methods

During 6 months, between November 2007 and May 2008, 11 children, aged 4–13 years (median 6 years), were treated with craniospinal irradiation. The diagnoses included medulloblastoma/PNET in 7, anaplastic ependymoma in 2 and atypical teratoid rhabdoid tumour in 2 patients. Patients received 25.05 or 35.07 Gy CSI in 15 or 21 fractions. The primary tumour site was subsequently boosted to the dose of 54–55.11 Gy. All children were immobilized in the prone position with an individual thermoplastic mask for the head. For 9 patients two treatment plans for spinal irradiation were created: the first one using photons and the second one using 18–22 MeV electron beams. In two cases the spine was localized too deep for the electron beam and these patients had only photon treatment plans.

The treatment planning was based on the 3 mm spaced CT slices from the top of the head to the level of the aortic arch and 9 mm elsewhere. The brain and the whole length of the thecal sac were contoured and defined as the CTV_{brain} and CTV_{spine}. PTV_{brain} encompassed a 4 mm margin around the CTV_{brain} in all directions. In PTV_{spine} the lateral margin was set to 10 mm because of the possibility of a child moving. Organs at risk that we additionally outlined were the thyroid, heart, lungs, liver, kidneys and vertebral body.

Our electron technique was reported previously.^{11,12} Our new photon-based technique is a two-isocentre treatment technique which could be applied for both the supine and

prone position. We performed two methods of planning: conformal 3D (for 7 patients) and forward planned segmented fields (for 4 patients). In our photon-based technique the brain with the upper cervical part of the spine (usually to the C3 level) was treated with 6 MV photon lateral beams with zero collimator angle. Custom blocks together with the MLC were used in both cranial fields to improve shielding of eye lens and the target conformity in the vicinity of the cribriform plate. The spine besides the thyroid region was treated with a 6 MV posterior photon beam. The width of the spinal fields was comparable with the width of the electron fields, which was about 5.5 cm. Between cranial and spinal beams two additional oblique fields were added in the thyroid region in order to reduce the dose to the thyroid gland. Such beam arrangement also helped to avoid irradiation of the mandible and oral cavity. These 15 MV photon beams, isocentric with cranial beams, covered the part of the spine in the vicinity of the thyroid gland. Gantry angles for cervical fields were 70° and 290°. The edges of these fields matched cranial fields from one side and the posterior spinal field from the other.

To ensure an even dose distribution at the match lines intrafraction feathering technique was used with two- and three-level intrafraction junctions for conformal 3D and forward planned segmented fields plans, respectively. Our standard CSI treatment plan with segmented fields comprised in total five beams. This number doubled for conformal 3D planning with two-level junctions. In our standard CSI treatment plan with segmented fields small weight segments were added to the cranial and spinal beams to improve dose homogeneity within the spine volume and to reduce the maximum dose to organs at risk. Four, three and five segments existed typically in each cranial, cervical and spinal field, respectively.

We compared target dose homogeneity and sparing of organs at risk for 20 treatment plans (11 photon and 9 electron). For all plans we calculated the standard deviation of the dose distribution in CTV_{spine} and PTV_{spine} for the total dose and the average doses to organs at risk as the percentage of the average dose in the CTV_{spine}. Additionally we compared the average values of standard deviation for photon conformal 3D and forward planned segmented field plans.

4. Results

Standard deviations of dose distribution in the spine are presented in Tables 1 and 2.

In the photon f-IMRT plans for the spine the dose homogeneity in CTV and PTV was better than in electron-based plans. However, the comparison of the values of the standard deviations of dose distribution in the spine does not reflect the existing radiobiological differences between plans. The photon beam junctions were moved within each fraction while the electron beam junctions were changed once during the whole CNS treatment, after the first half of the therapy. Better dose homogeneity in the spine for photons is radiobiologically enhanced by the introduction of a three-level intrafraction beam junction.

The average doses in the organs at risk are presented in Table 3.

Download English Version:

<https://daneshyari.com/en/article/1854273>

Download Persian Version:

<https://daneshyari.com/article/1854273>

[Daneshyari.com](https://daneshyari.com)