



Separability criteria with angular and Hilbert space averages



Kazuo Fujikawa^{a,b,*}, C.H. Oh^{b,c}, Koichiro Umetsu^d, Sixia Yu^c

^a Quantum Hadron Physics Laboratory, RIKEN Nishina Center, Wako 351-0198, Japan

^b Institute of Advanced Studies, Nanyang Technological University, Singapore 639798, Singapore

^c Centre for Quantum Technologies, National University of Singapore, Singapore 117543, Singapore

^d Laboratory of Physics, College of Science and Technology, Nihon University, Funabashi 274-8501, Japan

ARTICLE INFO

Article history:

Received 29 December 2015

Accepted 9 February 2016

Available online 18 February 2016

Keywords:

Separability

Entanglement

Two-particle correlation

ABSTRACT

The practically useful criteria of separable states $\rho = \sum_k w_k \rho_k$ in $d = 2 \times 2$ are discussed. The equality $G(\mathbf{a}, \mathbf{b}) = 4[\langle \psi | P(\mathbf{a}) \otimes P(\mathbf{b}) | \psi \rangle - \langle \psi | P(\mathbf{a}) \otimes \mathbf{1} | \psi \rangle \langle \psi | \mathbf{1} \otimes P(\mathbf{b}) | \psi \rangle] = 0$ for any two projection operators $P(\mathbf{a})$ and $P(\mathbf{b})$ provides a necessary and sufficient separability criterion in the case of a separable pure state $\rho = |\psi\rangle\langle\psi|$. We propose the separability criteria of mixed states, which are given by $\text{Tr} \rho \{ \mathbf{a} \cdot \boldsymbol{\sigma} \otimes \mathbf{b} \cdot \boldsymbol{\sigma} \} = (1/3)C \cos \varphi$ for two spin $1/2$ systems and $4\text{Tr} \rho \{ P(\mathbf{a}) \otimes P(\mathbf{b}) \} = 1 + (1/2)C \cos 2\varphi$ for two photon systems, respectively, after taking a geometrical angular average of \mathbf{a} and \mathbf{b} with fixed $\cos \varphi = \mathbf{a} \cdot \mathbf{b}$. Here $-1 \leq C \leq 1$, and the difference in the numerical coefficients $1/2$ and $1/3$ arises from the different rotational properties of the spinor and the transverse photon. If one instead takes an average over the states in the $d = 2$ Hilbert space, the criterion for two photon systems is replaced by $4\text{Tr} \rho \{ P(\mathbf{a}) \otimes P(\mathbf{b}) \} = 1 + (1/3)C \cos 2\varphi$. Those separability criteria are shown to be very efficient using the existing experimental data of Aspect et al. in 1981 and Sakai et al. in 2006. When the Werner state is applied to two photon systems, it is shown that the Hilbert space average can judge its inseparability but not the geometrical angular average.

© 2016 Elsevier Inc. All rights reserved.

* Corresponding author at: Quantum Hadron Physics Laboratory, RIKEN Nishina Center, Wako 351-0198, Japan.
E-mail address: fujikawa@phys.s.u-tokyo.ac.jp (K. Fujikawa).

1. Introduction

To characterize the entanglement, the notions of separability and inseparability, which are the characteristic properties of state vectors in quantum mechanics, are commonly used. On the other hand, the notion of local realism based on local non-contextual hidden-variables models is also used to test the properties such as locality and reduction [1,2]. Local hidden-variables models are generally different from quantum mechanics, and thus local realism tests the deviation from quantum mechanics also.

In the experimental study of local realism, which is commonly tested by CHSH inequality [2], it is customary to first confirm the consistency of the measured basic correlation such as the spin correlation $(\mathbf{a} \cdot \boldsymbol{\sigma} \otimes \mathbf{b} \cdot \boldsymbol{\sigma})$ with quantum mechanics and then test the CHSH inequality. If one confirms the consistency of $(\mathbf{a} \cdot \boldsymbol{\sigma} \otimes \mathbf{b} \cdot \boldsymbol{\sigma})$ for any unit vectors \mathbf{a} and \mathbf{b} with quantum mechanics, one can naturally apply the criterion of quantum mechanical separability to the correlation. In the present study, we discuss this aspect of separability test in quantum mechanics. It is well known that the Peres criterion of the positivity of partial transposed density matrix gives a necessary and sufficient condition of separability of general density matrix in $d = 2 \times 2$ [3] which we study in the present paper. However, it is also well-known from the days of Pauli [4] that the reconstruction of the state vector or density matrix from measured data is in general very involved [5]. It is thus practically useful to derive simpler criteria which do not require the precise state reconstruction. The purpose of the present paper is to derive such separability criteria. We assume the most general separable density matrix but we use a limited set of two-point correlations and thus obtain only the necessary condition of separability in general. Nevertheless, we illustrate that our criteria are very useful when applied to the past experimental data of Aspect et al. in 1981 [6] and Sakai et al. in 2006 [7].

To be specific, we study the general separable quantum mechanical states

$$\rho = \sum_k w_k \rho_k, \tag{1}$$

in $d = 2 \times 2 = 4$; all the states ρ_k are separable pure quantum states [8]. If the state is a separable pure state $\rho = |\psi\rangle\langle\psi|$, separability in (1) is quantified by the equality, which is necessary and sufficient,

$$G(\mathbf{a}, \mathbf{b}) \equiv 4[\langle\psi|P(\mathbf{a}) \otimes P(\mathbf{b})|\psi\rangle - \langle\psi|P(\mathbf{a}) \otimes \mathbf{1}|\psi\rangle\langle\psi|\mathbf{1} \otimes P(\mathbf{b})|\psi\rangle] = 0, \tag{2}$$

for arbitrary two projection operators $P(\mathbf{a})$ and $P(\mathbf{b})$.

In the case of a general mixed ρ , we derive the useful criteria of separable mixed states, namely

$$\text{Tr}\rho\{\mathbf{a} \cdot \boldsymbol{\sigma} \otimes \mathbf{b} \cdot \boldsymbol{\sigma}\} = (1/3)C \cos \varphi, \tag{3}$$

for two spin 1/2 systems, and

$$4\text{Tr}\rho\{P(\mathbf{a}) \otimes P(\mathbf{b})\} = 1 + (1/2)C \cos 2\varphi, \tag{4}$$

for two photon systems with linear polarization projectors $P(\mathbf{a})$ and $P(\mathbf{b})$, respectively: Here $-1 \leq C \leq 1$. The basic new ingredient in our derivation of these criteria (3) and (4) is that we take a geometric *angular average* of unit vectors \mathbf{a} and \mathbf{b} with fixed $\cos \varphi = \mathbf{a} \cdot \mathbf{b}$. It is shown that this angular averaging, which is originally motivated by the specific experiment [7], does not add a new burden to measurements by analyzing the existing experimental data. The difference in the numerical coefficients 1/3 and 1/2 in (3) and (4) arises from the difference in rotational properties; the spinor rotational freedom is 3-dimensional, which agrees with the freedom of the $d = 2$ Hilbert space, while the rotational freedom of the photon is two-dimensional, which differs from the freedom of the $d = 2$ Hilbert space since it is confined in a plane perpendicular to the momentum direction. If one instead takes an average over the states in the $d = 2$ Hilbert space, the formula for the photon is replaced by

$$4\text{Tr}\rho\{P(\mathbf{a}) \otimes P(\mathbf{b})\} = 1 + (1/3)C \cos 2\varphi. \tag{5}$$

This difference between the geometrical angular average and the Hilbert space average is interesting, and it has an interesting implication on the separability issue of the Werner state [8] which accommodates a specific local hidden-variables representation and thus satisfies CHSH inequality. In the case of photon, the Hilbert space average can judge the inseparability of the Werner state but the geometrical angular average cannot.

Download English Version:

<https://daneshyari.com/en/article/1854484>

Download Persian Version:

<https://daneshyari.com/article/1854484>

[Daneshyari.com](https://daneshyari.com)