

Whole body MR in patients with multiple myeloma

Alina PIEKAREK, Piotr SOSNOWSKI, Adam NOWICKI, Mieczysław KOMARNICKI

Received: 11.05.2009
Accepted: 13.07.2009
Subject: original article

Clinical Radiology Department

Address for correspondence:
Alina Piekarek
Clinical Radiology Department
Poznań University
of Medical Sciences
Długa 1/2
Poznań, Poland
e-mail: alinamed@hotmail.com

ABSTRACT

BACKGROUND: Multiple myeloma is a cancer of plasma cells which leads to bone marrow infiltration.

AIM: Whole-body MR is the most sensitive imaging method available to detect multiple myeloma lesions.

MATERIAL AND METHODS: MR scans were performed in 100 patients with multiple myeloma who were receiving treatment in the Haematology Clinic in Poznań in the years 2005–2006. Whole-body MR scans were performed with general coil 1.0 T in STIR sequences and T1 sequences, in coronal and sagittal planes with scanning area covering the head, neck, trunk and the limbs (FOV for specific regions was 36–48 cm). The bone lesions were classified as focal (monofocal/multifocal lesions), infiltrative, mixed and “salt and pepper” type. Depending on the size of the lesions the patients were included in one of three groups according to Salmon-Durie Plus classification.

RESULTS: Four main types of multiple myeloma were distinguished based on MR scans: focal (48 patients; monofocal in 10 patients), infiltrative (17 patients), mixed type (19 patients) and “salt and pepper” type (4 patients). The remaining 12 patients had no multiple myeloma lesions in the bone marrow. Additionally, in 18% of patients a soft tissue mass could be observed. According to Salmon-Durie Plus categorisation 27 subjects were classified as having stage I, 16 patients stage II and 57 patients stage III disease. In 12% of patients MR data changed the disease staging.

CONCLUSIONS: WB MR is a sensitive and effective diagnostic method with an important impact on staging and further treatment of multiple myeloma.

KEY WORDS: whole-body MR (WB MR); multiple myeloma (MM); Salmon-Durie Plus (S-D Plus)

BACKGROUND

Multiple myeloma (*myeloma multiplex*) is a malignant neoplasm of plasma cells which invade the bone marrow. Multiple myeloma constitutes about 1% of all neoplasms and about 13% of blood neoplasms. The disease may invade the skeleton and the kidneys, may impair immune system function and may lead to haematological abnormalities. Prognosis depends on disease stage, particularly at the time of diagnosis. Treatment involves chemotherapy and bone marrow transplant, as well as radiotherapy.

AIM

Whole-body MR (WB MR) is the most sensi-

tive imaging method available to detect multiple myeloma lesions.

MATERIALS AND METHODS

Whole-body MR scans were performed in 100 subjects with multiple myeloma (39 females and 61 males; age 32 to 79 years; mean age 59.22 years) who were receiving treatment in the Haematology Clinic in Poznań University of Medical Sciences in the years 2005–06. In 62 patients WB MR was performed before the initiation of the treatment, whereas the remaining 38 patients had their WB MR scans performed during multi-phase treatment (after chemotherapy, bone marrow trans-

plant and/or radiotherapy). The time period between WB MR and treatment completion ranged from 3 to 21 days (17 days on average) and it was dependent on the disease stage, overall patient condition and the availability of WB MR.

Whole-body MR scans were performed using a general coil 1.0 T MR apparatus in STIR sequences (TR 2700 ms, TE 50 ms, TI 110 ms) and T1 sequences (TR 450 ms, TE 50 ms), in coronal and sagittal planes, covering the head, neck, trunk and the upper and lower limbs to the knee level (FOV for specific regions ranged from 36 to 48 cm). Additional sequences were performed in 19 patients with sporadic lesions detected in the head, abdomen and the joints. Bone marrow status was graded 0–2, both for focal and for infiltrative lesions, where:

0 = No lesions

1 = Uncertain lesions (indirect signal in STIR and/or T1 sequences)

2 = Certain lesions (significantly high signal in STIR sequences and significantly low signal in T1 sequences)

The number of focal lesions for a specific body region was determined and the lesion area was subjectively categorized as “no infiltration”, “minor infiltration” and “extensive, homogeneous infiltration”. Depending on the extent of the lesions the patients were included in one of three groups according to the Salmon-Durie Plus classification. (Table 1)

RESULTS

Whole-body MR scans allowed four main types of multiple myeloma to be distinguished:

1. Focal type (high signal foci in STIR sequences and low signal foci in T1 sequences) was „observed in 48 patients, including monofocal type – detected in 10 patients. The smallest „foci of <1 cm diameter were found in skull bones, whereas the largest foci of >4 cm diameter were detected in pelvic bones (Fig. 1–2)*

2. Infiltrative type (homogeneous infiltration of high signal in STIR sequences and low signal in T1 sequences) – detected in 17 patients (Fig. 3)

Table 1. Salmon-Durie Plus classification of multiple myeloma lesions

I	0–4 foci and/or absence of infiltration in MR/PET scans
II	5–20 foci and/or minor infiltration in MR/PET scans
III	>20 foci and/or extensive, massive infiltration in MR/PET scans
A: creatinine <2 mg/dl B: creatinine >2 mg/dl	



Fig. 1. Multifocal lesions, Stage III: b – sagittal STIR image: high signal focal lesions in the skull, sternum and vertebral bodies with collapsed Th2, c – sagittal SE T1-W image: low signal lesions in the sternum and vertebral bodies

3. Mixed type (diffuse foci seen against the infiltrative lesion background) – detected in 19 patients

4. “Salt and pepper” type (with high signal fat islets in T1 sequences) – detected in 4 patients

The remaining 12 patients had no multiple myeloma lesions in the bone marrow.

In 18% of our patients a soft tissue mass

* Nother focal lesions were taken into account (for instance angiomas of thoracic and lumbar vertebrae with high signal T1 and STIR (T2-W)).

Download English Version:

<https://daneshyari.com/en/article/1855769>

Download Persian Version:

<https://daneshyari.com/article/1855769>

[Daneshyari.com](https://daneshyari.com)