



Towards a large deviation theory for strongly correlated systems

Guíomar Ruiz^{a,b,*}, Constantino Tsallis^{a,c}

^a Centro Brasileiro de Pesquisas Físicas and National Institute of Science and Technology for Complex Systems, Rua Xavier Sigaud 150, 22290-180 Rio de Janeiro, RJ, Brazil

^b Departamento de Matemática Aplicada y Estadística, Universidad Politécnica de Madrid, Pza. Cardenal Cisneros s.n., 28040 Madrid, Spain

^c Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

ARTICLE INFO

Article history:

Received 12 March 2012

Received in revised form 20 June 2012

Accepted 22 June 2012

Available online 28 June 2012

Communicated by C.R. Doering

Keywords:

Probability theory

Statistical mechanics

Entropy

ABSTRACT

A large-deviation connection of statistical mechanics is provided by N independent binary variables, the $(N \rightarrow \infty)$ limit yielding Gaussian distributions. The probability of $n \neq N/2$ out of N throws is governed by e^{-Nr} , r related to the entropy. Large deviations for a strong correlated model characterized by indices (Q, γ) are studied, the $(N \rightarrow \infty)$ limit yielding Q -Gaussians ($Q \rightarrow 1$ recovers a Gaussian). Its large deviations are governed by $e_q^{-Nr_q}$ ($\propto 1/N^{1/(q-1)}$, $q > 1$), $q = (Q - 1)/(\gamma[3 - Q]) + 1$. This illustration opens the door towards a large-deviation foundation of nonextensive statistical mechanics.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

In his 1902 historical book *Elementary Principles in Statistical Mechanics* [1], Gibbs emphasizes that systems involving long-range interactions are intractable within the Boltzmann–Gibbs (BG) theory, due to the divergence of the partition function. Amazingly enough, this crucial remark is often overlooked in most textbooks.

To solve related complexities it was proposed in 1988 [2–5] a generalization of the BG theory, currently referred to as nonextensive statistical mechanics. It is based on the nonadditive entropy

$$S_q = k_B \frac{1 - \sum_{i=1}^W p_i^q}{q - 1} \quad \left(q \in \mathcal{R}; \sum_{i=1}^W p_i = 1 \right), \quad (1)$$

which recovers $S_{BG} = -k_B \sum_{i=1}^W p_i \ln p_i$ for $q \rightarrow 1$. If A and B are two probabilistically independent systems (i.e., $p_{ij}^{A+B} = p_i^A p_j^B$, $\forall(i, j)$), definition (1) implies the nonadditive relation $\frac{S_q(A+B)}{k_B} = \frac{S_q(A)}{k_B} + \frac{S_q(B)}{k_B} + (1 - q) \frac{S_q(A)}{k_B} \frac{S_q(B)}{k_B}$. Moreover, if probabilities are all equal, we straightforwardly obtain $S_q = k_B \ln_q W$, with $\ln_q z \equiv \frac{z^{1-q} - 1}{1 - q}$ ($\ln_1 z = \ln z$). If we extremize (1) with a constraint on its width (in addition to normalization of the probabilities $\{p_i\}$), we obtain

$$p_i = \frac{e_q^{-\beta_q E_i}}{\sum_{j=1}^W e_q^{-\beta_q E_j}}, \quad (2)$$

e_q^z being the inverse function of the q -logarithmic function, i.e., $e_q^z \equiv [1 + (1 - q)z]^{1/(1-q)}$ ($e_1^z = e^z$); $\{E_i\}$ are the energy levels; β_q is the effective inverse temperature.

Complexity frequently emerges in natural, artificial and social systems. It may be caused by various geometrical–dynamical ingredients, which include nonergodicity, long-term memory, multifractality, and other spatial–temporal long-range correlations between the elements of the system. During the last two decades, many such phenomena have been successfully approached in the frame of nonextensive statistical mechanics. Predictions, verifications and various applications have been performed in high-energy physics [6–11], spin-glasses [12], cold atoms in optical lattices [13], trapped ions [14], anomalous diffusion [15], dusty plasmas [16], solar physics [17,18], relativistic and nonrelativistic nonlinear quantum mechanics [19], among many others.

Large-deviation estimates are important in order to quantify the rates of convergence to equilibrium. They are therefore relevant for addressing the rigorous derivation concerning simplified models to describe complex systems. In fact it is known since several decades that the mathematical foundation of BG statistical mechanics crucially lies on the theory of large deviations (see [20,21] and references therein). To attain the same status for nonextensive statistical mechanics, it is necessary to q -generalize the large deviation theory itself. The purpose of the present effort precisely is to make a first step towards that goal through the study of a simple model.

* Corresponding author at: Departamento de Matemática Aplicada y Estadística, Universidad Politécnica de Madrid, Pza. Cardenal Cisneros s.n., 28040 Madrid, Spain.
E-mail addresses: guiomar.ruiz@upm.es (G. Ruiz), tsallis@cbpf.br (C. Tsallis).

This Letter is organized as follows. In Section 2 we make some considerations about relative entropy associated with binary random variables, and a possible generalization for non-BG entropic random variables. In Section 3 we report some results of large deviation theory in a well-known model assuming independent probabilities and consequently a BG type relative entropy; we extend these results to a modified model that presents complexity due to strong correlations in Section 4. Finally, in Section 5 we summarize our results and conclusions.

2. Relative entropy

The *relative entropy* or *mutual information* for a random variable with discrete W events with probabilities $\{p_i\}$ ($i = 1, \dots, W$) is defined as

$$I_1 = - \sum_{i=1}^W p_i \ln \frac{p_i^{(0)}}{p_i}, \quad (3)$$

where $p_i^{(0)}$ is a reference distribution [22]. By choosing $\{p_i^{(0)}\}$ as the uniform distribution (i.e., $p_i^{(0)} = 1/W$), we have

$$I_1 = \ln W - \sum_{i=1}^W p_i \ln \frac{1}{p_i} = \ln W - \frac{S_1}{k_B}. \quad (4)$$

Definition (3) and the nonadditive entropy S_q naturally lead to the generalization [24]

$$I_q = - \sum_{i=1}^W p_i \ln_q \frac{p_i^{(0)}}{p_i} = \sum_{i=1}^W p_i \frac{[(p_i/p_i^{(0)})^{q-1} - 1]}{q-1}, \quad (5)$$

where by once again choosing as $\{p_i^{(0)}\}$ the equiprobability distribution (i.e., $p_i^{(0)} = 1/W$), we have

$$I_q = W^{q-1} \left[\ln_q W - \frac{S_q}{k_B} \right]. \quad (6)$$

For $W = 2$ (e.g. tossing a coin to have *head* or *tail*), I_1 reads

$$I_1 = \ln 2 + p_1 \ln p_1 + p_2 \ln p_2 \quad (7)$$

and by identifying $(p_1, p_2) \rightarrow (x, 1-x)$, we establish [23]

$$r_1(x) = \ln 2 + x \ln x + (1-x) \ln (1-x), \quad (8)$$

as the inset of Fig. 1 shows. Generalized mutual information (5) now reads

$$I_q(x) = \frac{1}{1-q} [1 - 2^{q-1} [x^q + (1-x)^q]], \quad (9)$$

which recovers expression (8) for $q \rightarrow 1$.

In the following, we will apply these results to the problem of studying large deviations when tossing N times a fair coin or, equivalently, tossing one time N independent fair coins (BG entropic system) and when tossing N strongly correlated fair coins (non-BG entropic system).

3. Large deviations in N uncorrelated coins

Let us now exhibit some large deviation theory results on a standard example which consists in tossing N times a fair coin. This well-known model presents independent probabilities of obtaining n ($n = 0, 1, \dots, N$) heads, that are given by

$$p_{N,n} = \binom{N}{n} \frac{1}{2^N}. \quad (10)$$




Fig. 1. Tossing N independent coins: the large-deviation probability $P(N; n/N < x)$ decays exponentially with N , and the slopes of the logarithms, $\ln P(N; n/N < x)$, provide the rate function numerical values $r_1(x)$, represented in the inset as dots. Continuous curve in the inset corresponds to Eq. (8).

The probability of having the ratio n/N smaller than x with $0 \leq x \leq 1/2$ (the case $1/2 \leq x \leq 1$ is totally symmetric) is given by

$$P(N; n/N < x) = \sum_{\{n | \frac{n}{N} < x\}} p_{N,n}. \quad (11)$$

and it is straightforward to obtain that, in the $N \rightarrow \infty$ limit,

$$P(N; n/N < x) \simeq e^{-Nr_1(x)} \quad (0 \leq x \leq 1/2), \quad (12)$$

where the subindex 1 in the rate function $r_1(x)$ will soon become clear. In fact, this exponential decay with N , deeply related with the exponential decay with energy of the BG weight (namely, the $q = 1$ particular case of Eq. (2)), is verified in Fig. 1. In Section 2, it has been shown the analytical calculation of the rate function $r_1(x)$ for the present trivial model (8). The high-precision numerical calculation can be verified in Fig. 1.

4. Large deviation in N correlated coins

Let us now extend the above model to the case where the coins present strong correlations, so as to introduce complexity through a specific system and study the large-deviation behavior in a simple non-BG system. To deal with this problem, it is appropriate to refer to the recently proposed families of probabilistic models [31, 32] having q -Gaussians as limiting probability distribution. Let us incidentally mention that probabilistic scale invariance is not sufficient for obtaining q -Gaussians. Indeed, in [28] the model is scale invariant but the limit distribution is not q -Gaussian, whereas in [31,32] the models are scale invariant and yield q -Gaussians.

The simple model that we have just reviewed yields, for $N \rightarrow \infty$, a Gaussian distribution. This conforms to the Central Limit Theorem, valid for sums of many independent random variables whose variance is finite. This classical theorem has been q -generalized [27] (see also [28], as well as [29,30]) for a special class of correlations referred to as q -independence (1-independence recovers standard independence). The attractors in the probability space of q -independent models are q -Gaussians, which precisely extremize the (continuous form of the) entropy S_q when an appropriately generalized q -variance is maintained fixed. It is then this class of correlations that we are going to focus on here in order to illustrate how the classical large deviation theory can be generalized. We adopt the specific class of binary variable models introduced in [31]. These models consist in discretized forms of Q -Gaussians

Download English Version:

<https://daneshyari.com/en/article/1867037>

Download Persian Version:

<https://daneshyari.com/article/1867037>

[Daneshyari.com](https://daneshyari.com)