



26th International Symposium on Superconductivity, ISS 2013

Anomalous Meissner Effect in Superconducting Junction with Spin-Active Interface

Takehito Yokoyama^{a*}, Yukio Tanaka^b, and Naoto Nagaosa^{c,d}

^aDepartment of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

^bDepartment of Applied Physics, Nagoya University, Nagoya, 464-8603, Japan

^cDepartment of Applied Physics, University of Tokyo, Tokyo 113-8656, Japan

^dRIKEN Center for Emergent Matter Science (CEMS), Wako 351-0198, Japan

Abstract

We investigate Meissner effect in normal metal/superconductor junctions where the interface is spin-active. We find that the orbital magnetic susceptibility of the normal metal depends on the temperature in an oscillatory fashion, accompanied by its sign change. Correspondingly, magnetic field and current density can spatially oscillate in the normal metal. These results stem from the generation of odd-frequency pairing due to the spin-active interface.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Peer-review under responsibility of the ISS 2013 Program Committee

Keywords: Odd frequency superconductivity; Meissner effect

Interface phenomena related to the superconductivity constitute a rich field of condensed matter physics. When a superconductor is attached to a normal metal, Cooper pairs penetrate into the normal metal which acquires superconducting correlation. This is called the proximity effect. As a result, for example, the normal metal has a gap in the density of states or shows Meissner effect [1, 2, 3]. In most cases, as lowering temperature, the proximity effect and hence the Meissner response become stronger. However, it has been reported that, at very low temperatures, the susceptibility of cylindrical structures shows a reentrant behavior and even has paramagnetic region. [4]

Recently, it has been clarified that in normal metal/superconductor junctions, if the interface is spin-active, induced superconducting pairing in the normal region can change its symmetry from even-frequency pairing to odd-

* Corresponding author. Tel.: +81-3-5734-2695; fax: +81-3-5734-2739
E-mail address: yokoyama@stat.phys.titech.ac.jp

frequency pairing[5, 6], similar to ferromagnet/superconductor junctions[7, 8]. The emergence of odd-frequency pairing is manifested as a zero energy peak in the density of states. [9, 10, 11] Here, even- or odd-frequency means that Cooper pair wavefunction is even or odd with respect to Matsubara frequency (or imaginary time).[12] If proximity induced pairing symmetry changes, the associated Meissner effect will also change qualitatively.

In this paper, we study Meissner response in the normal metal attached to superconductor where the interface is spin-active. We find that orbital magnetic susceptibility of the normal metal shows quite complex dependence on junction parameters. In particular, the magnetic susceptibility depends on the temperature in an oscillatory fashion, accompanied by its sign change. We also show that magnetic field and current density can spatially oscillate in the normal metal. These results are due to the generation of odd-frequency pairing which stems from the spin-active interface.[13]

We consider a junction consisting of a diffusive normal metal (DN) with a length L and resistance R_d , and a superconductor. The interface between the DN and the superconductor at $x = L$ has a resistance R_b (or tunneling conductance G_T) and the surface at $x = 0$ is specular. A weak external magnetic field H is applied in z -direction. We consider spin-active interface at $x = L$ which is described by the mixing conductances which reflect the spin rotation upon reflection and transmission at the interface. [14, 15] To evaluate the mixing conductances, we model magnetic barrier (interface) region as a rectangular potential V with the exchange field h in z -direction and the width d , following Ref.[15].

To study the Meissner response, we adopt the quasiclassical Green's function theory.[16] When a magnetic field is applied parallel to the interface, rich and nontrivial screening effect occurs. Within the linear response theory, the current distribution flowing in y - direction is given by [2]

$$j(x) = -8\pi e^2 N(E_F) DT \sum_{\sigma, \omega_n > 0} \sin^2 \theta_\sigma(x) A(x) \quad (1)$$

where $A(x)$, $N(E_F)$ and T denote the vector potential, the density of states at the Fermi energy and the temperature of the system, respectively. $\sin \theta_\sigma$ is the anomalous Green's functions.

By solving the Maxwell equation, we finally obtain the expression of the orbital magnetic susceptibility, [2]

$$-4\pi\chi = 1 + \frac{A(0)}{HL}$$

We set $h/E_F = 0.01$, $R_d/R_b = 10$ and $16\pi e^2 N(E_F)D^2 = 1000$. Below, ξ and T_c denote the superconducting coherence length and the transition temperature, respectively.

Figure 1 shows susceptibility at $L/\xi = 10$ with $V/E_F = 0.95$ as a function of the temperature T . We find that the susceptibility oscillates with temperature, and, over some region, paramagnetic state, namely that with positive χ , appears. When mixing conductance is present, odd-frequency pairing is generated in the DN region.[5, 6] This odd-frequency pairing makes it possible to oscillate magnetic field rather than suppress in the DN region[17], as explicitly shown below. Therefore, susceptibility could be positive when odd-frequency pairing correlation is dominant over even-frequency pairing in the DN.




Figure 1: Susceptibility $-4\pi\chi$ as a function of the temperature.

Download English Version:

<https://daneshyari.com/en/article/1874580>

Download Persian Version:

<https://daneshyari.com/article/1874580>

[Daneshyari.com](https://daneshyari.com)