



Comparison of maximal elastance and systolic wall thickening using arterial tonometry and gated myocardial SPECT in patients undergoing coronary artery bypass grafting

Won Jun Kang^a, Dong Soo Lee^{b,*}, Byeong Il Lee^c, Jin Chul Paeng^b, Jae Sung Lee^b, June-Key Chung^b, Myung Chul Lee^b

^a Department of Radiology, Division of Nuclear Medicine, Yonsei University College of Medicine, Seoul, Republic of Korea

^b Department of Nuclear Medicine, Seoul National University College of Medicine, 28 Yeongeon-dong, Jongno-gu, Seoul 110-744, Republic of Korea

^c Department of Nuclear Medicine, Chonnam National University Medical School, Gwangju, Republic of Korea

ARTICLE INFO

Keywords:

Myocardial contractility
Maximal elastance
Coronary artery bypass grafting
Gated SPECT

ABSTRACT

Myocardial SPECT using ^{99m}Tc-sestamibi, ^{99m}Tc-tetrofosmin, ²⁰¹Thallium is widely used in nuclear cardiology. Left ventricular systolic wall thickening (SWT) by SPECT and regional maximal elastance (rE_{\max}) using arterial tonometry were compared.

rE_{\max} was calculated from time–pressure and time–volume curves. In normal heart, improvement of SWT was $4.1 \pm 11\%$, while $6.0 \pm 16\%$ in dilated heart. Improvement of rE_{\max} was 0.67 ± 1.0 mmHg/mL in normal heart and 0.32 ± 0.7 mmHg/mL in dilated heart ($p < 0.05$). rE_{\max} can be an alternative variable as an index of regional contractility.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Left ventricular systolic wall thickening (SWT) measured by gated myocardial SPECT is used to evaluate regional left ventricular wall contractility (Germano et al., 1997; Sharir et al., 2001). Automatic quantification of SWT was commonly used, and its reproducibility is superior to echocardiography (Paeng et al., 2001).

However, SWT depends on left ventricular volume, because SWT is measured by percentage increase of myocardial thickness from end-diastole to end-systole. Dilated left ventricle usually has thin diastolic wall thickness, and minor improvement of wall thickness on systolic phase makes thickening score greater. Criteria of abnormal SWT according to end-diastolic wall thickness have not been clarified.

Maximal elastance (E_{\max}) is known to be another marker of myocardial contractility. E_{\max} is a volume-independent marker, and is not influenced by preload or afterload (Suga, 2003a, b; Suga and Paul, 1994). Recently, the method to calculate regional maximal elastance (rE_{\max}) using volume data from gated SPECT and pressure profile obtained from arterial tonometry was developed (Senzaki et al., 1996; Lee et al., 1999; Kim et al., 2001).

In this study, we compared rE_{\max} with systolic thickening in patients with ischemic heart disease, and investigated the

changes of rE_{\max} and systolic thickening after coronary artery bypass grafting (CABG).

2. Materials and methods

2.1. Subjects

Nineteen patients who have ischemic heart disease and underwent CABG were included. There were 15 males and 4 females, ranging in age from 55 to 77 y (60 ± 5 y). Fourteen patients had triple-vessel disease, 4 patients had two-vessel disease, and one had left anterior descending (LAD) artery disease.

2.2. Myocardial SPECT

Rest ²⁰¹Tl-dipyridamole stress ^{99m}Tc-sestamibi gated SPECT was performed before CABG. Patients refrained from consuming caffeine-containing beverages for 24 h and fasted for 4 h before the study. At 10 min after an injection of 111 MBq of ²⁰¹Tl, the rest SPECT image was acquired. Then, dipyridamole of 0.56 mg/kg was continuously injected over 4 min for the stress study. After 3 min of the completion of dipyridamole infusion, 925 MBq of ^{99m}Tc-sestamibi were injected. Gated SPECT with ^{99m}Tc-sestamibi was performed 90 min after the stress study.

SPECT was performed with a dual-head gamma camera equipped with a low-energy high-resolution collimator (Vertex

* Corresponding author. Tel.: +82 2 2072 2501; fax: +82 2 2072 9083.
E-mail address: dsl@plaza.snu.ac.kr (D.S. Lee).

EPIC; ADAC Laboratories). Thirty-two step-and-shoot images were acquired at intervals of 3° for 25 s per step. For gating, 16 frames per cardiac cycle with prefixed R–R intervals and 40% windows were used. Images were reconstructed with a Butterworth filter at a cutoff frequency of 0.35 and order 10 for ^{201}Tl and at a cutoff frequency of 0.45 and order 10 for $^{99\text{m}}\text{Tc}$ -sestamibi. Rest ^{201}Tl -dipyridamole stress $^{99\text{m}}\text{Tc}$ -sestamibi gated SPECT was repeated 3 months after CABG with the same protocol as that used for the preoperative study.

The reconstructed images were analyzed by use of an automatic quantifying software package (AutoQUANT; ADAC Laboratories) without manual intervention. For the regional analysis, a 20-segment model was adopted. Stress and rest segmental perfusion were quantified and expressed as percentages of maximal uptake. Segmental wall thickening was expressed as a percentage of the end-diastolic wall thickness. The left ventricular ejection fraction (LVEF), end-systolic volume (ESV) and end-diastolic volume (EDV) were also measured by quantitative gated SPECT.

2.3. Regional maximal elastance

Radial arterial pressure was measured during gated SPECT with tonometry (SphygmoCor, PWV Medical, Australia). Regional time-elastance curves were obtained using pressure data from radial artery tonometry and volume data from gated SPECT.

A time–pressure curve of the radial artery was acquired and transformed into central time–pressure curve. Assuming the isopressure in left ventricular cavity, time–pressure curve obtained from tonometry was used for every region. The regional elastance curve $E(t)$ was calculated using pressure curve $P(t)$ and volume

curve $V(t)$. We developed Cardiac SPECT Analyzer (CSA) software for automatic calculation of regional E_{max} .

2.4. Statistics

Differences between groups were evaluated using χ^2 test for categorical variables and unpaired t -test for continuous variables. Paired t -test was used to compare pre and post CABG changes of parameters. To assess correlation between variables, correlation coefficient was acquired.

3. Results

EDV and ESV decreased from 123.4 ± 42.6 to 93.6 ± 33.2 ml ($p < 0.01$) and from 66.3 ± 40.2 to 48.1 ± 25.9 ml ($p < 0.01$) after CABG, respectively. Ejection fraction (EF) improved from 50.3 ± 14.2 to $55.4 \pm 15.2\%$ ($p < 0.01$). Global E_{max} improved from 2.2 ± 1.2 to 2.5 ± 1.2 ($p < 0.01$) (Fig. 1).

Global E_{max} showed good correlation with EF in preoperative measurement ($r = 0.8333$, $p < 0.0001$). After CABG, correlation between global E_{max} and EF decreased but there was still statistical significance ($r = 0.6442$, $p = 0.0029$) (Fig. 2). Global E_{max} showed negative correlation with EDV both in preoperative and postoperative measurement ($r = -0.8606$, $p < 0.0001$ in preoperative, $r = -0.7740$, $p = 0.0001$ in postoperative) (Fig. 3).

rE_{max} and SWT showed good correlation between pre CABG and post CABG measurements (Fig. 4, $r = 0.8487$ $p < 0.0001$ in rE_{max} , $r = 0.7300$, $p < 0.0001$ in SWT). However, rE_{max} showed weak correlation with SWT both in pre CABG and post CABG




Fig. 1. Postoperative changes of EF(A), EDV(B) and global E_{max} (C).

Download English Version:

<https://daneshyari.com/en/article/1878173>

Download Persian Version:

<https://daneshyari.com/article/1878173>

[Daneshyari.com](https://daneshyari.com)