



Protein expression and isotopic enrichment based on induction of the Entner–Doudoroff pathway in *Escherichia coli*

Bosmat Refaeli, Amir Goldbourt*

School of Chemistry, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 69978, Tel Aviv, Israel

ARTICLE INFO

Article history:

Received 5 September 2012

Available online 17 September 2012

Keywords:

Entner–Doudoroff pathway

Protein expression

Nuclear magnetic resonance

Isotopic labeling

Sparse labeling

[1-¹³C]-gluconate

ABSTRACT

The Entner–Doudoroff pathway is known to exist in many organisms including bacteria, archaea and eukarya. Although the common route for carbon catabolism in *Escherichia coli* is the Embden–Meyerhof–Parnas pathway, it was shown that gluconate catabolism in *E. coli* occurs via the Entner–Doudoroff pathway. We demonstrate here that by supplying BL21(DE3) competent *E. coli* cells with gluconate in a minimal growth medium, protein expression can be induced. Nuclear magnetic resonance data of over-expressed ubiquitin show that by using [1-¹³C]-gluconate as the only carbon source, and ¹⁵N-enriched ammonium chloride, sparse isotopic enrichment in the form of a spin-pair carbonyl-amide backbone enrichment is obtained. The specific amino acid labeling pattern is analyzed and is shown to be compatible with Entner–Doudoroff metabolism. Isotopic enrichment serves as a key factor in the biophysical characterization of proteins by various methods including nuclear magnetic resonance, mass spectrometry, infrared spectroscopy and more. Therefore, the method presented here can be applied to study proteins by obtaining sparse enrichment schemes that are not based on the regular glycolytic pathway, or to study the Entner–Doudoroff metabolism during protein expression.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

The Entner–Doudoroff (ED) metabolic pathway has been discovered in *Pseudomonas saccharophila* in 1952 [1] and describes the initial steps of carbon catabolism in the cell. It was shown that when *Ps. saccharophila* is fed with glucose, which is radio-labeled at its C1 position, all the ¹⁴C signals are recovered as ¹⁴CO₂ that originates from carboxyl-labeled pyruvate. This result differs from the common Embden–Meyerhof–Parnas (EMP) glycolytic pathway [2], the more common route for the breakdown of carbon, which predicts that C1 is converted to the methyl carbon in pyruvate. It was later discovered that the pathway exists in many other *Pseudomonas* species [3], in other organisms [4] and most importantly in *Escherichia coli* [5]. In particular, when *E. coli* cells are fed with gluconate as their sole carbon source, the ED pathway becomes the dominant route for carbon catabolism [6]. Detailed studies of the ED pathway have been performed in order to study its key enzymes demonstrating that the key factors are the missing phosphofructokinase and active 6-phosphogluconate dehydratase (EDD) and KDPG aldolase (EDA) [7,8].

Isotopic labeling of proteins and other macromolecules serves as a key factor in the biophysical characterization of their structure, function and dynamics. A variety of methods, including nucle-

ar magnetic resonance, mass spectrometry, infrared spectroscopy and more, use isotopic enrichment in order to enhance sensitivity and resolution [9,10]. For example, over-expression [11] in *E. coli* using a minimal medium that is based on enriched isotopes is a common technique for producing proteins for NMR studies, enabling the characterization their structure and dynamics. Multi-dimensional NMR experiments are an essential tool in these studies [12,13] and require the use of NMR-active isotopes such as ²H, ¹³C and ¹⁵N. Sparse isotopic enrichment of proteins also bears many advantages [14–19]; the elimination of the non-enriched carbons from the spectra results in better resolution allowing the study of larger proteins; relayed transfers are minimized enabling more accurate structure determinations; coupling of attached carbons is removed in many cases producing narrower lines in magic-angle spinning (MAS) NMR experiments and easier interpretation of relaxation data in solution NMR [20]. More specifically, the first protein structure determination by MAS NMR was facilitated by the use of 1,3-¹³C and 2-¹³C enriched glycerol media [21], a labeling scheme that was developed earlier for measuring thioredoxin dynamics by solution NMR [22]. Additional examples are the successful quantitative dynamics and binding studies of the 670 kDa 20S proteasome [23] by solution NMR using keto-acid labeling strategies [14] and the combined cryo-EM/solid-state NMR structure elucidation of the type III secretion system, a hollow needle-like protein filament of *E. coli*, using 1-¹³C glucose and 2-¹³C glucose as precursors for carbon [24]. The labeling

* Corresponding author. Fax: +972 3 6409293.

E-mail address: amirgo@post.tau.ac.il (A. Goldbourt).

patterns generated by these precursors, and by other carbon precursors [14,15,19,25,26], are all based on the EMP pathway, which is the main route for the breakdown of carbon in *E. coli*. There is therefore an increasing importance in creating diverse enrichment schemes that will enable the studies of proteins with larger complexity and molecular weight, for unfolded proteins, membrane proteins, and aggregated proteins, all examples of systems that produce highly congested NMR spectra. As opposed to protein overexpression, the production of bacteriophage viruses depends on the specific host that they infect. It was demonstrated that when the filamentous bacteriophage Pf1 is produced in its specific host, *Pseudomonas aeruginosa strain K* (PAK), carbon catabolism follows the Entner–Doudoroff (ED) pathway and not the EMP pathway. Therefore, when $1\text{-}^{13}\text{C}$ glucose (and $^{15}\text{N}_4\text{Cl}$) was used as the sole carbon precursor for Pf1 growth, a $^{15}\text{N}\text{-}^{13}\text{C}$ spin-pair backbone labeling scheme was obtained [27] rather than the enrichment of mainly aliphatic carbons [25,28]. However, up to now, it was not possible to use this pathway in general for every protein of choice, since it is not possible to over express proteins in this pathogenic strain.

Here we demonstrate that the ED pathway can be induced in *E. coli* strains suitable for protein expression, and therefore the sparse labeling schemes that arise are different from those produced following the EMP pathway [2], the common route for glycolysis, and follow the ED pathway. As an example, a spin-pair backbone labeling scheme is shown, in which many but not all carbonyl carbons are enriched to various degrees. Some sidechain methyl and carboxyl groups are fractionally labeled as well.

2. Materials and methods

A pET15-ubiquitin plasmid bearing a His-tag and kanamycin resistance (104 residues) has been transformed into Novagen *E. coli* BL21(DE3) competent cells grown on a typical LB-agar plate using normal procedures [11]. Reference unlabeled samples for testing induction were grown in a minimal medium containing gluconate and ammonium chloride using common protein expression techniques [11]. A $[1\text{-}^{13}\text{C}]$ -gluconate labeled sample was obtained by growing a single colony in a starter LB medium. The preliminary culture was then transferred in a 1:100 ratio to a 1 L of M9 medium with unlabeled gluconate (5 g/L) and NH_4Cl (2 g/L). The unlabeled culture was pelleted at log phase, washed, and redissolved in similar concentrations of $1\text{-}^{13}\text{C}$ gluconate/ $^{15}\text{N}_4\text{Cl}$ enriched precursors within a minimal medium with a 1:4 volume ratio [29]. After 1 h the cells were induced with 0.8 mM IPTG, centrifuged, and lysed using a microfluidizer. Purification was carried out on a Profinity IMAC Ni-charged resin (BIO-RAD) followed by dialysis against a 20 mM sodium-citrate buffer (pH 4.1, 10% D_2O). In all cases, pure protein yields per one liter of unlabeled starting media were 14–18 mg. Solution NMR experiments were performed on a Bruker DRX-500 MHz using a dual probe (1D experiment) and a TXI probe with a Z-gradient (HSQC and HNCQ). For the HSQC experiment (acquired in 65 min.), 235/2048 points were acquired in t_1/t_2 for a total of 58/128 ms acquisition times. Data were linear predicted in the indirect dimension, apodized with a squared cosine and zero filled to 512 t_1 points. For HNCQ (17 h), 96/40/2048 points were acquired (17.5/11/128 ms) and processed with linear prediction in both indirect dimensions followed by cosine square apodization and zero filling to 512/256/2048 points. In both experiments 16 scans were acquired with a repetition rate of 1 s. The relative peak intensities, and therefore the relative enrichments (with respect to Val), were determined by taking both volume and peak height ratios (determined from Sparky [30]) of the HSQC and HNCQ spectra, and then averaging over peaks with preceding carbonyls belonging to similar amino acids. A few peaks

showed large deviations from the average, and were therefore dropped. $[1\text{-}^{13}\text{C}]$ -gluconate precursor was obtained from Cambridge isotope laboratories with an enrichment level of 99.4%.

3. Results and discussion

The principal step in the ED pathway (Fig. 1) is the dehydrogenation of glucose-6 phosphate (G6P) to 6-phosphogluconate (6PG), replacing the conversion to fructose-1,6-bisphosphate (FBP) that normally occurs in the EMP pathway. This step results from the lack of the enzyme phosphofructokinase (*pfk*) in organisms in which the ED pathway is naturally occurring. In the EMP pathway, FBP is converted to pyruvate and glyceraldehyde 3-phosphate (G3P/GAP). However in the ED pathway, 6PG is converted to 2-keto-3-deoxy-6-phosphogluconate (KDPG) in a reaction catalyzed by 6-phosphogluconate dehydratase (EDD) and KDPG is converted to pyruvate and to G3P in a reaction catalyzed by KDPG aldolase (EDA). In both pathways, G3P goes through a series of reactions that generate precursors for the synthesis of serine, glycine and cysteine (from 3PG, 3-phosphoglycerate) as well as phosphoenolpyruvate (PEP) that enters the pentose phosphate pathway (PPP) resulting in biosynthesis of aromatic amino acids and the TCA cycle generating oxaloacetate [2,31]. Although the final products of the two pathways are the same, i.e. pyruvate and G3P, the fate of the carbons of the precursor are different; while in EMP C-1 of glucose (marked ^{13}C in Fig. 1) occupies the methyl position in pyruvate, in the ED pathway it occupies the carbonyl carbon. The ED pathway is the preferred catabolic route in many *Pseudomonas* bacteria however, when *E. coli* is nurtured with gluconate, the formation of gluconokinase [6] is induced (and two additional enzymes of the ED pathway, EDD and EDA), leading to the production of 6PG, and therefore *E. coli* is forced to follow the ED pathway.

We have shown in the past that for the Pf1 filamentous bacteriophage [27], the host of which is *Ps. aeruginosa*, the naturally occurring ED pathway leads to carbonyl enrichment if the bacterial culture is fed with $[1\text{-}^{13}\text{C}]$ -glucose. Consequently, line narrowing and spectral simplification were obtained in a multi-dimensional 3D NCC magic-angle spinning solid-state NMR experiment. The resulting spectra were used for the identification of the chemical shifts of the isotopically labeled carbonyl carbons belonging to the capsid of the intact virus. Furthermore, Kay et al. have shown [32,33] that by feeding *E. coli* directly with $[1\text{-}^{13}\text{C}]$ -pyruvate, a carbonyl enrichment patterns is also obtained and can be used for measuring invisible and excited protein states using solution NMR.

In order to verify whether protein expression in a minimal medium containing gluconate follows the ED pathway, we transformed competent *E. coli* cells with a plasmid coding for ubiquitin. Cells were grown initially on unlabeled gluconate and ammonium chloride and the presence of over-expressed ubiquitin was then verified by SDS-PAGE (Supporting Fig. S1). Once expression was verified, a fresh cell culture was grown to log phase using the same unlabeled precursors and pelleted at log phase. These cells were then washed and resuspended in a minimal M9 medium containing $[1\text{-}^{13}\text{C}]$ -gluconate and $^{15}\text{N}_4\text{Cl}$ as the sole carbon and nitrogen sources, respectively, induced after 1 h and purified. We then performed NMR experiments in solution on the purified ubiquitin sample. The spectrum in Fig. 2, resulting from a ^{13}C single pulse experiment, shows that indeed the dominant signal comes from the carbonyl carbons, that the backbone $\text{C}\alpha$ carbons are completely suppressed and that sidechains of distinct amino acids have also been labeled. Most of these aliphatic resonances have already been associated with the existence of labeled carbonate ions in the growth medium [27]; they result from the release of $^{13}\text{CO}_2$ during several processes such as the entry of pyruvate into the TCA cycle, the conversion of α -ketoisovalerate to leucine and the entry of 6PG

Download English Version:

<https://daneshyari.com/en/article/1929492>

Download Persian Version:

<https://daneshyari.com/article/1929492>

[Daneshyari.com](https://daneshyari.com)