



Intracellular trafficking of superparamagnetic iron oxide nanoparticles conjugated with TAT peptide: 3-dimensional electron tomography analysis

Baiju G. Nair, Takahiro Fukuda, Toru Mizuki, Tatsuro Hanajiri, Toru Maekawa*

Bio-Nano Electronics Research Centre, Toyo University, Saitama 350-8585, Japan

ARTICLE INFO

Article history:

Received 28 March 2012

Available online 21 April 2012

Keywords:

Nanoparticles

Internalisation

Cell penetrating peptides

Endocytic pathways

Transmission electron microscopy

3-D tomography

ABSTRACT

Internalisation of nanoparticles conjugated with cell penetrating peptides is a promising approach to various drug delivery applications. Cell penetrating peptides such as transactivating transcriptional activator (TAT) peptides derived from HIV-1 proteins are effective intracellular delivery vectors for a wide range of nanoparticles and pharmaceutical agents thanks to their amicable ability to enter cells and minimum cytotoxicity. Although different mechanisms of intracellular uptake and localisation have been proposed for TAT conjugated nanoparticles, it is necessary to visualise the particles on a 3-D plane in order to investigate the actual intracellular uptake and localisation. Here, we study the intracellular localisation and trafficking of TAT peptide conjugated superparamagnetic iron oxide nanoparticles (TAT-SPIOs) using 3-D electron tomography. 3-D tomograms clearly show the location of TAT-SPIOs in a cell and their slow release from the endocytic vesicles into the cytoplasm. The present methodology may well be utilised for further investigations of the behaviours of nanoparticles in cells and eventually for the development of nano drug delivery systems.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Nanoparticles internalised in cells can be utilised in various biomedical areas such as hyperthermia, drug release, imaging and gene silencing [1–3]. To make the above operations successful, nanoparticles need to enter the cells and stay inside for a period of time, which in particular projects an important aspect of nano-drug delivery systems [4]. It is well known that nanoparticles can enter a cell via different methods; i.e., non specific uptake by endocytosis, direct injection of nanomaterials, electroporation and specifically targeted uptake of nanomaterials functionalised with various targeting moieties. Among the above methods, nanoparticles conjugated with specific ligands are highly promising for selective nanodrug delivery systems [5]. Cell penetrating peptide (CPP) is supposed to be one of the best ligands to improve active internalisation of nanoparticles into a target cell [6]. CPPs in general translocate efficiently across cell membranes, but the exact mechanism of the cell penetrating activity is still under investigation [7]. CPPs such as transactivating transcriptional activator (TAT) peptide derived from HIV can transport a large number of nanoparticles into a mammalian cell cytoplasm avoiding the normal endocytic pathways through the penetration of the cell membrane [8]. Complete lack of specificity of TAT peptides used to be a solemn matter of concern for targeting a cancer cell, but a selective

internalisation of nanoparticles into a cancer cell was successfully performed by combining TAT peptides with cell targeting peptides (CTPs) [9]. This kind of combined peptides can be utilised for the internalisation of an enormous amount of nanoparticles into cells and are also useful for various intracellular manipulation studies [10]. Internalisation of a massive number of nanoparticles into a target cell and their escape from the normal endocytic pathways are attractive features particularly for nano drug delivery systems [11]. Most of the studies on the internalisation of nanoparticles are based on the confocal and flow cytometry analyses. However, a complete visualisation of intracellular trafficking of nanoparticles using the above optical methods is quite difficult due to the high electrostatic interaction between nanoparticles and the surface of cells, which often leads to wrong interpretations of the data [12]. Transmission electron microscopic (TEM) studies support the identification and characterisation of nanoparticles internalised in a cell [13–16]. In the case of 2-D TEM images, however, it is extremely difficult to differentiate whether the nanoparticles are present in the resin or lying on the surface of cell sections without being internalised, or whether the nanoparticles are distributed in the 3-D volume of the section or not. 3-D electron tomography (ET) is a well-established technique in biological sciences. In 3-D ET, the specimen is rotated around an axis at a range of tilt angles perpendicular to the electron beam and multiple 2-D projection images of a 3-D sample are recorded. Finally, high resolution 3-D images are constructed from a set of 2-D images [17]. The 3-D structures of organelles and macromolecular assemblies [18] and the molecular organisation of the cytoplasm in thin sections of

* Corresponding author.

E-mail address: maekawa@toyo.jp (T. Maekawa).




Fig. 1. TEM images of superparamagnetic iron oxide nanoparticles (SPIONs) in a glioma cell. (A) Glioma cell (Control); (B) Clusters composed of HIV-1 TAT peptide conjugated SPIONs (TAT-SPIONs); (C) Bare SPIONs in a glioma cell; (D) TAT-SPIONs in a glioma cell. A large amount of TAT-SPIONs were internalised in vesicles of the glioma cell. Although bare SPIONs were also internalised in the cell, the quantity was much less compared to the TAT-SPIONs.

cells [19] were visualised by 3-D ET. The structure of a virus based on their outer layer glycoprotein spikes, interior nuclear caspid and tegument was also visualised [20]. 3-D ET also augmented the study on pathways exhibited by viruses at different stages of infection focusing on the nano ecology of the protein spikes present on the viral envelop [21]. For instance, the interaction between SIV/HIV-1 viruses and the surface of cells and the entry of these viruses into cells were clearly visualised using 3-D ET [22]. Moreover, the life cycle of a virus in a host cell and related intracellular pathways including endocytic pathways and nuclear entry at different stages of viral infection were successfully visualised by 3-D ET [23]. Interestingly, each ligand conjugated nanoparticle can be viewed as a miniature form of a virus [24,25] and the successful visualisation of the intracellular activities of viruses in a host cell by 3-D ET can be utilised as an inspiring tool for the visualisation of nanoparticles interacting with cells. Intracellular uptake and related localisation of nanomaterials such as carbon nanotubes and fullerenes without any modification with ligands were visualised by 3-D ET [26,27]. In the present study, we analysed the intracellular localisation and trafficking of HIV-1 TAT peptide conjugated superparamagnetic iron oxide nanoparticles (TAT-SPIONs) in a glioma cell by 3-D ET. We suppose that the present study gives more insight into the interactions between nanoparticles and cells and may encourage further investigations of the effect of the type of ligands and the size and material of nanoparticles on the intracellular trafficking of nanoparticles using 3-D ET.

2. Materials and methods

2.1. Preparation of TAT-SPIONs

Commercially available superparamagnetic iron oxide nanoparticles (nanomag[®]-D – spio PEG COOH, Micromod, Germany) of 50 nm average diameter coated with PEG were used for the experimental studies. An 11-mer HIV-1 TAT peptide (YGRKKRRQRRR) (Anaspec, USA) was conjugated with SPIONs using the carbodiimide coupling chemistry [28]. 50 μ l of SPIONs were treated with 250 mM 1-ethyl-3-[3-dimethylaminopropyl] carbodiimide hydrochloride (EDC) and 100 mM *N*-hydroxysulfosuccinimide (S-NHS) (Pierce, Thermo Scientific, USA) in MES buffer of pH 5.5 (BupH MES, Thermo Scientific, USA) and incubated for 15 min. Then,

1 mg of peptide in 100 μ l of 50 mM sodium tetraborate (Wako Pure Chemical, Japan) containing 100 mM sodium chloride was added to the carboxyl group activated SPIONs and the mixture was kept at 20 °C for 4 h. After the conjugation, free peptides were separated by centrifugal filters (Nanosep, 100K, Pall Corporation, USA). Peptide conjugated nanoparticles were washed three times with MES buffer and were resuspended in fresh PBS for cell interaction studies.

2.2. Cell internalisation protocols

Human glioblastoma (U251, JCRB, Japan) were grown on a tissue culture flask with 10% heat inactivated fetal bovine serum and 1% antibiotics in MEM (Sigma–Aldrich, USA) and after three passages, cells were maintained at 1×10^5 cells ml^{-1} . For internalisation studies, cells were grown on 60 mm culture plates to reach confluence of approximately 60%. Next, cells were washed with Dulbecco's PBS (Sigma–Aldrich, USA) and incubated with 25 μ l of TAT peptide conjugated SPIONs or bare SPIONs in the culture medium for 4 h.

2.3. 3-D electron tomography studies

Glioma cells incubated with either TAT-SPIONs or bare SPIONs were washed twice with PBS to remove any residual particles on the cells. Samples for TEM observations were prepared according to the protocol mentioned elsewhere [29]. After cutting the resin embedded samples in 70–80 nm thickness with an ultramicrotome using a 45° wedge angle diamond knife (Lieca Ultracut, Germany), the sections were put on a 300 mesh copper grid and observed by a TEM (JEM 2100, JEOL, Japan) at 120–200 kV. 3-D ET was conducted on the same cell sections as those observed in 2-D electron microscopy. The cell specimen was tilted at an angle from –60 to +60 at 1° increment for better resolution and electron micrographs were taken by a slow scan CCD camera. 3-D reconstruction of each 2-D image was carried out by composer and visualiser application software (TEMography™, System in Frontier Inc., Japan). Various quantitative structural analyses of cells were carried out by zooming, rotating, panning and tilting the 3-D images to understand the diverse behaviours of the molecular assemblies in a cell [30]. Finally, slicing of 3-D images on XZ and YZ planes was carried

Download English Version:

<https://daneshyari.com/en/article/1929769>

Download Persian Version:

<https://daneshyari.com/article/1929769>

[Daneshyari.com](https://daneshyari.com)