



## Identification of the amino acid sequence motif of $\alpha$ -synuclein responsible for macrophage activation

Saet-byul Lee<sup>a</sup>, Sang Myun Park<sup>b</sup>, Keun Jae Ahn<sup>a</sup>, Kwang Chul Chung<sup>c</sup>, Seung R. Paik<sup>d</sup>, Jongsun Kim<sup>a,\*</sup>

<sup>a</sup>Department of Microbiology and Brain Korea 21 Projects For Medical Science, Institute for Immunology and Immunological Diseases, Yonsei University College of Medicine, 134 Shinchon-dong, Seodaemun-gu, Seoul 120-752, Republic of Korea

<sup>b</sup>Department of Pharmacology, Ajou University School of Medicine, Suwon, Republic of Korea

<sup>c</sup>Department of Biology, College of Sciences, Yonsei University, Seoul, Republic of Korea

<sup>d</sup>School of Chemical and Biological Engineering, College of Engineering, Seoul National University, Seoul, Republic of Korea

### ARTICLE INFO

#### Article history:

Received 19 January 2009

Available online 8 February 2009

#### Keywords:

Synuclein

TNF- $\alpha$

Macrophage

Inflammation

Parkinson's disease

### ABSTRACT

$\alpha$ -Synuclein (Syn) is implicated in the pathogenesis of PD and related neurodegenerative disorders. Recent studies have also shown that  $\alpha$ -synuclein can activate microglia and enhance dopaminergic neurodegeneration. The mechanisms of microglia activation by  $\alpha$ -synuclein, however, are not well understood. In this study, we found that not only  $\alpha$ -synuclein but also  $\beta$ - and  $\gamma$ -synucleins activated macrophages (RAW 264.7) *in vitro*. Macrophages treated with synuclein proteins secreted TNF- $\alpha$  in a dose-dependent manner. Synuclein family proteins also increased mRNA transcription of COX-2 and iNOS. Two  $\alpha$ -synuclein deletion mutants, Syn $\Delta$ NAC and Syn61–140, activated macrophages, while deletion mutants Syn1–60 and Syn96–140 did not significantly activate them. Finally, we demonstrated that macrophage activation by  $\alpha$ -synuclein was accompanied by phosphorylation of ERK. These results suggest that synuclein family proteins can activate macrophages, and that macrophage activation needs both the N-terminal and C-terminal domains of  $\alpha$ -synuclein, but not the central NAC region.

© 2009 Elsevier Inc. All rights reserved.

### Introduction

$\alpha$ -Synuclein, an acidic, heat-resistant and unstructured protein of 140 amino acids long, is highly expressed in brain tissues and primarily localized at the presynaptic terminals of neurons [1,2].  $\alpha$ -Synuclein is widely expressed in the central nervous system (CNS). In addition,  $\alpha$ -synuclein is expressed in various tissues [2].  $\alpha$ -Synuclein consists of three distinct regions. The N-terminal region (1–60 amino acid residues) contains KTEGV repeats, which form amphipathic  $\alpha$ -helices similar to the lipid binding domain of apolipoproteins. The central region (61–95 amino acids residues) is a very hydrophobic NAC (non A $\beta$ -component of Alzheimer's disease) peptide, and the C-terminal region (96–140 amino acid residues) is primarily composed of acidic amino acids [2,3]. In addition to  $\alpha$ -synuclein,  $\beta$ -,  $\gamma$ -synucleins and synoretin, which belong to the synuclein family, have been identified in humans [4–7].

$\alpha$ -Synuclein has traditionally been considered a cytoplasmic protein [1]. This view was challenged recently by the finding that  $\alpha$ -synuclein has been detected in extracellular biological fluids, including human cerebrospinal fluid (CSF) and blood plasma in both healthy subjects and patients with Parkinson's disease [8–10]. Recent studies have also shown that  $\gamma$ -synuclein, as well

as  $\alpha$ -synuclein, is present in CSF. In particular,  $\alpha$ - and  $\gamma$ -synucleins are more abundant in aged subjects with neurodegenerative and vascular changes [11]. Moreover, a recent report demonstrated that  $\alpha$ -synuclein is rapidly secreted from cells via unconventional, ER/Golgi independent exocytosis [12], suggesting that extracellular  $\alpha$ -synuclein may function physiologically or pathologically. In agreement with these suggestions, it has been recently reported that extracellular  $\alpha$ -synuclein activates microglia, THP-1 cells and astrocytes [13–16]. Extracellular  $\alpha$ -synuclein also induces microglial phagocytosis [17].

In the present study, we investigated the molecular mechanisms by which  $\alpha$ -synuclein activates macrophages. To identify the amino acid sequence motif of  $\alpha$ -synuclein responsible for macrophage activation, we used synuclein family proteins and several types of  $\alpha$ -synuclein deletion mutants. We compared their influence on tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) secretion, and looked at Cytochrome c oxidase subunit 2 (COX-2) and inducible Nitric oxide synthase (iNOS) expression as macrophage activation markers. We also investigated the downstream signaling of  $\alpha$ -synuclein-induced macrophage activation.

### Materials and methods

**Materials.** Fetal bovine serum (FBS) and Dulbecco's modified Eagle medium (DMEM) were obtained from Gibco-Invitrogen

\* Corresponding author. Fax: +82 2 392 7088.

E-mail address: [jkim63@yuhs.ac](mailto:jkim63@yuhs.ac) (J. Kim).

(Carlsbad, CA). Lipopolysaccharide (LPS) and Polymyxin B sulfates were purchased from Sigma (St. Louis, MO). ECL solution and BCA protein assay kits were purchased from PIERCE (Rockford, IL). Antibodies to P44/42 mitogen activated protein kinase (MAPK), phospho-P44/42 MAPK, Stress-activated protein kinase/Jun amino-terminal kinase (SAPK/JNK), phospho-SAPK/JNK, p38 MAPK and phospho-p38 MAPK were obtained from Cell Signalling Technology (Danvers, MA). Antibody against  $\alpha$ -tubulin was obtained from Sigma (St. Louis, MO). The ELISA TNF kit was from BD science (555212 and 558874, Franklin Lakes, NJ). All other reagents used in this study were analytical grade and obtained from either Sigma (St. Louis, MO) or USB (Cleveland, OH).

**Protein expression and purification.** Synuclein proteins were overexpressed in *E. coli* (BI21), and the recombinant proteins were purified as described previously [18]. Briefly, the transformed bacteria were grown in LB medium with 0.1 mg/ml ampicillin at 37 °C to an A600 of 0.8, and then cultured for an additional 4 h after being induced with 0.5 mM IPTG. The cells were harvested by centrifugation at 8000g for 10 min, resuspended in 20 mM MES pH 6.0, and then disrupted by ultrasonication. The supernatant was purified with DEAE anion-exchange chromatography, followed by CM cation-exchange chromatography in 20 mM MES, pH6.0. The bound proteins were eluted with a linear gradient between 0.1 M and 0.5 M NaCl at a flow rate of 1.5 mL/min. All proteins were further purified on an FPLC gel-filtration column (GE healthcare, Sweden) pre-equilibrated with PBS, pH7.4. All proteins were concentrated and buffer changed with a Centricon apparatus (Satorius Stedim Biotech, Germany). Proteins were quantitated with the BCA assay, then filtered and stored at 4 °C until use.

**Cell culture.** RAW 264.7, a mouse macrophage cell line obtained from the Korean Type Culture Collection, was grown in DMEM with 10% FBS and maintained at 37 °C in a humidified atmosphere of 5% CO<sub>2</sub> and 95% air.

**ELISA.** RAW 264.7 cells ( $1 \times 10^5$  per well) were cultured in 24 well plates overnight. The medium was then removed and replaced with fresh DMEM containing 10% FBS. Cells were incubated with the indicated doses of LPS or indicated doses of recombinant protein with 10  $\mu$ g/ml polymyxin B sulfate for 18 h. The cell-free supernatants were then harvested, and the TNF- $\alpha$  levels in the supernatants were determined using ELISA kits according to each manufacturer's instructions.

**Activation of human primary macrophages.** Peripheral blood mononuclear cells (PBMC) from healthy volunteers were obtained through passage over Ficoll-hypaque gradient (GE Healthcare, Sweden). The cells were then washed and resuspended in RPMI 1640 medium with 10% FBS at a concentration of  $4 \times 10^6$  cells/ml. These cells were placed on 24 well plates, at 1ml/well, and incubated for 2 h at 37 °C in 5% CO<sub>2</sub>. Nonadherent cells were removed from the plates, and adherent cells were cultured in RPMI 1640 medium with 10% FBS for 4 days. Cells were incubated with the indicated doses of LPS or 5  $\mu$ M of recombinant synuclein proteins with 10  $\mu$ g/ml polymyxin B sulfate for 24 h. The cell-free supernatants were then harvested, and the TNF- $\alpha$  levels in the supernatants were determined using ELISA kits according to each manufacturer's instructions.

**RT-PCR.** RAW 264.7 cells ( $2 \times 10^6$ ) were incubated with the indicated doses of proteins for 6 h and then harvested. Total RNA was isolated using an RNeasy mini kit (Qiagen, Santa Clara, CA). To synthesize cDNA, 1  $\mu$ g of each RNA sample was mixed with 100 ng of random hexamers, 6  $\mu$ l of 5 $\times$  first strand buffer, 12  $\mu$ l of 2.5 mM dNTPs (TaKaRa, Shiga, Japan) and 200 units of murine Molony leukemia virus reverse transcriptase (Invitrogen) and incubated at 42 °C for 80 min. The reaction mixture was boiled at 95 °C for 5 min. cDNA was amplified by PCR using PCR PreMix (Bioneer, Seoul, Korea) and a pair of primers specific for the genes of interest. PCR was performed using the following specific oligonucleotide

primer sets: COX-2 forward, 5'-TTC TTC AAC CTC TCC TAC TAC-3', and reverse, 5'-GCA CGT AGT CTT CGA TCA CTA-3'; iNOS forward, 5'-ATG TCC GAA GCA AAC ATC ACA-3', and reverse, 5'-TAA TGT CCA GGA AGT AGG TGA-3'; TNF- $\alpha$  forward, 5'-CTA CTG AAC TTC GGG GTG ATC-3', and reverse, 5'-CAG TCG GCT AAA CGA TAG AGT-3'; GAPDH forward, 5'-GAT CAT CAG CAA TGC CTC CTC-3', and reverse, 5'-TGT GGT CAT GAG TCC TTC CA-3'.

**Western blot analysis.** RAW 264.7 cells ( $5 \times 10^6$ ) were incubated with the indicated doses of proteins for the indicated times and then harvested. The cells were lysed with lysis buffer (10 mM Tris-HCl, pH7.4, 150 mM NaCl, 2 mM EDTA, 1% Triton X-100, 1 mM PMSF, 15  $\mu$ g/ml leupeptin, 2 mM NaF and 2 mM NaVO<sub>4</sub>). The protein bands were transferred onto Polyvinylidene Fluoride (PVDF) membranes (Pall Corporation, Ann Arbor, MI, USA). The membranes were blocked with blocking buffer [Tris-buffered saline (TBS) containing 5% nonfat dried milk] for 2 h and incubated with the indicated primary antibodies for 1 h. After washing three times with TBS containing 0.1% Tween 20 (TBS-T), the membranes were incubated with secondary antibodies for 1 h. After washing three times with TBS-T, the membranes were developed using an ECL kit and then exposed to FUJI X-ray film (Tokyo, Japan).

## Results

### *Synuclein family proteins can induce TNF- $\alpha$ secretion in RAW 264.7 cells*

To elucidate whether synuclein proteins activate macrophages, RAW 264.7 murine macrophage cells were first incubated with  $\alpha$ -synuclein, and then the secretion of TNF- $\alpha$  in the cells was analyzed by ELISA. We used polymyxin B to rule out the effect of contaminating endotoxin in the solution of recombinant proteins. When cells were incubated with 10  $\mu$ g/ml polymyxin B, 10 ng/ml LPS-induced TNF- $\alpha$  secretion was completely inhibited (Fig. 1A), indicating that polymyxin B treatment could inhibit the endotoxin contamination in recombinant proteins. Under these conditions,  $\alpha$ -synuclein induced TNF- $\alpha$  secretion in RAW 264.7 cells in a dose-dependent manner (Fig. 1A). To rule out the effect of cell death by  $\alpha$ -synuclein, we also performed an MTT assay, and observed that  $\alpha$ -synuclein did not induce cell death of RAW 264.7 cells under the same conditions (data not shown).

We next used  $\beta$ - and  $\gamma$ -synucleins to clarify whether the effect of  $\alpha$ -synuclein on TNF- $\alpha$  secretion in RAW 264.7 cells was a general property of synuclein family proteins. As shown in Fig. 1,  $\beta$ - and  $\gamma$ -synucleins, like  $\alpha$ -synuclein, induced TNF- $\alpha$  secretion in RAW 264.7 cells in a dose-dependent manner (Fig. 1B and C). All synuclein proteins appeared to induce similar amounts of TNF- $\alpha$  secretion in RAW 264.7 cells. We also observed that  $\alpha$ -synuclein enhanced TNF- $\alpha$  secretion in BV-2 microglia cells (data not shown). Furthermore, when cells were incubated with  $\alpha$ -,  $\beta$ - and  $\gamma$ -synucleins, all three synuclein proteins induced the expression of TNF- $\alpha$ , COX-2 and iNOS mRNAs (Fig. 1D).

### *Effects of $\alpha$ -synuclein point mutants A30P, E46K and A53T on the induction of TNF- $\alpha$ secretion in RAW 264.7 cells*

The three point mutants of  $\alpha$ -synuclein (A30P, E46K and A53T) that are associated with a few cases of familial Parkinson's disease [19–21] have been thoroughly studied to determine the role of  $\alpha$ -synuclein in the pathogenesis of Parkinson's disease. These mutant forms of  $\alpha$ -synuclein appear to have different properties from the wild-type  $\alpha$ -synuclein with respect to aggregation patterns, binding to lipid membranes and toxicity to cells [22–24]. We investigated whether these  $\alpha$ -synuclein point mutants might function differently in the activation of macrophages. When RAW 264.7 cells were incubated with  $\alpha$ -synuclein point mutants, each mutant

Download English Version:

<https://daneshyari.com/en/article/1933819>

Download Persian Version:

<https://daneshyari.com/article/1933819>

[Daneshyari.com](https://daneshyari.com)