

Available online at www.sciencedirect.com



BBRC

Biochemical and Biophysical Research Communications 352 (2007) 153-157

www.elsevier.com/locate/ybbrc

Progressive mRNA decay establishes an *mkp3* expression gradient in the chick limb bud

Susana Pascoal, Raquel P. Andrade, Fernanda Bajanca, Isabel Palmeirim *

Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, 4710-057 Braga, Portugal

Received 30 October 2006 Available online 9 November 2006

Abstract

The apical ectodermal ridge (AER) controls limb outgrowth and patterning, such that its removal causes changes in mesodermal gene expression, cell death and limb truncation. Fibroblast growth factor (FGF) family members are expressed in the AER and can rescue limb bud outgrowth after AER removal. Cells localized underneath the AER are maintained in an undifferentiated state by the FGFs produced by the AER. *MAPK phosphatase 3 (mkp3)* is a downstream effector of FGF8 signalling during limb bud development and is expressed in the distal limb mesenchyme. The present work evidences a gradient of *mkp3* transcripts along the chick limb bud, in a distal to proximal direction. *mkp3* transcription occurs only in the most distal limb bud cells and its mRNA gradient throughout the limb results from progressive mRNA decay. We show that FGF8-soaked beads induce ectopic *mkp3* expression, indicating that AER-derived FGF8 protein may activate *mkp3* in the distal mesenchyme.

© 2006 Elsevier Inc. All rights reserved.

Keywords: FGF8; mkp3; mRNA decay; mRNA gradient; Limb bud distal mesenchyme; Chick embryo development

Initiation of limb bud development occurs at defined positions in the flank of the embryo through differential cell proliferation of mesenchymal cells of two embryonic origins, lateral plate and somitic mesoderm. Fibroblast growth factors (FGFs) are essential during embryonic development namely, for the regulation of cellular proliferation and differentiation [1].

Intracellular responses to FGFs are mediated by several signal transduction cascades, including the MAPK and the phosphatidylinositol-3-OH kinase (PI(3)K)/Akt pathways [2]. The *MAPK phosphatase 3 (mkp3)* encodes a dual-specificity phosphatase, known to be a negative regulator of the MAPK/ERK pathway [3–6]. Kawakami and colleagues (2003) found that *mkp3* is a downstream effector of the FGF8 signalling pathway during limb development, and that its induction is mediated through the PI(3)K pathway

[7]. Thus, *mkp3* mediates the cellular response to FGF8 signalling in the vertebrate limb [7].

The growing limb mesenchyme is surrounded by ectoderm, whose distal tip forms a specialized epithelial structure, the apical ectodermal ridge (AER) [8]. AER removal leads to limb truncation, evidencing that the AER is responsible for proximal-distal (p–d) limb outgrowth and its activity keeps the subjacent mesoderm in an undifferentiated state [8,9]. Several members of the FGF family, fgf2, fgf4, and fgf8 are expressed in the chick AER and rescue limb outgrowth and patterning following ridge removal [10–12].

FGFs are diffusible proteins whose signals are mediated by a group of four transmembrane proteins with intrinsic tyrosine kinase activity, known as FGF receptors (FGFRs) [13,14]. FGFRs are expressed in a variety of tissues and organs throughout vertebrate development with a distinct expression pattern for each receptor. FGFR1, 2, and 3 exist in two alternative splice variants: IIIb and IIIc. FGF4 and FGF8 are produced by the limb epithelium AER

^{*} Corresponding author. Fax: +351 253 604831.

E-mail address: ipalmeirim@ecsaude.uminho.pt (I. Palmeirim).

⁰⁰⁰⁶⁻²⁹¹X/\$ - see front matter @ 2006 Elsevier Inc. All rights reserved. doi:10.1016/j.bbrc.2006.10.186

and preferentially bind to the IIIc isoforms [15], which are expressed in the mesenchyme [16-18].

In this work, we show that FGF8 protein produced in the AER induces the expression of its downstream effector mkp3 in the distal limb mesenchyme. mkp3 presents an expression gradient in the limb mesenchyme and we show that this gradient results from the decay of mRNA transcribed by the distal limb bud progenitors.

Materials and methods

Eggs and embryos. Fertilised chick (*Gallus gallus*) eggs obtained from commercial sources were incubated at $37.2 \text{ }^{\circ}\text{C}$ in a 49% humidified atmosphere and staged according to the Hamburger and Hamilton (HH) classification [19].

RNA probes. Antisense digoxigenin-labelled RNA probes *mkp3* [7] was prepared as previously described.

An intronic *mkp3 in situ* hybridisation probe was generated. Reverse transcription and polymerase chain reactions (RT-PCR) were used to isolate the fourth chick *mkp3* intron, using the sense oligo 5'-AGCA GAACCCCCATTCCCTTCT-3' and the antisense oligo 5'-CCAG CATGGAAATCCTGCA-3'. The DNA fragment generated was cloned into the pCR[®]II-TOPO[®] vector (Invitrogen) and plasmid DNA was isolated. Antisense *mkp3* intronic RNA probe was produced using standard procedures.

RNA extraction and reverse transcription. Limb buds at stage HH24 were dissected into three portions of equivalent size along the p-d axis. Total mRNA was extracted from 5 distinct batches of each portion of the limb using the RNeasy Mini Kit Protect (Qiagen, Germany). Total mRNA quantification was done by spectrophotometry (NanoDrop Technologies, Inc., USA). Total RNA was digested with DNAse RNAse-Free (Promega, USA) according to manufacturer's instructions.

Quantitative real-time RT-PCR. Quantitative real-time RT-PCR was performed as previously described [20]. Primer design was based on available sequences in GenBank (NCBI-NLM-PubMed- Gene). The following intron-spanning primers were used, sense and antisense, respectively: MKP3 5'-ATCTCTTTCATAGATGAAGCGCGGG-3' and 5'-TGAGCTTCTGCATGAGGTAGGC-3'; FGF8 5'-CTGATCGGCAA GAGTAACGGC-3' and 5'-CCATGTACCAGCCCTCGTACTTG-3'; RALDH2 5'-CAAGGAGGAGATTTTTGGGCCTGTTC-3' and 5'-GGCCTTGTTTATGTCATTTGTAAAGACAGC-3'; the primers used for the reference gene 18S Ribosomal RNA were: 5'-TGCTCTTAACT GAGTGTCCCGC-3' and 5'-CCCGTTTCCGAAAACCAACAA-3'. Primer sets standard amplification curves were made for MKP3, FGF8, RALDH2 and 18S Ribosomal RNA with cDNA samples from chick limb buds at stage HH24, setting r = 0.99. In all the samples, gene expression was normalized for 18S Ribosomal RNA and for limb region 1 values.

Statistical analysis. All quantitative data are presented as mean \pm SEM. Statistical analysis was performed by one-way ANOVA on ranks and the Student–Newman–Keuls test was used for post-test analysis. Statistical significance was set at $p \leq 0.05$.

Whole-mount in situ hybridization. Embryos and limb buds were collected from stage HH24 and fixed overnight at 4 °C in fresh 4% formaldehyde 2 mM EGTA in phosphate-buffered saline (PBS), rinsed in PBT (PBS, 0.1% Tween 20), dehydrated through a methanol series and stored in 100% methanol at -20 °C. Whole-mount *in situ* hybridization was performed according to the previously described procedure [21].

Immunohistochemistry (IHC). Forelimb buds were collected from stage HH25 chick embryos and immediately fixed in 4% paraformaldehyde in 0.1 M phosphate buffer (PB) with 0.12 mM CaCl₂ and 4% sucrose overnight at 4 °C, washed in buffer only, and then in PB with 15% sucrose, both overnight at 4 °C. Embryos were then incubated in PB with 15% sucrose and 7.5% gelatin for 1 h at 30 °C, immediately frozen in liquid nitrogen chilled isopentane, and stored at -80 °C until sectioned. Approximately 10-µm-thick serial sections were collected on Super Frost slides, and blocked for 30 min with 10% normal goat serum in PBS containing 1% bovine serum albumin. The primary anti-FGF8 antibody (KM1334 from BioWa Inc.; [22,23]) was diluted at 1:500 in the blocking solution and incubated overnight at 4 °C, followed by washing in PBS, and detected with Alexa Fluor 488-conjugated anti-rabbit IgG, all F(ab')2 fragments (Molecular Probes) diluted 1:1000 in blocking solution, followed by PBS washes. Slides were stained with 4',6-diamidino-2-pheny-lindole (DAPI, Sigma), mounted in Vectashield (Vector Laboratories) and sealed after cover-slipping.

Bead implants. A window was cut in fertilised eggs and the vitelline membrane was removed. Heparin acrylic beads (Sigma) soaked in FGF8 protein at 1 mg/ml (R&D Systems) or PBS were microsurgically implanted in the proximal limit of the zone of *mkp3* expression in the right limb buds of stage HH24 embryos. The embryos (n = 12) were incubated for 4 h and hybridized with *mkp3* probe. The same experiment was performed using AG1-X2 ion exchange beads (Bio-Rad) incubated in 4 mg/ml SU5402 (a drug known to specifically block the kinase activity of FGF receptors) (Calbiochem) or in DMSO (n = 6).

Imaging. Embryos processed for *in situ* hybridization were photographed in PBT/0.1% azide, using a Sony DXC-390P-3CCD colour video camera coupled to a Leica MZFLIII microscope. Embryos processed for IHC were photographed using an Olympus DP70 camera coupled to an Olympus BX61 microscope with epifluorescence. Images were treated in Adobe Photoshop CS.

Results and discussion

We performed mkp3 in situ hybridization studies in chick limb buds and observed that mkp3 shows a graded distribution of its transcripts throughout the limb mesenchyme (Fig. 1). *mkp3* expression is stronger in the most distal part of the limb (underneath the AER), fading out in the direction of the more proximal part (Fig. 1A). Realtime polymerase chain reaction (real-time RT-PCR) was performed to characterise the observed mkp3 mRNA gradient in the limb bud. Limb buds at stage HH24 were dissected into three portions of equivalent size along the p-d axis (Fig. 1A). Total mRNA was extracted from multiple batches of each portion of the limb (n = 5) and *mkp3* mRNA content variation in the three regions was analysed by real-time RT-PCR. This high-resolution analysis clearly demonstrates a gradient of *mkp3* transcripts in the limb bud, in a distal to proximal direction (Fig. 1B).

Real-time RT-PCR experiments showed that fgf8 is strongly expressed in the distal tip of the limb bud, and no transcripts can be significantly measured in the more proximal limb regions (Fig. 1B). We could also observe an opposing gradient of raldh2 expression, which presents higher transcript levels in the proximal limb region (Fig. 1C). An antagonism between FGF and RA signalling pathways is thought to operate during the development of several structures (see e.g. [24,25]). Namely, during somitogenesis, FGF8 and RA are mutually inhibitory [26]. In fact, it has been shown that FGFs promote limb distalization by counteracting the RA pathway. This happens through inhibition of the proximalizing RA signal required to maintain the Meis activity, providing a molecular basis for the role of RA in specifying proximal limb fates in different vertebrates [27].

To further characterise limb bud *mkp3* expression gradient, we constructed an intronic *in situ* hybridisation probe

Download English Version:

https://daneshyari.com/en/article/1938412

Download Persian Version:

https://daneshyari.com/article/1938412

Daneshyari.com