



Eel green fluorescent protein is associated with resistance to oxidative stress



Aki Funahashi^a, Masaharu Komatsu^{b,*}, Tatsuhiko Furukawa^{c,d}, Yuki Yoshizono^b, Hikari Yoshizono^b, Yasuhiro Orikawa^b, Shota Takumi^{b,e}, Kazuhiro Shiozaki^b, Seiichi Hayashi^b, Yoshio Kaminishi^a, Takao Itakura^a

^a Laboratory of Marine Biotechnology, Faculty of Fisheries, Kagoshima University, 4-50-20 Shimoarata, Kagoshima 890-0056, Japan

^b Division of Food and Chemical Biology, Faculty of Fisheries, Kagoshima University, 4-50-20 Shimoarata, Kagoshima 890-0056, Japan

^c Department of Molecular Oncology, Graduate School of Medical and Dental Sciences, Kagoshima University, Kagoshima 890-8544, Japan

^d Center for the Research of Advanced Diagnosis and Therapy of Cancer, Graduate School of Medical and Dental Sciences, Kagoshima University, Kagoshima 890-8544, Japan

^e Department of Domestic Science, Kagoshima Women's College, Kagoshima 890-8565, Japan

ARTICLE INFO

Article history:

Received 22 October 2015

Received in revised form 16 December 2015

Accepted 19 December 2015

Available online 30 December 2015

Keywords:

Anguilla japonica

Bilirubin

Eel

eelGFP

Oxidative stress

ABSTRACT

Green fluorescent protein (GFP) from eel (*Anguilla japonica*) muscle (eelGFP) is unique in the vertebrates and requires bilirubin as a ligand to emit fluorescence. This study was performed to clarify the physiological function of the unique GFP. Investigation of susceptibility to oxidative stress was carried out using three types of cell lines including jellyfish (*Aequorea coerulescens*) GFP (jfGFP)-, or eel GFP (eelGFP)-expressing HEK293 cells, and control vector-transfected HEK293 cells. Binding of eelGFP to bilirubin was confirmed by the observation of green fluorescence in HEK293-eelGFP cells. The growth rate was compared with the three types of cells in the presence or absence of phenol red which possessed antioxidant activity. The growth rates of HEK293-CV and HEK293-jfGFP under phenol red-free conditions were reduced to 52 and 31% of those under phenol red. Under the phenol red-free condition, HEK293-eelGFP had a growth rate of approximately 70% of the phenol red-containing condition. The eelGFP-expressing cells were approximately 2-fold resistant to oxidative stress such as H₂O₂ exposure. The fluorescence intensity partially decreased or disappeared after exposure to H₂O₂, and heterogeneous intensity of fluorescence was also observed in isolated eel skeletal muscle cells. These results suggested eelGFP, but not jfGFP, coupled with bilirubin provided the antioxidant activity to the cells as compared to non-bound free bilirubin.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

The 27 kDa green fluorescent protein (GFP) from jellyfish *Aequorea Victoria* (jfGFP), which excitation and emission maxima were 475 and 508 nm, respectively, was discovered and isolated by Shimomura (1979). The jfGFP contains a chromophore induced from the amino acid sequence of -Ser-Tyr-Gly- within the protein structure by cyclization, dehydration, and aerial oxidation (Tsien, 1998; Miyawaki et al., 2003).

We (Hayashi and Toda, 2009) have discovered the green fluorescent protein from eel (*Anguilla japonica*) muscle (eelGFP), with excitation

and emission maxima of 493 and 527 nm, respectively. The deduced amino-acid sequences of eelGFP contained no chromophore such as -Ser-Tyr-Gly-, but exhibited a high similarity to the fatty acid binding protein (FABP) family. Additionally, both eelGFP and FABPs were a lower molecular weight of 14–16 kDa (Hertzel and Bernlohr, 2000; Zimmerman and Veerkamp, 2002; Kumagai et al., 2013) than jfGFP. The FABP family consists of FABP1 - FABP9 which act on different organs and functions (Zimmerman and Veerkamp, 2002). FABPs possess multi-functions including intercellular uptake and transport of fatty acids, regulation of gene transcription and enzyme activity, oxidation of peroxisome and mitochondria (Furuhashi and Hotamisligil, 2008). Thereafter, Kumagai et al. (2013) indicated that *A. japonica* UnaG, which was identical to eelGFP, required hydrophobic antioxidant bilirubin as a ligand with strongly high affinity, K_d = 98 pM, and specificity, to emit green fluorescence. Bilirubin is an endogenous product of heme metabolism in vertebrates. Heme is degraded by heme oxygenase to biliverdin, which is then reduced by biliverdin reductase to form unconjugated bilirubin (Maines, 1988). In mammals, bilirubin forms a complex with albumin, K_d = 87 nM (Kumagai et al., 2013), in blood circulation, and is transported to the liver (Brodersen, 1979). Since bilirubin is a hydrophobic compound, it is conjugated by glucuronyl

Abbreviation: APF, 2-[6-(4'-amino)phenoxy-3 H-xanthen-3-on-9-yl]benzoic acid; eelGFP, eel green fluorescent protein; E-MEM, Eagle's Minimum Essential Medium; FABP, fatty acid binding protein; FBS, fetal bovine serum; hROS, highly oxidative reactive oxygen species; Gly, glycine; jfGFP, jellyfish green fluorescent protein; H₂O₂, hydrogen peroxide; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide; OH, hydroxyl radical; ROS, reactive oxygen species; Ser, serine; Tyr, tyrosine; UnaG, unagi (eel) green fluorescent protein.

* Corresponding author at: Division of Food and Chemical Biology, Faculty of Fisheries, Kagoshima University, Shimoarata 4-50-20, Kagoshima, 890-0056, Japan.

E-mail address: komatsu@fish.kagoshima-u.ac.jp (M. Komatsu).

transferase to a hydrophilic compound, which possess solubility in the bile for excretion (Ostrow and Schmid, 1963; Zucker et al., 1994; Patra and Pai, 1997). The accumulation of excess bilirubin in the circulation after opposite vectorial transport from hepatocytes to the blood causes a jaundice in human, especially in neonates, resulting in basal ganglia with neurologic dysfunction (Dennerly et al., 2001). On the other hand, it is suggested that bilirubin possess a physiologically important role to protect lipids (Frei et al., 1988; Hulea et al., 1995), proteins (Neuzil et al., 1993; Minetti et al., 1998), and nucleotides both intracellularly and extracellularly (Sedlak et al., 2009) against oxidative stress as a potent antioxidant. Interestingly, the antioxidant activity of bilirubin coupled with serum albumin is higher than that of non-bound free style bilirubin (Wu et al., 1991; Kapitulin, 2004). The eelGFP-bound bilirubin is also suggested to possess an antioxidant activity, however, there is no information concerned with the functional properties of eelGFP. Hence, we investigated the effect of the eelGFP coupled with bilirubin on exposure to oxidative stress in the eelGFP-expressing cells.

2. Materials and methods

2.1. Preparation of vector and transfection

Total RNA was extracted from the muscle of eel *Anguilla japonica* and used for cDNA synthesis. The 417-bp target of eelGFP (AB731138) was amplified from cDNA by using the forward primer (5'-AAACTCCGAGATGGTCGAGAAATTTGTT-3') and reverse primer (5'-AAACTCCGAGTCATTCGTCGCCCTCCG-3') (underlined letters show *Xho*I recognition site).

The PCR fragment was ligated into the *Xho*I-digested pcDNA3.1(+) mammalian expression vector (Invitrogen Corp., Carlsbad, CA USA). Human embryonic kidney cells (HEK293) were transfected with eelGFP-pcDNA3.1(+) vector (HEK293-eelGFP) using Lipofectamine™ 2000 (Thermo Fisher Scientific Inc., MA, USA) according to the manufacturer's directions. As a control, HEK293 cells were transfected with empty pcDNA3.1(+) vector (HEK293-CV). In addition, HEK293 cells were transfected with pEGFP-C2 vector (Clontech, Mountain View, CA, USA) (HEK293-jfGFP), and then employed as a further comparison with the functional property of HEK293-eelGFP cells. Stable transfectants were selected and subcloned in the presence of 400 µg/ml G418.

2.2. Cell culture

HEK293-eelGFP, HEK293-jfGFP, and HEK293-CV cells were cultured in Eagle's Minimum Essential Medium (E-MEM) supplemented with 10% fetal calf serum (FCS), 100 units/ml penicillin, 100 µg/ml streptomycin, and 400 µg/ml G418 for 3–5 days under 100% humidity and 5% CO₂ at 37 °C.

2.3. Cell growth rate measurement

Cells were seeded at 2.0×10^5 cells/well (3 mL/well) using 12-well plates, and cultured in Dulbecco's Modified Medium (D-MEM) containing 10% FCS, 100 units/ml penicillin, 100 µg/ml streptomycin, and 6.0 mM L-glutamine with or without phenol red. The number of cells were counted at 0, 24, 48, 72, and 96 h to measure the growth rate of the cells.

2.4. Investigation of sensitivity to oxidative stress

HEK293-eelGFP (1.3×10^4 cells/well), HEK293-jfGFP (1.0×10^4 cells/well), and HEK293-CV (1.0×10^4 cells/well) were seeded into 96-well plates and cultured in D-MEM-10% FCS without phenol red at 37 °C under 5% CO₂ for 48 h. After exposure to H₂O₂ serially diluted from 0 to 200 µM, the cells were cultured at 37 °C under 5% CO₂ for 72 h.

The cell viability was measured to assess the sensitivity of the cells to H₂O₂, and determined by colorimetric assay using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) as described previously (Carmichael et al., 1987). The survival fractions (%) of three independent experiments were calculated from the ratio of cell numbers in each concentration of H₂O₂ to H₂O₂-free controls. The IC₅₀ values were measured as the concentration of H₂O₂ that reduced the number of cells to 50% of that in control medium.

2.5. Isolation of muscle cells and observation of green fluorescence

Muscle cells were isolated by the collagenase digestion method (Rosenblatt et al., 1995; Alam et al., 2004) described in our previous report (Hayashi and Toda, 2009). The green fluorescence of cultured eel muscle cells was then observed using a fluorescence stereomicroscope (MZ10F, Leica, Wetzlar, Germany).

2.6. Conversion of H₂O₂ to the highly oxidative reactive oxygen species (hROS) such as hydroxyl radical

Non fluorescent aminophenyl fluorescein (APF) is converted to the green fluorescent compound fluorescein after reaction with hROS such as hydroxyl radical, but is not detectable with H₂O₂ (Setsukinai et al., 2003). HEK293-CV cells (1.2×10^4 cells/well) were seeded into 96-well plates and cultured for 24 h. Then, cells were exposed to 200 µM H₂O₂ for 4 h in the CO₂ incubator. After exposure to H₂O₂, cells were additionally treated with the hROS specific detection probe APF for 24 h in the CO₂ incubator. We tried to detect the hROS such as hydroxyl radical after conversion mediated by the Fenton reaction (Imlay et al., 1988) from H₂O₂ in the HEK293-CV cells. The green fluorescence of the fluorescein product was observed using EVOS Flويد imaging station (Thermo Fisher Scientific Inc., MA, USA).

3. Results

3.1. Cloning of eelGFP

We have successfully cloned eelGFP mRNA (GenBank accession no. AB731138) from Japanese eel muscle. The open reading frame sequence of eelGFP mRNA was completely identical to that of *UnaG* mRNA (GenBank accession no. AB763906) (Kumagai et al., 2013).

3.2. FCS-dependent expression of green fluorescence of eelGFP

Detection of green fluorescence expression using HEK293-eelGFP cells was carried out after cell culture for 120 h in medium containing




Fig. 1. The relationship between cell density and fluorescent expression in HEK293-eelGFP cells depends on FCS concentration. The FCS concentration (0, 1, 2, 10%)-dependent green fluorescence expression was measured using a Tali Image Cytometer (Thermo Fisher Scientific Inc., MA, USA) after cell culture for 120 h.

Download English Version:

<https://daneshyari.com/en/article/1977128>

Download Persian Version:

<https://daneshyari.com/article/1977128>

[Daneshyari.com](https://daneshyari.com)